
Update in Use of Continuous Perineural Catheters for Postoperative Analgesia

Spencer S. Liu, MD

Recent evidence from randomized controlled trials (RCTs) demonstrates that continuous perineural techniques offer the potential benefits of prolonged analgesia with fewer side effects. This refresher course will summarize pertinent anatomy, technical aspects, and current evidence from RCTs for the indications and efficacy of continuous perineural techniques for postoperative analgesia.

Surgery for Upper Extremities

Evidence for Potential Benefits

RCTs demonstrate that use of continuous interscalene analgesia has superior analgesic efficacy to placebo and intraarticular infusions for hospitalized patients (1–3). Compared with IV patient-controlled analgesia (PCA) for open shoulder surgery, RCTs consistently demonstrate that use of continuous interscalene analgesia not only reduced requirements for postoperative opioids (4,5) but also provided better analgesia, reduced opioid-related side effects, and provided better patient satisfaction for at least the first 48 h after inpatient surgery. Although a case series of 100 patients suggested enhanced physical rehabilitation after shoulder surgery with continuous interscalene analgesia (6), effects on success of physical rehabilitation or duration of hospitalization are unknown. Development of portable disposable and electronic pumps has increased interest in continuous perineural analgesia for outpatient upper extremity surgery (7). RCTs have begun to establish superior efficacy of continuous peripheral catheter techniques for postoperative analgesia after ambulatory surgery. For shoulder surgery, 20 patients were recently randomized to receive continuous interscalene analgesia with either 0.2% ropivacaine or saline for 48 h with a disposable infusion pump at 8 mL/h with patient boluses (2 mL) allowed every 15 min (Fig. 1). Rescue analgesia was provided with oral opioids. During the infusion period, patients receiving ropivacaine had better analgesia, used less oral opioid, had less nausea, sedation, and pruritus, and had better sleep patterns (8). Continuous infraclavicular analgesia for brachial plexus analgesia has

been studied for outpatient upper extremity surgery (9). Thirty patients were randomized to receive either saline or 0.2% ropivacaine with the same disposable pump. Again during the infusion period, patients receiving ropivacaine had better analgesia, used less oral opioid, had less nausea, sedation, and pruritus and better sleep patterns.

Update on Techniques

Recent work in ultrasound imaging with **high-frequency linear arrays** demonstrates clear images of brachial plexus anatomy at the interscalene, supraclavicular, infraclavicular, axillary, and mid-humeral approaches. Direct visualization of neural structure allows visualization of block placement and may improve efficiency of perineural catheter placement (10). For shoulder surgery, the interscalene approach is typically used. However, the classic (Winnie) approach at C6 directs the needle almost perpendicular to the neural bundle. This orientation is satisfactory for single-shot blocks but may increase difficulty of placing a catheter parallel to the neural bundle. Recent modifications to improve the parallel orientation of needle/catheter and neural bundle include the intersternocleidomastoid and modified lateral interscalene approaches. Prospective surveys for both of these techniques suggest satisfactory success rates for catheter placement with the inter-sternocleidomastoid (63 of 70 patients) and modified lateral approach (602 of 700 patients) (11,12).

Another recent technical development is the commercial release of stimulating catheters. Verification of correct catheter placement has been previously reported with fluoroscopy, ultrasound, and computed tomographic (CT) scans. All of these techniques may be cumbersome. The stimulating catheters allow direct and immediate functional confirmation of perineural catheter location and may aid in guidance of catheter placement. Preliminary experience in 64 upper extremity perineural stimulating catheters suggests utility of this technique and also that stimulating characteristics of the catheter are **different** from the needle (**1.6 mA** versus **0.5 mA**) (13).




Figure 1. Use of 0.2% ropivacaine for outpatient continuous interscalene analgesia reduces pain, opioid consumption, side effects, and improves sleep after upper extremity surgery. *different. Values from Ilfeld BM, Morey TE, Wright TW, et al. Continuous interscalene brachial plexus block for postoperative pain control at home: a randomized, double-blinded, placebo-controlled study. *Anesth Analg* 2003;96:275–81.

Risks

Prospective surveys have begun to define potential risks associated with upper extremity perineural catheters. Reports enrolling over 900 patients undergoing continuous interscalene analgesia for 2–5 days observed an approximately 0.7% incidence of catheter site infection and an approximately 0.2% incidence of neurological complications after 6 mo (11).

Lower Extremity Surgery

Prospective clinical trials support the use of continuous femoral analgesia after total knee replacement (TKR) (14–16). Continuous femoral analgesia provides comparable or better analgesia with fewer side effects than IV PCA and epidural analgesia for at least the first 48 h after surgery. The improved analgesia provided by continuous femoral nerve blocks resulted in

faster short-term functional recovery of knee flexion during rehabilitation than IV PCA but without significant differences between the two groups after 6–12 wk (Table 1). Patients undergoing outpatient lower extremity surgery have also been recently studied (17,18). Thirty patients were randomized to receive either saline or 0.2% ropivacaine with the same disposable infusion pump via a popliteal fossa catheter. Again during the infusion period, patients receiving ropivacaine had better analgesia, used less oral opioid, had less nausea, sedation, and pruritus, and had better sleep patterns (Fig. 2). Use of a popliteal fossa catheter may also improve ability to perform outpatient lower extremity surgery. A similar study enrolling 24 patients observed similar benefits and was able to discharge more patients on the same day with 0.25% bupivacaine (40%) versus saline infusions (0%) (19).

Update on Techniques

Use of high-frequency linear arrays is also helping to improve visualization of femoral and sciatic nerves. Case series have described the successful use of ultrasound to guide femoral and popliteal blocks, and direct visualization may also improve catheter placement (20,21). Stimulating catheters have also been used for continuous femoral and sciatic catheters (66 patients) with good success and similar stimulating characteristics as upper extremity placement (13). However, when compared with nonstimulating catheters in patients undergoing continuous femoral analgesia after major knee surgery (TKR and anterior collateral ligament repair), efficacy of stimulating catheters was not different from regular femoral nerve catheters (22). Several recent RCTs have examined different techniques for continuous perineural analgesia for TKR. Use of the posterior psoas compartment technique had been proposed to produce better block of the lumbar plexus than the femoral 3-in-1 approach. However, a RCT examining 3-in-1 (femoral) catheters

Table 1. Continuous Femoral Sheath Analgesia for 48 After Total Knee Replacement Reduces Side Effects and Accelerates Functional Recovery and Patient Discharge from Rehabilitation Center

	IV patient controlled analgesia	Epidural analgesia	Femoral sheath analgesia
Incidence of nausea (%)	21*	5	12
Incidence of dysesthesia (%)	0	5	30*
Time until discharge from rehabilitation center (days)	50 (30–80)*	37 (30–45)	40 (31–60)
Knee flexion (degrees)			
Day 5	60* (50–70)	85 (75–100)	80 (65–85)
Discharge	80* (65–90)	90 (78–100)	90 (70–95)
1 month	90 (85–100)	105 (100–120)	95 (95–100)
3 month	125 (100–125)	130 (115–130)	125 (105–125)

* = different from other two groups ($P < 0.05$). Data from *Anesthesiology* 1999;91:8. Intravenous patient controlled analgesia = morphine 1 mg bolus with 7-min lockout. Epidural analgesia = 1% lidocaine, 2 μ g/mL clonidine, and 0.03 mg/mL morphine at 0.1 mL/kg/h. Femoral analgesia = 1% lidocaine, 2 μ g/mL clonidine, and 0.03 mg/mL morphine at 0.1 mL/kg/h.




Figure 2. Use of 0.2% ropivacaine for outpatient continuous sciatic analgesia reduces pain, opioid consumption, side effects, and improves sleep after lower extremity surgery. *different. Values from Ilfeld BM, Morey TE, Wang RD, Enneking FK. Continuous popliteal sciatic nerve block for postoperative pain control at home: a randomized, double-blinded, placebo-controlled study. *Anesthesiology* 2002;97:959–65.

versus psoas compartment (posterior) approach observed **no differences** in pain scores or analgesic consumption (23). Thus, the techniques appear equivalent with the femoral approach probably being technically easier. Within the femoral approach, a RCT has compared use of a nerve stimulator versus the loss of resistance fascia iliaca technique for placement of non-stimulating catheters. The **fascia iliaca technique was equally effective and required less time** than the nerve stimulator technique (24).

Risks

One prospective survey of 211 femoral catheters noted a **1.4% incidence of infectious** complications and a **0.4% incidence of neurological** complications after 12 mo (25).

Agents for Continuous Perineural Analgesia

Local Anesthetics

There are insufficient data to determine an optimal analgesic solution for the various types of continuous

plexus analgesia. Lidocaine, bupivacaine, and ropivacaine have all been used as the primary local anesthetic for continuous plexus analgesia, with bupivacaine and ropivacaine being the most commonly used agents. The use of bupivacaine (0.1% to 0.25%) typically does not result in toxic blood levels when used for postoperative analgesia for 24–72 h in current regimens (Table 2). Typical **venous** total bupivacaine levels during continuous brachial plexus analgesia are **0.5–1.0 $\mu\text{g}/\text{mL}$** (2) and during continuous lumbar plexus analgesia are 0.5–1.8 $\mu\text{g}/\text{mL}$ (26), whereas levels **greater than 2 $\mu\text{g}/\text{mL}$** are considered toxic. The use of ropivacaine may provide several advantages over bupivacaine and levobupivacaine for providing continuous plexus analgesia. Studies suggest that ropivacaine produces less motor block compared with bupivacaine, which may result in improved participation in postoperative rehabilitation. A comparison of continuous interscalene analgesia with ropivacaine 0.2% versus bupivacaine 0.15% observed equivalent analgesia in both groups, but significantly **less motor block with ropivacaine** (27). The decreased cardiotoxicity of ropivacaine may provide an additional safety margin over both bupivacaine and levobupivacaine. **Animal** studies comparing ropivacaine, levobupivacaine, and bupivacaine suggest **cardiac toxicity** ratios of approximately **1.0:1.7:3.0** (28). Typical perineural infusions of **ropivacaine** 0.2% at 6–12 mL/h for 48 h results in peak plasma levels of approximately **1.7–2.5 $\mu\text{g}/\text{mL}$** , with **toxic** levels considered to be approximately **4 $\mu\text{g}/\text{mL}$** (29). In addition to local anesthetics, analgesic regimens may include clonidine or opioids. However, efficacy of these additives has not been demonstrated (30,31).

Delivery of Continuous Plexus Analgesia

Continuous plexus analgesia may be provided with boluses, continuous infusion, PCA, or a combination of background infusion and PCA boluses. Evidence is accumulating that **patient-controlled** regimens (background infusion plus patient-controlled boluses or patient-controlled boluses only) **may be advantageous for delivery of continuous plexus analgesia**. RCTs indicate that use of a background infusion + PCA provides

Table 2. Typical Delivery Regimens for Continuous Perineural Catheters for Postoperative Analgesia

Catheter	Agent	Infusion rate (mL/h)	PCA bolus (mL)	Lock out (min)
Interscalene	Bupivacaine	5–8	2–4	15–20
Infraclavicular	0.15–0.2% or Ropivacaine 0.2%			
Femoral	Bupivacaine	5–10	*	*
Popliteal	0.1–0.25% or Ropivacaine 0.2%	*	5–10	30–60
		5–8	2	15–20

superior analgesia, reduces local anesthetic consumption, and improves patient satisfaction when compared with infusion or PCA only administration for continuous interscalene, infraclavicular, and popliteal analgesia (32–34). RCTs indicate similar findings in femoral catheters for PCA delivery but do not support the addition of a background infusion for femoral analgesia (35). Typical dosing regimens are listed in Table 2.

References

- Touminen M, Pitkanen M, Rosenberg PH. Postoperative pain relief and bupivacaine plasma concentrations. *Acta Anaesthesiol Scand* 1987;31:276–8.
- Klein SM, Grant SA, Greengrass RA, et al. Interscalene brachial plexus block with a continuous catheter insertion system and a disposable infusion pump. *Anesth Analg* 2000;91:1473–8.
- Delaunay L, Souron V, Lafosse L, et al. Analgesia after arthroscopic rotator cuff repair: subacromial versus interscalene continuous infusion of ropivacaine. *Reg Anesth Pain Med*. 2005;30:117–22.
- Borgeat A, Tewes E, Biasca N, et al. Patient-controlled interscalene analgesia with ropivacaine after major shoulder surgery: PCIA vs. PCA. *Br J Anaesth* 1998;81:603–5.
- Borgeat A, Perschak H, Bird P, et al. Patient-controlled interscalene analgesia with ropivacaine 0.2% vs. patient-controlled intravenous analgesia after major shoulder surgery: effects on diaphragmatic and respiratory function. *Anesthesiology* 2000;92:102–8.
- Cohen NP, Levine WN, Marra G, et al. Indwelling interscalene catheter anesthesia in the surgical management of stiff shoulder: a report of 100 consecutive cases. *J Shoulder Elbow Surg* 2000;9:268–74.
- Ilfeld BM, Enneking FK. Continuous peripheral nerve blocks at home: a review. *Anesth Analg* 2005;100:1822–33.
- Ilfeld BM, Morey TE, Wright TW, et al. Continuous interscalene brachial plexus block for postoperative pain control at home: a randomized, double-blinded, placebo-controlled study. *Anesth Analg* 2003;96:275–81.
- Ilfeld BM, Morey TE, Enneking FK. Continuous infraclavicular perineural infusion with clonidine and ropivacaine compared with ropivacaine alone: a randomized, double-blinded, controlled study. *Anesth Analg*. 2003;97:706–12.
- Chan VW, Perlas A, Rawson R, Odukoya O. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg* 2003;97:1514–7.
- Borgeat A, Dullenkopf A, Ekatothramis G, Nagy L. Evaluation of the lateral modified approach for continuous interscalene block after shoulder surgery. *Anesthesiology* 2003;99:436–42.
- Pham-Dang C, Gunst JP, Gouin F, et al. A novel supraclavicular approach to brachial plexus block. *Anesth Analg* 1999;85:111–6.
- Pham-Dang C, Kick O, Collet T, et al. Continuous peripheral nerve blocks with stimulating catheters. *Reg Anesth Pain Med* 2003;28:83–8.
- Capdevilla X, Barthelet Y, Biboulet P, et al. Effects of perioperative analgesic technique on the surgical outcome and duration of rehabilitation after major knee surgery. *Anesthesiology* 1999;91:8–15.
- Singelyn F, Deyaert M, Pendeville E, et al. Effects of patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous three-in-one block on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty. *Anesth Analg* 1998;87:88–92.
- Chelly JE, Greger J, Gebhard R, et al. Continuous femoral nerve blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *J Arthroplasty* 2001;16:436–45.
- Ilfeld BM, Morey TE, Wang RD, Enneking FK. Continuous popliteal sciatic nerve block for postoperative pain control at home: a randomized, double-blinded, placebo-controlled study. *Anesthesiology* 2002;97:959–65.
- Zaric D, Boysen K, Christiansen J, et al. Continuous popliteal sciatic nerve block for outpatient foot surgery: a randomized, controlled trial. *Acta Anaesthesiol Scand* 2004;48:337–41.
- White PF, Issioui T, Skrivanek GD, et al. The use of a continuous popliteal sciatic nerve block after surgery involving the foot and ankle: does it improve the quality of recovery? *Anesth Analg* 2003;97:1303–9.
- Marhofer P, Schrogendorfer K, Wallner T, et al. Ultrasonographic guidance reduces the amount of local anesthetic for 3-in-1 blocks. *Reg Anesth Pain Med* 1998;23:584–8.
- Sinha A, Chan VW. Ultrasound imaging for popliteal sciatic nerve block. *Reg Anesth Pain Med* 2004;29:130–4.
- Morin AM, Eberhart LH, Behnke HK, et al. Does femoral nerve catheter placement with stimulating catheters improve effective placement? A randomized, controlled, and observer-blinded trial. *Anesth Analg*. 2005;100:1503–10.
- Kaloul I, Guay J, Cote C, Fallaha M. The posterior lumbar plexus (psoas compartment) block and the three-in-one femoral nerve block provide similar postoperative analgesia after total knee replacement. *Can J Anaesth* 2004;5:45–51.
- Morau D, Lopez S, Biboulet P, et al. Comparison of continuous 3-in-1 and fascia iliaca compartment blocks for postoperative analgesia: feasibility, catheter migration, distribution of sensory block, and analgesic efficacy. *Reg Anesth Pain Med* 2003;28:309–14.
- Cuvillon P, Ripart J, Lalourcey L, et al. The continuous femoral nerve block catheter for postoperative analgesia: bacterial colonization, infectious rate, and adverse effects. *Anesth Analg* 2001;93:1045–9.
- Anker-Moller E, Spangsberg N, Dahl JB, et al. Continuous blockade of the lumbar plexus: a comparison of the plasma concentrations and analgesic effect of bupivacaine 0.25% and 0.125%. *Acta Anaesthesiol Scand* 1990;34:468–72.
- Borgeat A, Kalberer F, Jacob H, et al. Patient-controlled interscalene analgesia with ropivacaine versus bupivacaine 0.15% after major open shoulder surgery: the effects on hand motor function. *Anesth Analg* 2001;92:218–23.
- Groban L, Deal DD, Vernon JC, et al. Cardiac resuscitation after incremental overdose with lidocaine, bupivacaine, levobupivacaine, and ropivacaine in anesthetized dogs. *Anesth Analg* 2001;92:37–43.
- Kaloul I, Guay J, Cote C, et al. Ropivacaine plasma concentrations are similar during continuous lumbar plexus blockade using the anterior three-in-one and the posterior psoas compartment techniques. *Can J Anaesth* 2004;51:52–6.
- Ilfeld BM, Morey TE, Enneking FK. Continuous infraclavicular perineural infusion with clonidine and ropivacaine compared with ropivacaine alone: a randomized, double-blinded, controlled study. *Anesth Analg* 2003;97:706–12.
- Ilfeld BM, Morey TE, Thannikary LJ, et al. Clonidine added to a continuous interscalene ropivacaine perineural infusion to improve postoperative analgesia: a randomized, double-blind, controlled study. *Anesth Analg* 2005;100:1172–8.
- Singelyn F, Seguy S, Gouverneur JM. Interscalene brachial plexus analgesia after open shoulder surgery: continuous versus patient-controlled infusion. *Anesth Analg* 1999;89:1216–20.
- Ilfeld BM, Morey TE, Enneking FK. Infraclavicular perineural local anesthetic infusion: a comparison of three dosing regimens for postoperative analgesia. *Anesthesiology* 2004;100:395–402.
- Ilfeld BM, Thannikary LJ, Morey TE, et al. Popliteal sciatic perineural local anesthetic infusion: a comparison of three dosing regimens for postoperative analgesia. *Anesthesiology* 2004;101:970–7.
- Singelyn FJ, Gouverneur JM. Extended “three-in-one” block after total knee arthroplasty: continuous versus patient-controlled techniques. *Anesth Analg* 2000;91:176–80.