



Case Report

Ultrasound-guided serratus plane block for ED rib fracture pain control^{☆,☆☆}



Abstract

Rib fractures are common and painful; providing effective pain relief promotes optimal pulmonary function, thereby preventing complications such as pneumonia and respiratory failure. Opioids, long considered the criterion standard analgesic, have significant drawbacks including respiratory depression, suppression of the cough reflex, and delirium. Regional anesthetic techniques such as intercostal and paravertebral blocks or epidurals can be time-consuming with significant risks and remain uncommon in the typical emergency department. The **serratus anterior plane block (SAPB)** holds promise to be a technically simple, relatively safe, and effective intervention for emergency treatment of rib fracture pain. The technical skill required is on par with an ultrasound-guided femoral nerve block for hip fracture. At its most simple, the block involves using ultrasound to inject a **20- to 40-mL bolus of** local anesthetic into the **space between the surface of the ribs (external intercostal)** and the **overlying muscle (serratus anterior) centered over the rib fractures**. Once injected, the **local first anesthetizes the lateral cutaneous branches** of the intercostal nerves, then as the bellowing motion of the thoracic wall promotes dispersal, it **soaks towards the intercostal space** and the **root intercostal nerves, spreading up and down several thoracic levels** to achieve multilevel analgesia. Further research will clarify questions about the optimal location of SAPB for various rib fracture patterns, the volume and concentration of local anesthetic, the role of injected adjuncts, and expected duration of analgesia. Herein we present the first description of a SAPB successfully used for rib fracture analgesia in the emergency department.

The efficacy of a serratus anterior plane block (SAPB) for rib fracture analgesia is based on the unique anatomy of the intercostal nerves that supply the chest wall from the parietal pleura to the skin [1–4]. The **intercostal nerves** emerge from the **anterior rami** of the **spinal thoracic nerves** to form a **branching network of interconnected nerves** that **communicate across multiple dermatomal levels**. Local freely **travels across the serratus muscle** however, **further study** is needed to determine the **extent of anesthetic spread** from the **superficial extrathoracic space** to the **internal intercostal muscle plane (origin of intercostal nerves)** (Fig. 1). Because of this communication between muscular planes, the SAPB can be performed with injection either **below or beneath the serratus anterior muscle**. The **SAPB** is in the **same family** as the pectoral nerve block, **transabdominus plane**, and **rectus sheath blocks** that **exploit this unique anatomy to produce large areas of thoracic anesthesia with targeted**

thoracic muscular plane injections [2]. These blocks are simple, relatively low risk, and easy to learn for emergency providers already familiar with ultrasound-guided procedures.

Patient 1. An 82-year-old man fell onto a table fracturing right ribs 4–9. He reported severe pain despite intravenous (IV) morphine.

Patient 2. A 65-year-old woman struck by a car while riding a bicycle, resulting in left 5–7 rib fractures. On hospital day 3, she complained of severe pain despite around-the-clock IV hydromorphone. The emergency department (ED) team was consulted by trauma to assist with pain management.

Serratus anterior plane block technique

Informed consent. We explained that the block would last 6 to 8 hours, with the possibility of severe rebound pain, block failure, and local anesthetic toxicity (LAST).

Equipment. Cardiac monitoring, an IV line, and **access to lipid emulsion** was confirmed. A **3.5-in. 20-gauge Touhy needle**, extension tubing, and a 30-mL syringe loaded with 0.5% ropivacaine and 3 mL of 1% lidocaine loaded in a 5-mL syringe with a 27-gauge needle were prepared. An adhesive sterile probe cover, sterile gloves, and sterile drape were assembled.

Positioning. Both patients were placed in the lateral decubitus position (the block can also be performed in the **supine** position). The **injection** was **targeted** toward the **posterior axillary line** at the level **of T5** (Fig. 2).

Injection. The needle was advanced **in-plane**, under ultrasound guidance to the plane **deep** to the serratus anterior muscle (case 1, Fig. 3) and to the plane just **superficial** to the serratus muscle (case 2, Fig. 4). After negative aspiration, a **test dose of 3 mL** confirmed proper needle tip placement with opening of the intramuscular layer. Injection continued under ultrasound guidance with aliquots of 3 to 5 mL after negative aspiration until **30 mL of 0.5% ropivacaine** was administered. Neither patient experienced untoward cardiac events or had signs of LAST; both had appropriate **lung sliding** on ultrasound **postprocedure**. At 30 minutes after the block, patient 1 was able to cough and laugh without pain (dynamic pain 8/10 to 0/10) and did not request analgesics until the following morning (>12 hours after the block). She was discharged the following day. Patient 2 had resting pain of 9/10 in severity. Thirty minutes after SAPB, he spontaneously dressed himself and requested to be discharged home complaining of only minimal pain. He was convinced to accept admission and reported excellent analgesia throughout the night. The next morning (**10 hours after the block**), his pain returned and epidural was placed by the anesthesia service (who are unfamiliar with the SAPB).

Our cases suggest a major role for the SAP in the ED treatment of rib fracture pain. **Rib fractures** are both common (**10%–38% of ED blunt trauma patients**) and painful. Enhancing analgesia with SAP may help improve pulmonary function, avoid adverse outcomes, and reduce mortality [1,3–8]. Current evidence on rib fracture analgesia suggests improved outcomes with an opioid-sparing multimodal approach that integrates regional anesthesia [8].

☆ Prior presentations: None.

☆☆ Funding sources/disclosures: None.




Fig. 1. Innervation of the thorax by the intercostal nerves. A, The intercostal nerves originate from anterior rami of the thoracic spinal nerves and travel adjacent to intercostal artery within the intercostal muscles. The lateral cutaneous branches pierce the internal and external intercostal muscles at the midaxillary line to innervate the muscles and skin of the lateral trunk. When the SAPB is performed, local anesthetic (blue highlight) is deposited directly in contact with the lateral cutaneous branches. The significant relief from fracture pain with the SAPB implies that the local anesthetic soaks into the intercostal space to reach the root intercostal nerves that supply the rib periosteum and parietal pleura. B, Cross-sectional anatomy of the intercostal nerve as it travels from the anterior ramus of the spinal thoracic nerve toward the sternum. Approximate location of local anesthetic (blue highlight) in SAPB is shown. Henry Gray, *Anatomy of the Human Body*, 1918.




Fig. 2. Surface anatomy and patient positioning for the SAPB. A, The serratus anterior muscle originates as muscle branches from the first to ninth ribs (giving the muscle its saw tooth appearance) and inserts onto the medial border of the scapula. In the superficial axillary region, the serratus is found lateral to the pectoralis muscle and medial to the latissimus dorsi muscle. B, Anterior approach. The patient is placed in a supine position with the ultrasound machine positioned for easy line-of-site with an in-plane injection centered over the rib fracture level. C, Posterior approach. The patient is placed in a lateral decubitus position.




Fig. 3. Case 1: **injection deep** to the anterior serratus muscle. A, The patient in left **lateral decubitus position** for an in-plane, **posterior approach** SAPB. B and C, Ultrasound still image showing the space deep to the serratus anterior muscle with bolus local anesthetic injection.

Further study is needed to determine if improved long-term outcomes can be achieved with early initiation of SAPBs for rib fractures in the ED.

We applied the SAPB to emergency rib fracture patients based on the initial description by **Blanco et al [2]** and several additional case reports describing the successful use of the **SAPB for breast surgery [9-13]**, after thoracotomy pain [14-16], and rib fractures [17]. Our initial experience suggests that while the SAPB provides excellent coverage for superficial procedures, it is **not reliable as the sole anesthesia for deeper procedures**

such as **thoracotomy tube placement**. This may be a result of **inconsistent spread** of local anesthetic to the **root intercostal nerves**. The major risk is LAST and can be avoided by using dilute anesthetic (eg, 30 mL of 0.25 bupivacaine) and standard precautions. Pneumothorax would require a major operator error because the **block target** is **superficial to the rib**. Nerve injury is unlikely. Further study is needed to **modify the technique** for specific fracture patterns, such as, **posterior** vs anterior ribs. Injecting in the cephalad-caudal plane or multiple injections at different thoracic




Fig. 4. Case 2: **injection superficial** to the anterior serratus muscle. A, The patient in left lateral decubitus position for an in-plane, posterior approach SAPB with approximate location of ribs shown (Henry Gray, *Anatomy of the Human Body*, 1918). B and C, Ultrasound still image showing the space superficial to the serratus anterior muscle with bolus local anesthetic injection.

levels may promote better spread to multiple thoracic levels. Use of catheters and or adjuvants such as dexamethasone may play a future role in achieving prolonged analgesia. In summary, our preliminary experience suggests the SAPB is technically easy and superior to IV opioids for ED treatment of rib fracture pain.

Edward Durant, MD, MPH
Brittany Dixon, MD
Josh Luftig, PA

Department of Emergency Medicine, Highland Hospital—Alameda Health System, Oakland, CA

Daniel Mantuani, MD
Andrew Herring, MD*

Department of Emergency Medicine, Highland Hospital—Alameda Health System, Oakland, CA

Department of Emergency Medicine

University of California, San Francisco, San Francisco, CA

*Corresponding author. Department of Emergency Medicine
Highland Hospital—Alameda Health System, 1411 East 31st
Street, Oakland, CA 94602-1018

Tel.: +1 510 437 8497; fax: +1 510 437 8322

E-mail address: andrew.a.herring@gmail.com

<http://dx.doi.org/10.1016/j.ajem.2016.07.021>

References

- [1] Ziegler DW, Agarwal NN. The morbidity and mortality of rib fractures. *Trauma Acute Care Surg* 1994;37(6):975–9.
- [2] Blanco R, Parras T, McDonnell JG, Prats-Galino A. Serratus plane block: a novel ultrasound-guided thoracic wall nerve block. *Anaesthesia* 2013;68(11):1107–13.
- [3] Sirmali M, Türüt H, Topçu S, Gülhan E, Yazici U, Kaya S, et al. A comprehensive analysis of traumatic rib fractures: morbidity, mortality, and management. *Eur J Cardiothorac Surg* 2003;24(1):133–6.
- [4] Fligel BT, Luchette FA, Reed RL, Esposito TJ, Davis KA, Santiello JM, et al. Half-a-dozen ribs: the breakpoint for mortality. *Surgery* 2005;138(4):717–23.
- [5] Mackersie RC, Karagianes TG, Hoyt DB, Davis JW. Prospective evaluation of epidural and intravenous administration of administration of fentanyl for pain control and restoration of ventilatory function following multiple rib fractures. *J Trauma* 1991;31:443–9.
- [6] Moon MR, Luchette FA, Gibson SW, Crews J, Sudarshan G, Hurst JM, et al. Prospective, randomized comparison of epidural versus parenteral opioid analgesia in thoracic trauma. *Annals of Surgery* 1999;229(5):684–5.
- [7] Wu CL, Jani ND, Perkins FM, Barquist E. Thoracic epidural analgesia versus intravenous patient-controlled analgesia for the treatment of rib fracture pain after motor vehicle crash. *J Trauma* 1999;47:564–7.
- [8] Bulger EM, Edwards T, Klotz P, Jurkovich GJ. Epidural analgesia improves outcome after multiple rib fractures. *Surgery* 2004;136:426–30.
- [9] Ho AM, Karmaker MK, Critchley LA. Acute pain management of patients with multiple fractured ribs: a focus on regional techniques. *Curr Opin Crit Care* 2011;17(4):323–7.
- [10] Pérez Herrero MA, López Álvarez S, Fadrigue Fuentes A, Manzano Lorefice F, Bartolomé Bartolomé C, González de J. Quality of postoperative recovery after breast surgery. General anaesthesia combined with paravertebral versus serratus-intercostal block. *Rev Esp Anesthesiol Reanim* 2016.
- [11] Takimoto K, Nishijima K, Ono M. Serratus plane block for persistent pain after partial mastectomy and axillary node dissection. *Pain Physician* 2016;19(3):E481–6.
- [12] Bhoi D, Pushparajan HK, Talawar P, Kumar A, Baidya DK. Serratus anterior plane block for breast surgery in a morbidly obese patient. *J Clin Anesth* 2015.
- [13] Bouzinac A, Brenier G, Dao M, Delbos A. Bilateral association of pecs I block and serratus plane block for postoperative analgesia after double modified radical mastectomy. *Minerva Anesthesiol* 2015;81(5):589–90.
- [14] Broseta AM, Errando C, De Andrés J, Díaz-Cambronero O, Ortega-Monzó J. Serratus plane block: the regional analgesia technique for thoracoscopy? *Anaesthesia* 2015;70(11):1329–30.
- [15] Fujiwara S, Komasa N, Minami T. Pectoral nerve blocks and serratus-intercostal plane block for intractable postthoracotomy syndrome. *J Clin Anesth* 2015;27(3):275–6.
- [16] Madabushi R, Tewari S, Gautam SK, Agarwal A, Agarwal A. Serratus anterior plane block: a new analgesic technique for post-thoracotomy pain. *Pain Physician* 2015;18(3):E421–4.
- [17] Kunhabdulla NP, Agarwal A, Gaur A, Gautam SK, Gupta R, Agarwal A. Serratus anterior plane block for multiple rib fractures. *Pain Physician* 2014;17(5):E651–3.