

Mechanisms of action, physiological effects, and complications of hypothermia

Kees H. Polderman, MD

Background: Mild to moderate hypothermia (32–35°C) is the first treatment with proven efficacy for postischemic neurological injury. In recent years important insights have been gained into the mechanisms underlying hypothermia's protective effects; in addition, physiological and pathophysiological changes associated with cooling have become better understood.

Objective: To discuss hypothermia's mechanisms of action, to review (patho)physiological changes associated with cooling, and to discuss potential side effects.

Design: Review article.

Interventions: None.

Main Results: A myriad of destructive processes unfold in injured tissue following ischemia–reperfusion. These include excitotoxicity, neuroinflammation, apoptosis, free radical production, seizure activity, blood–brain barrier disruption, blood vessel leakage, cerebral thermopooling, and numerous others. The severity of this destructive cascade determines whether injured cells will survive or die. Hypothermia can inhibit or mitigate all of these mechanisms, while stimulating protective systems such as early gene activation. Hypothermia is also effective in mitigating intracranial hypertension and reducing brain edema. Side effects include immunosuppression with increased infection risk, cold diuresis and hypovolemia, electrolyte disorders, insulin resis-

tance, impaired drug clearance, and mild coagulopathy. Targeted interventions are required to effectively manage these side effects. Hypothermia does not decrease myocardial contractility or induce hypotension if hypovolemia is corrected, and preliminary evidence suggests that it can be safely used in patients with cardiac shock. Cardiac output will decrease due to hypothermia-induced bradycardia, but given that metabolic rate also decreases the balance between supply and demand, is usually maintained or improved. In contrast to deep hypothermia ($\leq 30^\circ\text{C}$), moderate hypothermia does not induce arrhythmias; indeed, the evidence suggests that arrhythmias can be prevented and/or more easily treated under hypothermic conditions.

Conclusions: Therapeutic hypothermia is a highly promising treatment, but the potential side effects need to be properly managed particularly if prolonged treatment periods are required. Understanding the underlying mechanisms, awareness of physiological changes associated with cooling, and prevention of potential side effects are all key factors for its effective clinical usage. (*Crit Care Med* 2009; 37[Suppl.]:S186–S202)

KEY WORDS: hypothermia; normothermia; fever; mechanisms; neuroprotection; side effects; neurological injury; cardiac arrest; traumatic brain injury

When applying hypothermia in a clinical setting, it is important to be aware of the underlying mechanisms through which the treatment exerts its effects. Understanding these issues can help guide clinical decisions regarding the required depth and duration of the treatment, and help in understanding physiological changes and side effects occurring during mild (34°C–35.9°C), moderate (32°C–33.9°C), mod-

erate-deep (30°C–31.9°C), and deep (<30°C) hypothermia (1). Insufficient understanding of these mechanisms is likely to decrease therapeutic efficacy, and can at worst lead to treatment failure. This is illustrated by early experiences with induced hypothermia in the treatment of cardiac arrest and traumatic brain injury (TBI), and other indications in the 1940s, 1950s, and 1960s (2–15). At that time, it was presumed that hypothermia exerted its effects exclusively through temperature-dependent reductions in metabolism, leading to decreased demand for oxygen and glucose in the brain. Patients were routinely treated with deep hypothermia (<30°C), based on the (erroneous) assumption that this was the sole mechanism of action, and that large reductions in temperature were required to achieve efficacy. In addition, core temperatures that were actually achieved varied considerably both between patients and within the same

patient, because the available cooling and rewarming methods were not very reliable. The most frequently used cooling methods were placement of slabs of ice, ice pads, and refrigerated water on the patient's skin, and sometimes the opening of windows of the hospital wards during the winter months (to the disconcertment of some members of the medical and nursing staff) (11). Because intensive care facilities did not become widely available until the early 1960s, the treatment initially had to be applied in general wards (2–11). Duration of cooling varied considerably, ranging from 2 to 3 days to up to 10 days.

Despite this lack of well-controllable cooling methods and the use of temperatures <30°C (with a much greater risk for severe side effects compared with temperatures $\geq 30^\circ\text{C}$), many of these studies reported benefits compared with “expected outcome” or historical controls (2–11). However, these benefits were

From the Clinical Research, Investigation, and Systems Modeling of Acute Illness (CRISMA) Laboratory, Department of Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh, United States.

The author has not disclosed any potential conflicts of interest.

For information regarding this article, E-mail: k.polderman@tip.nl or poldermankh@ccm.upmc.edu

Copyright © 2009 by the Society of Critical Care Medicine and Lippincott Williams & Wilkins

DOI: 10.1097/CCM.0b013e3181aa5241

variable and uncertain, and as is to be expected there were significant problems in patient management. In the late 1960s, interest among clinicians in therapeutic usage of hypothermia seems to have simply fizzled out; there is a 38-yr gap between the publication of the first two case series describing use of hypothermia in cardiac arrest patients and the next study (8, 9, 16). These experiences illustrate the difficulties that can be encountered during use of controlled hypothermia.

Interest in cooling was rekindled in the early 1980s by the positive results from numerous animal experiments, some of which are discussed below. Many important insights regarding effective use of hypothermia and the mechanisms underlying its protective effects were gained at that time. An important breakthrough was the realization that neurological outcome could be improved by using mild to moderate hypothermia (31°C–35°C) rather than deep hypothermia ($\leq 30^\circ\text{C}$), with far fewer and less severe side effects. The basis for this new insight was the observation that the protective effects of hypothermia were not (at least not exclusively or mainly) due to decreased oxygen and glucose consumption in the brain. Indeed, the mechanisms involved are far more complex.

A cascade of destructive events and processes begins at the cellular level in the minutes to hours following an initial injury. The injury can be ischemic or traumatic. Ischemia may be induced by obstruction of a blood vessel, by pressure/edema, or by other mechanisms. This destructive cascade is similar regardless of the cause, although there are differences in regard to how long the cascade is sustained. These processes—which have been collectively described in terms such as postresuscitation disease, reperfusion injury, and secondary brain injury—may continue for hours to many days after the initial injury and can be retriggered by new episodes of ischemia (17).

A key point is that all of these processes are *temperature dependent*; they are all stimulated by fever, and can all be mitigated or blocked by mild to moderate hypothermia. The wide-ranging effects of hypothermia may explain why it has proved to be clinically effective, whereas studies with agents that affect just one of the destructive processes have been far less successful (18–22).

The second important development was the advent of intensive care units (ICUs) to care for critically ill patients.

The side effects of hypothermia can be far more easily managed in this setting, and the methods for inducing and maintaining hypothermia have improved significantly since the 1950s. These new insights and improvements in facilities have set the stage for a successful comeback of hypothermia as a clinical treatment.

Studies on mechanisms underlying hypothermia's protective effects point to four key factors determining success or failure of cooling treatment. These are:

- *Speed of induction* of hypothermia; outcomes in animal experiments are far better when cooling is initiated rapidly after injury.
- *Duration* of cooling (depending on the severity of the initial injury and the time interval until target temperature is reached).
- *Speed of rewarming* (this should be slow lest the destructive processes be reinitiated; this happens frequently if rewarming speeds are high).
- Proper management and *prevention of side effects*.

The equation is complicated by the fact that the relative contributions of different mechanisms to the ongoing injury may differ. There may be differences between various types of injury (e.g., traumatic vs. purely postischemic), between different patients (depending on genetic factors, comorbidities, gender, etc.), and even within the same patient over time. Thus, the available windows of opportunity for therapeutic interventions such as hypothermia may vary, and the same may apply to the required duration for the cooling treatment to be (fully) effective. For all these reasons, a better understanding of underlying mechanisms may help us to better target our treatments, and help improve outcome.

Many of the available data come from animal experiments; translating these results into clinical practice poses some limitations. Firstly, the relative importance of injury mechanisms may differ between humans and animals, and also between different animal species. Thus, if a specific treatment decreases the degree of histological brain damage in a specific area in rats, this treatment may have similar histological effects in humans but may not make a clinically significant difference in outcome (23, 24). Secondly, some treatments may target specific destructive mechanisms playing a central

role in injury development in a particular animal model, whereas this particular mechanism is far less important in humans (25–27). Such a treatment would be highly effective in that animal model, but would prove far less effective in a clinical trial.

Such differences may result in overestimation of the potential positive effects of hypothermia; the brains of some animals, especially small animals such as rodents, appear to be more responsive to neuroprotection than the human brain (28, 29). On the other hand, protective effects may also be underestimated when looking at only one mechanism and/or only one animal model. For example, many animal studies have focused on the so-called *neuroexcitotoxic cascade*, which is central to the development of postischemic injury in many primitive mammalian animal models (mice, guinea pigs, rats, etc.). This mechanism plays out within relatively short time frames, [± 2 hrs or less] (25–30). However, clinical studies in cardiac arrest patients have reported significant improvements in outcome even when induction of hypothermia was delayed for up to 8 hrs (17); this indicates that other, more protracted mechanisms must play a role in determining outcome in human brain injury. Thus, focusing exclusively on neuroexcitotoxicity would significantly underestimate the available time window for application of hypothermia.

Nevertheless, in spite of such limitations there is much that we can learn from animal experiments, in particular if a treatment has shown value in more than one animal model and/or in primates (orangutans, chimpanzees, etc.). What we have learned regarding ischemia–reperfusion and posttraumatic injuries, and the protective effects of hypothermia, is summarized in Figure 1 and are now discussed in more detail.

Underlying Mechanisms, Physiological Sequelae, and Consequences for Patient Management

Decrease in Cerebral Metabolism. When hypothermia was first used in a clinical setting it was presumed that its protective effects were due purely to a slowing of cerebral metabolism, leading to reduced glucose and oxygen consumption. Indeed this assumption is not completely incorrect: Cerebral metabolism decreases by 6% to 10% for each 1°C




Figure 1. Schematic depiction of the mechanisms underlying the protective effects of mild to moderate hypothermia. *TxA2*, thromboxane A2.

reduction in body temperature during cooling (27, 29, 31–35). Early studies attempted to reduce metabolism as far as possible by using deep hypothermia ($\leq 30^{\circ}\text{C}$). Our current understanding is that a reduction in metabolic rate is only one of many mechanisms underlying hypothermia's protective effects.

Physiological Consequences and Patient Management. When core temperature drops to 32°C , the metabolic rate decreases to 50% to 65% of normal; oxygen consumption and CO_2 production will decrease by the same percentage. If ventilator settings are left unchanged this will lead to hyperventilation, with potentially adverse consequences such as cerebral vasoconstriction (36, 37). The decrease in oxygen consumption will increase oxygen levels in the blood, posing a (theoretical) risk during reperfusion because high oxygen levels may increase reperfusion injury (38). Therefore, ventilator settings should be adjusted as temperature decreases, and frequent checks of blood gasses are necessary especially in the induction phase. During maintenance the feeding rate should also be adjusted to reflect lower metabolic demands.

Other metabolic changes include increased fat metabolism, leading to an increase in the levels of glycerol, free fatty acids, ketonic acids, and lactate. This in turn can cause mild metabolic acidosis.

This phenomenon does not require treatment; pH rarely decreases below 7.25, and the drop in extracellular pH is not reflected in intracellular pH, which increases slightly under hypothermic conditions (33).

Another consequence of hypothermia is decreased insulin secretion and, in many patients, a moderate (and sometimes severe) insulin resistance. This can lead to hyperglycemia and/or a significant increase in doses of insulin required to maintain glucose levels within an acceptable range. This temperature-dependence of insulin requirements is particularly important in the rewarming phase, when insulin sensitivity may increase rapidly, leading to hypoglycemia if insulin doses are not decreased. Thus, glucose levels should be checked frequently particularly during rewarming and, to a lesser extent, the induction phase of hypothermia (1). For this and other reasons, rewarming rates following hypothermia treatment should be slow and controlled. In our unit the target rewarming rate is $0.25^{\circ}\text{C}/\text{hr}$ for postcardiac arrest patients and $0.1^{\circ}\text{C}/\text{hr}$ for patients following severe TBI.

Apoptosis, Calpain-Mediated Proteolysis and Mitochondrial Dysfunction. Following a period of ischemia and reperfusion, cells can become necrotic, fully or partially recover, or enter a path leading to so-called apoptosis or programmed cell

death. Whether apoptosis will develop is determined by cellular processes such as mitochondrial dysfunction, other disorders in cellular energy metabolism, and release of so-called caspase enzymes. Numerous studies have shown that hypothermia can interrupt the apoptotic pathway, thereby preventing cellular injury from leading to apoptosis (39–43). Hypothermia seems to affect mainly the early stages of the apoptotic process and apoptosis initiation (40). Its effects include inhibition of caspase enzyme activation (39–41, 43, 44), prevention of mitochondrial dysfunction (42), decreased overload of excitatory neurotransmitters, and modification of intracellular ion concentrations. (The latter mechanisms are discussed in more detail below.)

Apoptosis begins relatively late in the postperfusion and/or posttrauma phase, while continuing for a period of 48 to 72 hrs or even longer (43–45). Thus, apoptosis is one of the mechanisms that can be mitigated (and perhaps even prevented) for some time after injury, providing (at least in theory) a wide window of opportunity for therapeutic interventions such as hypothermia. For this and other reasons, influencing apoptotic pathways could play an important role in providing neuroprotection and in mitigating postresuscitation injury in humans.

Ion Pumps and Neuroexcitotoxicity. A large body of evidence suggests that hypothermia inhibits harmful excitatory processes occurring in brain cells during ischemia–reperfusion. Levels of high-energy metabolites such as adenosine triphosphate and phosphocreatine decrease within seconds when oxygen supply to the brain is interrupted (27). The breakdown of adenosine triphosphate and the switch of intracellular metabolism to anaerobic glycolysis lead to an increase in intracellular levels of inorganic phosphate, lactate, and H^+ , resulting in both intra- and extracellular acidosis and an influx of calcium (Ca^{2+}) into the cell. Loss of adenosine triphosphate and acidosis also inhibits the mechanisms that normally deal with excessive intracellular Ca^{2+} by sequestering Ca^{2+} from the cell, further aggravating intracellular Ca^{2+} overload. These problems are compounded by failure of adenosine triphosphate-dependent $\text{Na}^+\text{-K}^+$ pumps and K^+ , Na^+ , and Ca^{2+} channels, leading to an additional influx of Ca^{2+} (46). The excess Ca^{2+} induces mitochondrial dysfunction (increasing intracellular calcium influx yet further, in a vicious cycle) and

activates numerous intracellular enzyme systems (kinases and proteases). In addition, immediate early genes are activated and a depolarization of neuronal cell membranes occurs, with a release of large amounts of the excitatory neurotransmitter glutamate into the extracellular space (25). This leads to prolonged and excessive activation of membrane glutamate receptors, further stimulating Ca^{2+} influx through activation of Ca^{2+} channels in another vicious cycle. Under normal circumstances, neurons are exposed to only very brief pulses of glutamate; prolonged glutamate exposure induces a permanent state of hyperexcitability in the neurons (the *excitotoxic cascade*), which can lead to additional injury and cell death. Furthermore, high levels of glutamate can be neurotoxic, especially in energy-deprived cells. Glutamate receptor activation can persist for some time after reperfusion, even when glutamate levels have returned to normal; this may be another important mediator of brain cell death.

In summary, ischemia and reperfusion lead to an interruption of a delicate balance between calcium influx and sequestration at the cellular level. Numerous animal experiments have clearly demonstrated that key destructive processes of the neuroexcitatory cascade (such as calcium influx, accumulation of glutamate, and the release of its coagonist glycine) can be prevented, interrupted, or mitigated by hypothermia (23, 25, 45–53). Even a relatively small decrease in temperature can significantly improve ion homeostasis, whereas the occurrence of fever can trigger and stimulate these destructive processes.

It is unclear how long the window of opportunity to interrupt this cascade is. Disruptions in Ca^{2+} homeostasis begin in the minutes after injury but may continue for many hours (sometimes even days). Furthermore, the mechanism can be reinitiated by new episodes of ischemia. Thus, in theory this mechanism may be susceptible to therapeutic interventions provided these are initiated in the hours following injury. However, some animal experiments suggest that neuroexcitotoxicity can be blocked or reversed only if the treatment is initiated in the very early stages of the neuroexcitatory cascade (26–27, 30, 54). Other studies have reported somewhat wider time frames, ranging from 30 mins to up to 6 hrs (55–67). Thus, the period during which the neuroexcitatory cascade can be modified *in vivo* remains unclear; indeed,

this may vary between different species and different types of injury. Some authors have suggested that the therapeutic window could be extended by combining hypothermia with other, more specific treatments such as caspase inhibitors or other experimental compounds (41, 68).

Immune Response and Inflammation. In most types of brain injury, a significant and protracted inflammatory response begins about 1 hr following a period of ischemia–reperfusion. Proinflammatory mediators such as tumor necrosis factor- α and interleukin-1 are released in large quantities by astrocytes, microglia, and endothelial cells; this rise begins ± 1 hr after injury and they remain elevated for up to 5 days (27, 69). This stimulates the chemotaxis of activated leukocytes across the blood–brain barrier, leading to an accumulation of inflammatory cells in the injured brain, as well as the appearance of adhesion molecules on leukocytes and endothelial cells. Simultaneously, there is an activation of the complement system which begins in the very early stages after brain injury and further stimulates the passage of neutrophils and (in later stages) monocytes–macrophages (69). These inflammatory and immunologic responses occur especially during reperfusion and are accompanied by free radical production. This can cause significant (additional) injury through the phagocytic actions of macrophages, synthesis of toxic products, and further stimulation of immune reactions in a vicious cycle.

However, it should be realized that this inflammatory response is to some extent physiological, and may have a dual role in the sense that some inflammatory mediators have neuroprotective properties, whereas others are neurotoxic (69–72). Nevertheless, there is strong evidence suggesting that certainly a disproportionate and persistent production of cytokines and leukocyte infiltration can significantly increase the risk and extent of brain cell injury and infarction (68–74). Especially the interleukin-1 family appears to be important in this regard (74). Of note, this effect is to some extent time dependent, with the destructive aspects of inflammation outweighing the potential benefits especially in the later stages of injury (69–72). Thus, again, there may be a potential time window for therapeutic interventions to interrupt or mitigate this process before it becomes destructive.

Numerous animal experiments and some clinical studies have shown that hypothermia suppresses ischemia-in-

duced inflammatory reactions and release of proinflammatory cytokines (75–78). Hypothermia also prevents or mitigates reperfusion-related DNA injury, lipid peroxidation, and leukotriene production (47–49), and it decreases the production of nitric oxide, which is a key agent in the development of postischemic brain injury (25). In addition, hypothermia can impair neutrophil and macrophage function, and (at temperatures $<32^{\circ}\text{C}$ – 33°C) decrease white blood cell count.

In animal experiments, the extent of brain injury and infarct size can be significantly attenuated if any of these processes is mitigated or interrupted (25); given that hypothermia can affect all of these steps there is (at least in theory) a huge potential for improving outcome. Furthermore, given that the inflammatory response begins relatively late (>1 hr following ischemia–reperfusion), and given that the destructive processes take some time to develop fully and continue for prolonged periods of time, there appears to be a clear therapeutic window for use of hypothermia to favorably affect this specific mechanism.

Free Radical Production. A destructive process that is closely linked to but distinct from the mechanisms discussed above is the release of free oxygen radicals following ischemia–reperfusion. Mediators such as superoxide (O_2^-), peroxynitrite (NO_2^-), hydrogen peroxide (H_2O_2), and hydroxyl radicals (OH^-) play an important role in determining whether injured cells will recover or die (47, 50, 79, 80). Free radicals can oxidize and damage numerous cellular components. Although brain cells have various enzymatic and nonenzymatic antioxidant mechanisms that prevent this type of injury under normal circumstances, free radical production following ischemia–reperfusion is so great that these defensive mechanisms are likely to be overwhelmed, leading to peroxidation of lipids, proteins, and nucleic acids.

Under hypothermic conditions the quantity of free radicals that is generated is significantly reduced, although free radical production is not completely prevented (47, 50, 79, 80). This allows the endogenous antioxidative (protective) mechanisms to better cope with free radicals that are being released, thereby preventing or significantly mitigating oxidative damage. This allows the cell to repair itself and recover, rather than suffering permanent damage and/or dying. The degree of inhibition of free radical produc-




Figure 2. Schematic depiction of the potential role of intracranial pressure as both a marker of ongoing brain injury and a potential cause of additional injury, and of the factors that may contribute to a rise in intracranial pressure. CSF, cerebrospinal fluid.

tion appears to be more or less linearly linked to the temperature; i.e., the lower the temperature, the lower the amount of free radicals (79).

Vascular Permeability, Blood–Brain Barrier Disruption, and Edema Formation. Ischemia–reperfusion and/or traumatic injury can lead to significant disruptions in the blood–brain barrier, which can facilitate the subsequent development of brain edema (81–83). Therapeutic interventions such as mannitol administration in TBI or stroke can add to blood–brain barrier disruption (82). Mild hypothermia significantly reduces blood–brain barrier disruptions (81–83), and also decreases vascular permeability following ischemia–reperfusion, further decreasing edema formation (84). The concept of a membrane- and blood–brain barrier–stabilizing effect of hypothermia is supported by the observation that hypothermia decreases extravasation of hemoglobin following TBI (85). Ischemia–reperfusion affects blood–brain barrier integrity in other ways, including decreased fluidity and integrity of cell membranes and increased vascular permeability of microvascular endothelial cells in the brain, mediated by vascular endothelial growth factor *via* release of nitric oxide (86, 87).

These processes of membrane disintegration and hypoxia-induced leakage can be mitigated or reversed by hypothermia (87). Hypothermia also decreases cytotoxic edema *via* the mechanisms de-

scribed above (dampening inflammatory responses and free radical production, improving ion homeostasis, etc.), and through other mechanisms, which are summarized in Figure 2. Brain edema can be monitored by measuring intracranial pressure (ICP), which can be regarded as a final common pathway for the destructive processes leading to brain edema (88) [Fig. 2].

The key role of brain edema and intracranial hypertension as a cause of (additional) neurological injury in severe TBI and ischemic stroke is well recognized (89, 90). Brain edema can also increase brain injury in acute encephalitis and meningitis (91), and some evidence suggests that edema-related injury can play a role in some patients with posthypoxic injury following cardiac arrest (92).

As intracranial hypertension appears to be both a marker for ongoing neurological injury and a cause of additional injury (88), it seems plausible that treatments that reduce ICP will also help improve neurological outcome. ICP is frequently used as a parameter to guide treatments and assess short-term therapeutic efficacy in patients with TBI (89, 93). Hypothermia has been used to treat brain edema and reduce ICP in a wide range of neurological injuries, including TBI, ischemic stroke, hepatic encephalopathy, meningitis, encephalitis, and subarachnoid hemorrhage (17, 28).

In theory, this mechanism of action would offer a relatively wide therapeutic

window to mitigate neurological injuries. Brain edema may develop rapidly, but more typically arises several hours after injury, often peaking after 24 to 72 hrs (94). It should be noted that although hypothermia has been shown to decrease ICP in many clinical studies, the results in improving survival and neurological outcomes have been mixed (17, 88), with one large trial failing to show any benefits during hypothermia treatment in spite of hypothermia-induced decreases in ICP (95). These varying results may be (partly) linked to the management of hypothermia's side effects (17, 88); these issues are discussed more extensively below. As explained above, the speed of induction, duration and depth of hypothermia, and rewarming speeds all play important roles in determining whether or not lasting protective effects are achieved (1, 17, 88, 96, 97).

Intra- and Extracellular Acidosis and Cellular Metabolism. The diminished integrity of cell membranes, the failure of various ion pumps, development of mitochondrial dysfunction, inappropriate activation of numerous enzyme systems with cellular hyperactivity, and the disruption of various other intracellular processes all contribute to the development of *intracellular acidosis*, a factor that powerfully stimulates many of the above-mentioned destructive processes (98, 99). Ischemia–reperfusion also leads to substantial rises in cerebral lactate levels (100–102). All of these factors can be significantly attenuated by hypothermia (98–103).

In addition, *brain glucose utilization* is affected by ischemia–reperfusion, and there is evidence suggesting that hypothermia can improve brain glucose metabolism; i.e., the ability of the brain to utilize glucose (104, 105). Similar observations have been made in severe TBI, for which animal experiments and some clinical studies suggest that an initial increase in cerebral glucose metabolism in the hours following trauma is followed by a deep and persistent decrease in metabolic rate, with a depression of mitochondrial oxidative phosphorylation and glucose utilization that can last for several weeks (106–108). Induced hypothermia applied during or after reperfusion increases the speed of metabolic recovery, with a better preservation of high-energy phosphates and reduced accumulation of toxic metabolites (106–108).

Brain Temperature and Cerebral Thermopooling. Even in healthy individ-

uals, the temperature of the brain is slightly higher than the measured core temperature (109–111). This difference can increase significantly in patients following neurological injury, with gradients ranging from 0.1°C to more than 2°C (112–118), although this phenomenon does not occur in all brain-injured patients (118). These differences may increase even further when a patient develops fever, a very frequent phenomenon in patients with neurological injuries (119–123). In addition, there may be small temperature differences between different areas of the brain even in healthy individuals, with active areas having a slightly higher temperature (124–127). These differences increase significantly when brain injury occurs, with injured areas becoming more hyperthermic than noninjured areas (128). The reason for this is that some of the destructive mechanisms discussed above lead to excessive generation of heat. These include excitotoxicity (which, as described above, induces “cellular hyperactivity” with extreme activation of enzyme systems, with concomitant generation of heat), the neuroinflammatory response, free radical production, and a process known as “cerebro thermopooling” (128, 129), meaning that excess heat generated in injured areas of the brain becomes more difficult to remove *via* the normal heat dissipation mechanisms (lymph and venous drainage) due to development of local brain edema. The heat thus becomes trapped in injured areas, further adding to hyperthermia-related injury.

Thus, a vicious cycle can arise: Brain injury leads to general overheating of the brain; more overheating occurs especially in injured areas; this leads to local and sometimes general brain edema, which then makes it more difficult to remove the excess heat. Due to this chain of events, brain temperatures in injured brain areas may exceed measured core temperatures by as much as 2°C to 4°C.

This is relevant because there is strong evidence from animal studies that (external) induction of hyperthermia significantly increases the risk and extent of neurological injury (17, 130–136). Hyperthermia increases the likelihood that ischemic areas become apoptotic or necrotic, even when fever develops (or is externally induced) some time after the initial injury. Baena et al (131) reported a 2.6-fold increase in neuronal injury in the rat hippocampus if mild warming

(1°C–2°C above normal for 3 hrs) was induced 24 hrs after a brief episode of forebrain ischemia. This suggests that fever can be detrimental even when it is of short duration and occurs on the day after injury. Others have reported similar observations in different animal models (132–136). The effects of fever appear to be more pronounced when they coincide with an episode of cerebral ischemia, suggesting that ischemic brain cells become even more susceptible to the harmful effects of hyperthermia (135–136).

Numerous clinical studies have confirmed that fever is indeed an independent predictor of adverse outcome in stroke, TBI, and postanoxic injury following cardiac arrest (121–123, 137–142). Schwarz et al (121) found that even very mild hyperthermia (>37.5°C!) within the first 72 hrs was independently associated with poor outcome in patients with intracerebral hemorrhage, and that there was a linear relationship between the severity and duration of fever and the risk for adverse outcome. In patients with ischemic stroke, fever is associated with a 3.4-fold increase in risk for adverse outcome (140), a higher brain infarct volume (141), and increased mortality (142). Although it has not been conclusively proven that these relationships are causal (i.e., that fever increases injury rather than just being a marker), this is strongly suggested by the temporal relationship and the results from animal studies (130–136).

Coagulation Activation and Formation of Microthrombi. Cardiopulmonary arrest and resuscitation are accompanied by a marked activation of coagulation, which can lead to intravascular fibrin formation with blockage of the microcirculation in the brain and heart (143–146). Administration of anticoagulants such as heparin and recombinant tissue-type plasminogen activator improves microcirculatory reperfusion and survival in animal experiments (146, 147); in addition, thrombolysis can improve cerebral tolerance to ischemia (148). Initial clinical reports suggesting that administration of thrombolytic agents could improve neurological outcome and survival in cardiac arrest (149) have not been confirmed in larger studies (150). Nevertheless, activation of coagulation seems to play an important role in developing ischemia–reperfusion injury, and although reversing this mechanism by itself may not improve outcome, this may be different if additional mechanisms are targeted simultaneously.

Hypothermia has some anticoagulatory effects. Mild platelet dysfunction occurs at temperatures $\leq 35^\circ\text{C}$, and some inhibition of the coagulation cascade develops at temperatures $\leq 33^\circ\text{C}$; platelet count can also decrease during cooling (1, 151–154). In theory, this anticoagulation effect may constitute yet another neuroprotective mechanism. This remains speculative given that no studies directly addressing this issue have been performed.

Vasoactive Mediators. Several studies have shown that hypothermia affects the local secretion of vasoactive substances such as endothelin, thromboxane A₂ (TxA₂), and prostaglandin I₂ in the brain and at other sites. Endothelin and TxA₂ are powerful vasoconstrictive agents, whereas prostaglandin I₂ is a vasodilator. TxA₂ also stimulates platelet aggregation (155–157). TxA₂ and prostaglandin I₂ play an important role in regulating local cerebral blood flow, and a balance between the two is required to maintain homeostasis (155). This local homeostasis can be disrupted following an ischemic or traumatic event, with a relative increase in production of TxA₂ (158–161). This disruption in equilibrium can lead to vasoconstriction, hypoperfusion, and thrombogenesis in injured areas of the brain.

Animal experiments and small clinical studies suggest that this relative predominance of local vasoconstrictors can be corrected or modified by hypothermia (49, 162, 163). Aibiki et al (162) reported that moderate hypothermia (32°C–33°C) led to a reduction in prostanoid production and attenuation of the imbalance between TxA₂ and prostaglandin I₂ in patients with severe TBI. Chen et al (163) found decreased production of the vasoconstrictor endothelin-1 during induction of hypothermia.

This and other preliminary evidence suggests that hypothermia may favorably affect vasoactive mediator secretion in brain-injured patients. However, regulation of cerebral perfusion is a highly complex issue, especially in the injured brain; local circulation can be influenced by the presence or absence of cerebral autoregulation, ventilator settings and blood gas management, systemic blood pressure, fluid management including osmotic therapy, etc. Thus, the impact of hypothermia on this equation needs to be studied in greater detail.

Improved Tolerance for Ischemia. Hypothermia improves the tolerance for

The three phases of hypothermia treatment




Figure 3. Graphic depiction of the three phases of hypothermia treatment. The induction phase should last between 30 and 120 mins; rapid cooling may lead to a small overshoot, which should be accepted provided it is no greater than 1°C. The maintenance phase usually lasts 24 hrs in cardiac arrest patients (may be longer for other indications) and should be characterized by no or minimal fluctuations in temperature. The rewarming phase should be slow and controlled, with rewarming rates of 0.2°C to 0.5°C in cardiac arrest patients and lower rewarming rates for other indications. Fever should be prevented after rewarming.

ischemia in various animal models (164–166). For this reason, it is widely used in the perioperative setting, especially in major vascular surgery, cardiothoracic surgery, and neurosurgical interventions (17). An ability to better withstand periods of ischemia would be an important protective mechanism, because many types of neurological injury can be complicated by ischemic events in the days following the initial insult.

For example, in patients with TBI a combination of biological factors (cytotoxic and vasogenic edema) and mechanical factors (blockage of spinal fluid drainage) can lead to cerebral edema and intracranial hypertension in the hours and days following initial injury (Fig. 2); cerebral edema can then cause additional ischemic injury. In patients with subarachnoid hemorrhage the development of vasospasms in the days and weeks following a bleeding episode can lead to significant ischemic injury. The risk for ischemic injury occurring in the postadmission period may also apply to postcardiac arrest patients, in whom cerebral ischemia may persist for several hours following successful resuscitation even when arterial oxygen levels are normal (167). Thus, a hypothermia-induced increase in tolerance for ischemia could be another powerful protective mechanism.

Suppression of Epileptic Activity. Nonconvulsive status epilepticus (i.e., epileptic activity without obvious clinical signs and symptoms) occurs frequently in

patients with postanoxic encephalopathy, stroke, TBI, and subarachnoid hemorrhage (168–172). It remains uncertain whether nonconvulsive status is a direct cause of additional injury; however, mounting evidence suggests that brain injury can be significantly increased when nonconvulsive status occurs in the acute phase of brain injury; i.e., while the destructive processes outlined above are ongoing (168, 169). In other words, although the role of nonconvulsive status by itself has not been fully clarified, when it occurs in combination with ongoing brain injury the two are synergistically detrimental (168, 169).

Evidence from various sources shows that hypothermia can suppress epileptic activity. Small case series report successful use of cooling to treat grand mal seizures (173–175). Animal studies have shown that external warming increases the extent of epilepsy-induced brain injury, that prevention or prompt treatment of fever reduces these injuries, and that induction of hypothermia further decreases seizure-induced brain injury (176–179). This antiepileptic effect offers yet another pathway through which hypothermia could provide neuroprotection, and by which fever may increase neurological injury.

Spreading Depressionlike Depolarizations. Some animal studies have suggested that neuronal damage in various types of neurological injury can be significantly increased by so-called *spreading*

depressionlike depolarizations (23, 179). This has been studied mainly in models for TBI and stroke; its potential role in the relatively brief global ischemic brain injury that characterizes cardiac arrest is unclear. Hypothermia can suppress spreading depressions in different types of neurological injury (23, 180–182), providing yet another mechanism by which hypothermia could improve outcome.

Influence on Genetic Expression. Hypothermia increases the expression of so-called *immediate early genes*, which are a part of the protective cellular stress response to injury, and stimulates the induction of *cold shock proteins*, which can protect the cell from ischemic and traumatic injury (97). This provides yet another mechanism through which hypothermia can provide neuroprotection.

Thus, hypothermia favorably affects a whole range of destructive mechanisms unfolding in the injured brain after ischemia or trauma. This wide range of effects probably explains its efficacy, in contrast to interventions that target just one or two of these mechanisms and have mostly proved unsuccessful in clinical trials. Different mechanisms may play a greater or smaller role in different types of brain injury, and/or during different phases of ongoing brain injury. A more detailed understanding of these processes will help us apply therapeutic hypothermia and controlled normothermia more effectively.

The other key element for successful use of therapeutic cooling is awareness and proper management of the physiological consequences and side effects. These issues are dealt with in the next section.

Physiological Aspects of Cooling

Phases of Cooling Treatment. A hypothermia treatment cycle can be divided into three distinct phases (Fig. 3):

- The induction phase, when the aim is to get the temperature below 34°C and then down to the target temperature as quickly as possible;
- The maintenance phase, when the aim is to tightly control core temperature, with minor or no fluctuations (maximum, 0.2°C–0.5°C);
- The rewarming phase, with slow and controlled warming (target rate, 0.2°C–0.5°C/hr for cardiac arrest patients and 0.1°C–0.2°C/hr for other

categories such as patients with severe TBI) (183).

Each of these phases has specific management problems. The risk for immediate side effects such as hypovolemia, electrolyte disorders, and hyperglycemia is greatest in the induction phase (17, 184, 185). This phase presents the greatest patient management problems, requiring frequent adjustments in ventilator settings; dosing of sedation, insulin, and vasoactive drugs; and fluid and electrolyte administration (185). The risks can be reduced by rapid induction of hypothermia; i.e., minimizing the duration of the induction phase and reaching the more stable maintenance phase as quickly as possible. Rapid cooling can be achieved by combining different cooling methods, for example, surface cooling with infusion of 1500–3000 mL of cold (4°C) fluids using a pressure bag (186). Furthermore, some of the new intravascular devices may have more rapid cooling rates than previously used methods (1).

The maintenance phase is characterized by increased stability of the patient, with a decreased shivering response and less risk for hypovolemia and electrolyte loss. In this phase, attention should shift toward prevention of longer-term side effects such as nosocomial infections and bedsores.

In the rewarming phase, the patient's temperature should be increased very slowly, for a number of reasons. Firstly, rapid rewarming can cause electrolyte disorders (in particular, hyperkalemia) caused by shifts from the intracellular to the extracellular compartment. This can be largely prevented by slow and controlled rewarming. Secondly, insulin sensitivity can increase during rewarming, and slow rewarming decreases the risk for hypoglycemia if the patient is being treated with insulin. Thirdly, some animal experiments and clinical observations suggest that rapid rewarming could lead to loss of some or even all of the protective effects of hypothermia (187–191). Significant decreases in jugular venous oxygen saturation have been reported during rapid rewarming of patients following cardiac surgery under hypothermic conditions, indicating hypoxia of the brain (190); more slow rewarming led to a decrease in the incidence and severity of jugular bulb desaturations (190). In another study, Bissonnette et al (191) reported that patients who were rapidly rewarmed following cardiopulmonary bypass surgery often developed severe brain hyperthermia, even when core temperature mea-

sured at other sites remained normal. Although none of these experiments directly addressed optimal rewarming rates for cardiac arrest patients treated with hypothermia, these and other findings indicate that rapid rewarming can increase ischemia and aggravate destructive processes in the injured brain (1).

Another important concept is the maintenance of strict normothermia following the rewarming phase. As discussed above, fever is independently linked to adverse outcome in all types of neurological injury, including postanoxic injury following cardiac arrest (17). In addition, cerebrovascular reactivity may be impaired following hypothermia treatment (192), increasing the potential harmful effects of fever.

Physiological Changes and Side Effects of Cooling

The induction of mild hypothermia induces numerous changes throughout the body. The most important physiological changes and side effects, and their consequences for patient management, are discussed below.

Shivering and Cutaneous Vasoconstriction. A separate article in this supplement is devoted to shivering (193); therefore, this topic is discussed only briefly here.

In awake patients, shivering induces unfavorable effects such as increased oxygen consumption and metabolic rate, excess work of breathing, and increased heart rate with increased myocardial oxygen consumption (1, 184, 194–197). In the perioperative setting, hypothermia has been linked to an increased risk for morbid cardiac events, particularly in older patients with heart disease (194–196). However, these adverse consequences are linked to the hemodynamic and respiratory responses rather than to shivering *per se*; these responses can be largely suppressed through administration of sedatives, anesthetics, opiates, or other drugs during therapeutic cooling (1). For example, in contrast to the tachycardia associated with perioperative (accidental) hypothermia, heart rates are markedly reduced during therapeutic cooling (1). Nevertheless, it is important to prevent or aggressively treat shivering because it significantly complicates hypothermia induction, and leads to an undesirable increase in metabolic rate and oxygen consumption (1). Appropriate sedation and analgesia will also cause va-

sodilation, which can facilitate heat loss through surface cooling.

Relatively effective and rapidly acting antishivering agents include fentanyl, alfentanil, meperidine, dexmedetomidine, propofol, clonidine, and magnesium (1). Specific advantages and disadvantages of these and other agents are reviewed in more detail elsewhere (1). Brief-acting paralyzing agents can be useful in the induction phase, especially when hypothermia is initiated outside the ICU setting, but prolonged paralysis is usually unnecessary (1). The effectiveness of the shivering response decreases with age; therefore, the doses of drugs required to suppress shivering are usually lower in older patients. Warm-air skin counterwarming can be used as an accessory method to lower the shivering threshold, and to decrease drug doses required to prevent shivering (198, 199).

In our unit, the standard approach is to administer a loading dose of magnesium (30 mmol) and fentanyl (50–100 µg) when cooling is initiated, together with continuous infusion of either midazolam or propofol (the latter only in hemodynamically stable patients). If shivering occurs, another bolus of fentanyl (50–150 µg) is given, and more magnesium if serum Mg^{2+} is ≤ 2.0 mmol/L. In some patients, the dose of sedatives will be temporarily increased during induction and/or a bolus dose of midazolam (5–10 mg) will be given. If shivering persists, one of the following drugs is given, depending on the clinical circumstances: clonidine, meperidine, ketanserin, or (in rare cases) a brief-acting paralyzing agent. All patients should be carefully monitored for shivering during all phases of hypothermia treatment. Use of a recently developed shivering assessment scale (200) may be considered for this purpose.

The importance of adequate sedation and suppression of hypothermia-induced stress responses is underscored by observations from animal experiments suggesting that some or all of hypothermia's neuroprotective effects can be lost if cooling is used in nonsedated animals. One research group observed no protective effects when cooling was used after global anoxia in unsedated newborn piglets (201); when the experiment was repeated in the same model under adequate sedation, significant improvements in outcome were noted in hypothermia-treated animals (202). The authors concluded

that stress effects of cooling in unsedated animals could negate protective effects. Although this phenomenon has not been well studied in humans, it seems reasonable to avoid a cooling-related stress response.

Metabolism, Blood Gases, Glucose and Electrolytes. Hypothermia decreases the metabolic rate by $\pm 8\%$ per 1°C drop in core temperature, with a parallel decrease in oxygen consumption and production of carbon dioxide. This means that ventilator settings need to be adjusted frequently especially in the induction phase of hypothermia, to avoid accidental hyperventilation that can cause cerebral vasoconstriction. Blood gas values are temperature dependent, and if blood samples are warmed to 37°C before analysis (as is common in most laboratories), Po_2 and Pco_2 will be overestimated and pH underestimated in hypothermic patients (1). This topic has been reviewed elsewhere (203).

Blood gas management when CO_2 values measured at 37°C are kept constant (for example, at 40 mm Hg) regardless of the patient's actual core temperature is called *alpha-stat* management. The alternative is *pH-stat* management, in which Pco_2 is corrected for temperature and is maintained at a prespecified value. This implies that the temperature-corrected pH will remain constant during pH-stat management and will increase during alpha-stat management (1). Which of these two methods is superior remains to be determined, and may depend in part on whether or not cerebral autoregulation is preserved (1). Strictly applied alpha-stat management could lead to hyperventilation and cerebral vasoconstriction; strict pH-stat management could lead to hypercapnia, cerebral vasodilation, and increased ICP. Which option is preferable may depend on the precise nature of the neurological injury and the presence or absence of brain edema. Excessive hypocapnia can increase ischemia in injured areas of the brain, while excessive hypercapnia can increase brain edema; thus both can cause additional brain injury. These issues are more extensively addressed in a recently published review (1).

For accurate temperature correction, blood samples should be analyzed at the patient's real temperature; this can be most easily accomplished by performing on-site analysis in the ICU. If this is not possible, the blood gas values can be estimated in the following way.

In a sample assayed at 37°C :

- Subtract 5 mm Hg Po_2 per 1°C that the patient's temperature is $<37^\circ\text{C}$;
- Subtract 2 mm Hg Pco_2 per 1°C that the patient's temperature is $<37^\circ\text{C}$;
- Add 0.012 pH units per 1°C that the patient's temperature is $<37^\circ\text{C}$.

Thus, a sample from a patient with a core temperature of 33°C analyzed in the lab at 37°C with the results pH 7.45, Pco_2 35 mm Hg, and Po_2 80 mm Hg would have the following temperature-corrected values: pH 7.50, Pco_2 27 mm Hg, and Po_2 60 mm Hg.

In our unit, we use temperature-corrected blood gas values but avoid severe hypercapnia. Target Pco_2 values at 32°C (our usual target temperature in cardiac arrest patients) are 32 to 36 mm Hg, which would be 42 to 46 mm Hg in an uncorrected sample.

Hypothermia can also decrease insulin sensitivity and the amounts of insulin secreted by the pancreas. This can lead to hyperglycemia and/or an increase in the doses of insulin required to maintain glucose levels within target range. Sustained hyperglycemia has been linked to adverse outcome in critically ill patients, and may pose additional risks in patients with neurological injuries (39), although the optimal targets for glucose control remain controversial (39). Prevention and/or prompt correction of severe hyperglycemia should be part of the therapeutic strategy during hypothermia treatment. Furthermore, it should be realized that doses of insulin required to maintain normoglycemia are likely to decrease when the patient is rewarmed; this means that hypoglycemia can easily develop in the rewarming phase as insulin sensitivity is restored, particularly if the patient is rewarmed (too) quickly.

Cooling can also affect *electrolyte levels*. A combination of hypothermia-induced intracellular shift and tubular dysfunction leading to an increase in renal excretion of electrolytes can lead to depletion of magnesium, potassium, and phosphate during cooling (185). These electrolyte disorders can increase the risk for arrhythmias and other potentially harmful complications (204, 205). There is evidence suggesting that magnesium could play a role in mitigating various types of brain injury (204, 205); hypophosphatemia has been linked to respiratory problems and increased infection risk (206). Thus, electrolyte levels should

be kept in the high-normal range during and after hypothermia treatment.

Extra care should be taken during re-warming because hyperkalemia may develop during this phase, due to release of potassium sequestered to the intracellular compartment during hypothermia induction (1). Hyperkalemia can be prevented by slow and controlled re-warming, allowing the kidneys to excrete the excess potassium. Thus, the risk for rebound hyperkalemia in the re-warming phase should not be regarded as a reason to accept hypokalemia in the induction and maintenance phase. However, in patients with anuria or severe oliguria, renal replacement therapy should be started before the patient is rewarmed.

Cardiovascular and Hemodynamic Effects. Hypothermia has complex and opposing effects on the myocardium and myocardial contractility, depending on the patient's volume status and whether or not the patient is adequately sedated. Under normal circumstances, mild hypothermia will decrease heart rate and increase myocardial contractility in sedated and euvoletic patients (1, 207–211). Systolic function will improve but a mild degree of diastolic dysfunction may occur in some patients (1, 207–211). Blood pressure remains stable or increases slightly in most patients during mild hypothermia (1, 186, 211–213). Cardiac output decreases along with the heart rate; however, the hypothermia-induced decrease in metabolic rate usually equals or exceeds the decrease in cardiac output, so that the balance between supply and demand remains constant or improves (1).

The heart rate can be artificially increased by external pacing, or through administration of chronotropic drugs such as isoprenaline or dopamine. However, this is usually unnecessary; furthermore, myocardial contractility may be adversely affected (210, 211, 214). Two animal studies (210, 214) and one clinical study in patients undergoing cardiac surgery (211) reported that while increasing heart rate under normothermic conditions increased myocardial contractility and cardiac output, increasing the heart rate under hypothermic conditions significantly decreased myocardial contractility. When heart rates were not artificially increased, induction of mild hypothermia improved myocardial contractility (211).

Thus, in most patients the heart rate should be allowed to decrease along with the patient's core temperature; further-

more, it should be realized that normal values for heart rate change with temperature.

Despite these data, hypothermia is often still viewed as a (potential) cause of hypotension and (additional) myocardial dysfunction. There are several reasons for this:

Firstly, in contrast to mild hypothermia, deep hypothermia (<30°C) does decrease myocardial contractility (1, 215). Secondly, hypothermia can cause hypovolemia by inducing “cold diuresis” through a combination of increased venous return, activation of atrial natriuretic peptide, decreased levels of antidiuretic hormone and renal antidiuretic hormone receptor levels, and tubular dysfunction (185, 216–221). The increase in venous return is caused by constriction of peripheral vessels (particularly in the skin) due to hypothermia-induced increases in plasma norepinephrine levels and activation of the sympathetic nerve system. This leads to a shift of the blood from peripheral (small) veins to deeper veins and the core compartment of the body, increasing the venous return. In this way, hypothermia can induce hypovolemia, which if uncorrected can cause hypotension as well as electrolyte depletion and an increase in blood viscosity. The risk for hypovolemia increases significantly if the patient is simultaneously treated with diuretic agents such as mannitol. However, hypotension can be quite easily prevented by avoiding or promptly correcting hypovolemia, and by avoiding excessive stimulation of the heart rate.

Three studies in pediatric patients enrolling a total of 127 patients and two small studies in adult patients enrolling 18 patients have reported successful use of mild hypothermia (32°C–33°C) as rescue therapy to reverse refractory cardiac shock following cardiac surgery (222–226). Two additional studies have reported successful use of hypothermia in patients who remained comatose after a massive cardiac arrest leading to severe cardiac shock (227, 228). Rates of favorable outcome in patients with cardiac shock were 61% (14 of 23) in the first study and 39% (11 of 28) in the second study. Oddo et al (229) recently reported that the presence of cardiac shock on admission did not correlate with adverse outcome in a prospective series of 74 patients with witnessed cardiac arrest treated with hypothermia, regardless of initial rhythm or hemodynamic status. These observations suggest that cardiac

shock should not be regarded as a counterindication for hypothermia treatment.

Hypothermia also causes changes in the heart rhythm and electrocardiogram. The hypothermia-induced increase in venous return discussed above initially leads to mild sinus tachycardia. Tachycardia can be much more pronounced if the patient is insufficiently sedated, and the shivering response with increased oxygen consumption and heat generation can lead to more marked tachycardia.

This is soon followed by sinus bradycardia when core temperature drops below 35.5°C. The heart rate decreases progressively as core temperature drops, and at 33°C the normal heart rate will be 45 to 55 bpm (although there is wide interpatient variability). The underlying mechanism is a decrease in the rate of spontaneous depolarization of cardiac pacemaker cells (including those of the sinus node), as well as prolongation of the duration of action potentials and a mild decrease in the speed of myocardial impulse conduction. The most common electrocardiogram changes are prolonged PR intervals, increased QT interval, and widening of the QRS complex. Sometimes so-called Osborne waves can be seen, although this is relatively rare during mild hypothermia.

These electrocardiogram changes usually do not require treatment. It is often assumed that mild hypothermia increases the risk for arrhythmias and decreases the chance of successful treatment of arrhythmias, because the hypothermic myocardium becomes less responsive to antiarrhythmic drugs and more difficult to defibrillate. However, in general this applies only to deep hypothermia ($\leq 28^\circ\text{C}$); experimental evidence suggests that mild hypothermia can actually decrease the risk for arrhythmias, by stabilizing cell membranes and increasing the likelihood of successful defibrillation if arrhythmias do occur. Two studies have assessed the likelihood of successful defibrillation under hypothermic conditions compared with normothermia in a swine model. Both found higher rates of return of spontaneous circulation, increased likelihood of successful defibrillation, fewer shocks required to reach return of spontaneous circulation, and lower incidence of postdefibrillation asystole during mild hypothermia (230, 231). Similar observations have been reported in a rabbit model (232). Various case studies describe successful use of hypothermia to treat arrhythmias (junctional ectopic

tachycardia) in young infants (233–235). All of this indicates that mild hypothermia decreases rather than increases the risk for arrhythmias, and facilitates rather than complicates the treatment of arrhythmias.

However, this changes if core temperature drops below 28°C (in rare cases, <30°C in patients with severe electrolyte disorders and/or severe myocardial ischemia). More profound hypothermia does increase the risk for arrhythmias, usually beginning with atrial fibrillation, which can progress to more severe arrhythmias including ventricular tachycardia and ventricular fibrillation if temperature decreases further. Thus, a *de novo* development of atrial fibrillation in a patient with a core temperature <30°C should be viewed as a warning sign, and the patient should be rewarmed to $\geq 30^\circ\text{C}$ immediately. In general, a temperature drop below 30°C should be strenuously avoided during therapeutic cooling in the ICU, but cooling treatment should not be withheld because of the presence or risk for arrhythmias.

In addition, attending physicians and nurses should be aware that the myocardium becomes more sensitive to mechanical manipulations during deep hypothermia. For example, if a physician decides to perform chest compressions because of severe bradycardia at a core temperature <28°C, this manipulation can induce arrhythmias including ventricular fibrillation (1). An additional problem is that if arrhythmias do develop during deep hypothermia, they become significantly more difficult to treat, because the hypothermic myocardium can become less responsive to antiarrhythmic drugs (1). Furthermore, other treatments for atrial fibrillation, such as electric cardioversion, are frequently unsuccessful at low temperatures, and may actually trigger a conversion from atrial fibrillation to ventricular fibrillation. Cardioversion for nonlife-threatening arrhythmias should be avoided at core temperatures <34°C.

Coronary Perfusion and Ischemia. Hypothermia reduces metabolic rate and induces bradycardia, thereby providing protective effects for the ischemic myocardium. In addition, it has been shown that mild hypothermia (35°C) induces coronary vasodilation and increases myocardial perfusion in healthy volunteers (236, 237). In contrast, hypothermia can induce vasoconstriction in severely atherosclerotic coronary arteries (237). In addition, hypothermia can cause shiver-

ing, tachycardia, and increased myocardial oxygen consumption in insufficiently sedated patients. Indeed, accidental hypothermia in the perioperative setting increases the risk for morbid cardiac events (194, 195). Thus, preventing adverse effects depends on avoiding a stress response and effective suppression of shivering.

Some animal experiments and preliminary clinical studies have suggested that inducing mild hypothermia in the early stages following myocardial infarction could mitigate myocardial injury (17). This issue needs to be addressed in larger studies, but the available evidence certainly does not suggest that hypothermia increases myocardial injury.

Coagulation. Mild hypothermia can induce mild coagulopathy (1). Temperatures below 35°C can cause platelet dysfunction and a mild decrease in platelet count; at temperatures <33°C, other steps in the coagulation cascade, such as the synthesis and kinetics of clotting enzymes and plasminogen activator inhibitors, may also be affected (1). These effects have been studied mostly *in vitro*. Clinical observations suggest that the risk for severe bleeding associated with therapeutic cooling is relatively small; none of the studies in patients with cardiac arrest, severe TBI, or stroke have reported significant bleeding problems, although it should be emphasized that actively bleeding patients were excluded from all of these studies (1, 17). Schefold and coworkers (238) recently assessed the risk and severity of bleeding during simultaneous use of mild hypothermia and thrombolysis. They found that bleeding risks were similar to historical controls treated with thrombolytics alone, although there was a trend toward more red blood cell units being required to reach target hematocrit in hypothermic patients who developed bleeding complications and needed transfusions. However, neurological outcomes were significantly better in patients treated with both thrombolytics and hypothermia, even in those who developed bleeding complications (238). The authors conclude that bleeding risks should not be viewed as a reason to withhold hypothermia treatment.

These factors should be taken into account when deciding whether or not to use hypothermia in patients who are actively bleeding, or who have a high bleeding risk. If possible the (potential) source of bleeding should be (surgically) con-

trolled before cooling is initiated. If this is not possible, the risks of bleeding should be weighed against the benefits of cooling, and careful consideration should be given to the appropriate depth of hypothermia for that patient. Very mild hypothermia (35°C) does not affect coagulation, and can be safely used even if bleeding risks are high. Temperatures of 33°C to 35°C affect platelet function only; if surgical procedures are performed under hypothermic conditions, platelet transfusion may be considered. Coagulation factors other than platelet function are affected only when temperatures decrease below 33°C.

Drug Clearance. The speed of most enzyme-mediated reactions is temperature-dependent; therefore, the speed of these reactions is significantly slowed by hypothermia. One of the consequences is a decrease in the rates of drug metabolism by the liver (17, 239). Hypothermia-induced reductions in clearance have been demonstrated for a number of commonly used ICU medications, including vasoactive drugs such as epinephrine and norepinephrine; opiates such as morphine, fentanyl, and remifentanyl; sedatives such as propofol, volatile anesthetics, barbiturates, and midazolam; neuromuscular blocking agents such as rocuronium, atracurium, and vecuronium; and other drugs such as phenytoin, nitrates, and some beta-blockers (1, 239). Other drugs metabolized by the liver are likely to be affected in similar ways.

The net effects of hypothermia on drug action may be more complex than simply increasing the concentration of active metabolites. Temperature can also directly affect the body's response to specific drugs. For example, the effects of vasoactive drugs, such as epinephrine and norepinephrine, on blood pressure can be slightly blunted by hypothermia. Hypothermia-induced changes in the volume of distribution and renal function may also influence how drugs work under hypothermic conditions.

However, in most cases the effect of hypothermia will be to increase drug levels, thereby enhancing drug potency and effect duration. From a practical perspective, it seems reasonable to make empirical dose adjustments in drugs with relatively short half-lives that are metabolized by the liver (e.g., midazolam). When an increase in depth of sedation is required, or an episode of shivering needs to be treated, it is preferable to use bolus doses rather than to increase maintenance dose.

Drugs with long half-lives (e.g., amiodarone) are far less affected by these mechanisms, because the amounts eliminated in 24 hrs are low regardless of temperature; therefore, effects of amiodarone are less temperature-dependent, and clearance is only slightly decreased under hypothermic conditions.

Risk for Infections. Hypothermia inhibits the proinflammatory response through inhibition of leukocyte migration and phagocytosis and decreased synthesis of proinflammatory cytokines (1). Indeed, suppression of harmful neuroinflammation is one of the mechanisms through which hypothermia may exert its protective effects; the downside of this protective mechanism is an increased risk for infections. Development of (accidental) hypothermia in the perioperative setting has been linked to increased risk for infections of the respiratory tract and (surgical) wounds (1, 240–242). Some evidence suggests that this may even apply to controlled normothermia; i.e., suppression of a febrile response. For therapeutic cooling (i.e., cooling of sedated and mechanically ventilated patients) the risk for nosocomial infections appears to be closely linked to treatment duration (higher risk if treatment is continued for >24 hrs) and to the category of patients (lowest risk in cardiac arrest patients, highest risk in patients with severe stroke). Most of the studies reporting increased risks of infection during therapeutic cooling also observed that final outcomes did not appear to be adversely affected, even when infections occurred (17). The suppression of inflammatory responses in the lung has also been used therapeutically to improve oxygenation in patients with severe acute respiratory distress syndrome (17).

Strategies to deal with increased infection risks include considering antibiotic prophylaxis. Recent publications suggest that the use of selective decontamination of the digestive tract can significantly reduce the risk for nosocomial infections and decrease mortality in ICU patients (243, 244). Clinical studies with therapeutic cooling in settings in which selective decontamination of the digestive tract was used have reported low infection rates (186, 216), even when hypothermia was used for prolonged periods (216).

Patients should be carefully monitored for signs of infection during therapeutic temperature management. Some normal signs of infection are absent or

suppressed during hypothermia; this obviously applies to development of fever but may also apply to increased C-reactive protein level and elevated white blood cell count. One telltale sign of developing infection can be a sudden increase in the "workload" of the cooling device; i.e., a decrease in water temperature and increase in noise levels of the device that is being used to cool the patient. Such increases in cooling power may indicate either shivering or an attempt by the body to generate fever, and thus the presence of an infection.

Many treatment protocols (including in our ICU) include daily blood surveillance cultures to screen for bacteremia. The threshold for antibiotic treatment should be low. In patients developing infections after hypothermia treatment, fever should be treated symptomatically, to prevent new or additional neurological injuries (17).

The risk for wound infections may be further increased by hypothermia-induced cutaneous vasoconstriction. The same applies to the development of bedsores, which once formed will be more likely to progress and become infected because of the immune-suppressed state and complete immobilization of the patient. Nurses and physicians should be aware of these additional risks and take extra precautionary measures. Extra attention should also be paid to catheter and central line insertion sites, as local infections also are more likely to develop here.

Other Effects. Hypothermia is associated with impaired bowel function and delayed gastric emptying. There is no specific strategy to deal with this issue; routine measures such as insertion of duodenal probes and/or administration of metoclopramide or low-dose erythromycin may be considered, taking care that the combination of hypothermia with these drugs does not cause QT prolongation (1). Hypothermia often causes a mild rise in serum amylase; however, the risk for clinically significant pancreatitis is extremely low (1).

Many other changes in laboratory measurements can develop during hypothermia. Apart from those already mentioned (hyperglycemia; low electrolyte levels; decreased platelet and white blood cell counts; increased levels of glycerol, free fatty acids, ketonic acids, lactate, and amylase), these may include a rise in liver enzymes (serum glutamic oxaloacetic transaminase and serum glutamic pyru-

vic transaminase) and increased levels of cortisol, norepinephrine, and epinephrine (184).

CONCLUSION

Hypothermia is the first treatment with a proven ability to reverse postischemic injury in the clinical setting. As such, it represents a major breakthrough, the importance of which is only just beginning to be fully appreciated. However, there are numerous pitfalls in using therapeutic cooling to maximum efficacy, particularly if prolonged treatments are required to improve outcome, as seems to be the case for stroke and severe TBI. Understanding the mechanisms that underlie hypothermia's protective effects, awareness of the physiology of cooling, and knowing and understanding the potential side effects are all key factors in effectively using this therapeutic strategy. There are still many misconceptions surrounding therapeutic cooling; some risks are imagined or overestimated, whereas others are not fully appreciated.

Temperature management strategies are likely to play an increasingly important role in the care of (neuro)critically ill patients. In view of the numerous potential applications for induced hypothermia or controlled normothermia, it is increasingly important that physicians and nurses caring for critically ill patients, particularly those with neurological injuries, be aware of these issues, to help them avoid the pitfalls and risks of this treatment.

REFERENCES

1. Polderman KH, Herold I: Therapeutic hypothermia and controlled normothermia in the intensive care unit: Practical considerations, side effects, and cooling methods. *Crit Care Med* 2009; 37:1101-1120
2. Fay T: Observations on prolonged human refrigeration. *NY State J Med* 1940; 40: 1351-1354
3. Fay T: Observations on generalized refrigeration in cases of severe cerebral trauma. *Assoc Res Nerv Ment Dis Proc* 1945; 24: 611-619
4. Bigelow WG, Callaghan JC, Hopps JA: General hypothermia for experimental intracardiac surgery. The use of electrophrenic respirations, an artificial pacemaker for cardiac standstill and radio-frequency rewarming in general hypothermia. *Ann Surg* 1950; 132:531-539
5. Botterell EH, Lougheed WM, Scott JW, et al: Hypothermia and interruption of carotid, or carotid and vertebral circulation,

- in the management of intracranial aneurysms. *J Neurosurg* 1956; 13:1-42
6. Botterell EH, Lougheed WM, Morley TP, et al: Hypothermia in the surgical treatment of ruptured intracranial aneurysms. *J Neurosurg* 1958; 15:4-18
7. Bigelow WC: Methods for inducing hypothermia and rewarming. *Ann NY Acad Sci* 1959; 80:522-532
8. Benson DW, Williams GR, Spencer FC, et al: The use of hypothermia after cardiac arrest. *Anesth Analg* 1959; 38:423-428
9. Williams GR, Spencer FC: The clinical use of hypothermia after cardiac arrest. *Ann Surg* 1959; 148:462-468
10. Rosomoff HL: Protective effects of hypothermia against pathologic processes of the nervous system. *Ann NY Acad Sci* 1959; 80:475-486
11. Fay T: Early experiences with local and general refrigeration of the human brain. *J Neurosurg* 1959; 16:239-259
12. Lazorthes G, Campan L: Moderate hypothermia in the treatment of head injuries. *Clin Neurosurg* 1964; 12:293-299
13. Hamby WB: Intracranial surgery for aneurysm: Effect of hypothermia upon survival. *J Neurosurg* 1963; 20:41-45
14. Safar P: Long-term resuscitation in intensive care units. *Anesthesiology* 1964; 25: 216-222
15. Rosomoff HL, Safar P: Management of the comatose patient. *Clin Anesth* 1965; 1:244-258
16. Bernard SA, Jones BM, Horne MK: Clinical trial of induced hypothermia in comatose survivors of out-of-hospital cardiac arrest. *Ann Emerg Med* 1997; 30:146-153
17. Polderman KH: Induced hypothermia and fever control for prevention and treatment of neurological injuries. *Lancet* 2008; 371: 1955-1969
18. Jastremski M, Sutton-Tyrrell K, Vaagenes P, et al: Glucocorticoid treatment does not improve neurological recovery following cardiac arrest. Brain Resuscitation Clinical Trial I Study Group. *JAMA* 1989; 262: 3427-3430
19. Brain Resuscitation Clinical Trial II Study Group: Randomized clinical study of thiopental loading in comatose survivors of cardiac arrest. *N Engl J Med* 1986; 314: 397-403
20. Brain Resuscitation Clinical Trial II Study Group: A randomized clinical study of a calcium-entry blocker (lidoflazine) in the treatment of comatose survivors of cardiac arrest. *N Engl J Med* 1991; 324:1225-1231
21. Helfaer MA, Ichord RN, Martin LJ, et al: Treatment with the competitive NMDA antagonist GPI 3000 does not improve outcome after cardiac arrest in dogs. *Stroke* 1998; 29:824-829
22. Silbergleit R, Haywood Y, Fiskum G, et al: Lack of a neuroprotective effect from N-acetylcysteine after cardiac arrest and resuscitation in a canine model. *Resuscitation* 1999; 40:181-186

23. Auer RN: Non-pharmacologic (physiologic) neuroprotection in the treatment of brain ischemia. *Ann NY Acad Sci* 2001; 939: 271–282
24. Corbett D, Nurse S: The problem of assessing effective neuroprotection in experimental cerebral ischemia. *Prog Neurobiol* 1998; 54:531–548
25. Siesjo BK, Bengtsson F, Grampp W, et al: Calcium, excitotoxins, and neuronal death in brain. *Ann NY Acad Sci* 1989; 568: 234–251
26. Dietrich WD, Busto R, Alonso O, et al: Intrischemic but not post-ischemic brain hypothermia protects chronically following global forebrain ischemia in rats. *J Cereb Blood Flow Metab* 1993; 13:541–549
27. Small DL, Morley P, Buchan AM: Biology of ischemic cerebral cell death. *Prog Cardiovasc Dis* 1999; 42:185–207
28. Polderman KH: Application of therapeutic hypothermia in the ICU: Opportunities and pitfalls of a promising treatment modality. Part I: Indications and evidence. *Intensive Care Med* 2004; 30:556–575
29. Milde LN: Clinical use of mild hypothermia for brain protection: A dream revisited. *J Neurosurg Anesthesiol* 1992; 4:211–215
30. Takata K, Takeda Y, Morita K: Effects of hypothermia for a short period on histological outcome and extracellular glutamate concentration during and after cardiac arrest in rats. *Crit Care Med* 2005; 33: 1340–1345
31. Hagerdal M, Harp J, Nilsson L, et al: The effect of induced hypothermia upon oxygen consumption in the rat brain. *J Neurochem* 1975; 24:311–316
32. Palmer C, Vannucci RC, Christensen MA, et al: Regional cerebral blood flow and glucose utilization during hypothermia in newborn dogs. *Anesthesiology* 1989; 71:730–737
33. Aoki M, Nomura F, Stromski ME, et al: Effects of pH on brain energetics after hypothermic circulatory arrest. *Ann Thorac Surg* 1993; 55:1093–1103
34. Ehrlich MP, McCullough JN, Zhang N, et al: Effect of hypothermia on cerebral blood flow and metabolism in the pig. *Ann Thorac Surg* 2002; 73:191–197
35. Erecinska M, Thoresen M, Silver IA: Effects of hypothermia on energy metabolism in mammalian central nervous system. *J Cereb Blood Flow Metab* 2003; 23: 513–530
36. Coles JP, Minhas PS, Fryer TD, et al: Effect of hyperventilation on cerebral blood flow in traumatic head injury: Clinical relevance and monitoring correlates. *Crit Care Med* 2002; 30:1950–1959
37. Coles JP, Fryer TD, Coleman MR, et al: Hyperventilation following head injury: Effect on ischemic burden and cerebral oxidative metabolism. *Crit Care Med* 2007; 35: 568–578
38. Lin JS, Chen YS, Chiang HS, et al: Hypoxic preconditioning protects rat hearts against ischemia/reperfusion injury: Role of erythropoietin on progenitor cell mobilization. *J Physiol* 2008; 586(Pt 23):5757–5769
39. Povlishock JT, Buki A, Koizumi H, et al: Initiating mechanisms involved in the pathobiology of traumatically induced axonal injury and interventions targeted at blunting their progression. *Acta Neurochir Suppl (Wien)* 1999; 73:15–20
40. Xu L, Yenari MA, Steinberg GK, et al: Mild hypothermia reduces apoptosis of mouse neurons in vitro early in the cascade. *J Cereb Blood Flow Metab* 2002; 22:21–28
41. Adachi M, Sohma O, Tsuneishi S, et al: Combination effect of systemic hypothermia and caspase inhibitor administration against hypoxic-ischemic brain damage in neonatal rats. *Pediatr Res* 2001; 50: 590–595
42. Ning XH, Chen SH, Xu CS, et al: Hypothermic protection of the ischemic heart via alterations in apoptotic pathways as assessed by gene array analysis. *J Appl Physiol* 2002; 92:2200–2207
43. Liou AK, Clark RS, Henshall DC, et al: To die or not to die for neurons in ischemia, traumatic brain injury and epilepsy: A review on the stress-activated signaling pathways and apoptotic pathways. *Prog Neurobiol* 2003; 69:103–142
44. Raghupathi R, Graham DI, McIntosh TK: Apoptosis after traumatic brain injury. *J Neurotrauma* 2000; 17:927–938
45. Leker RR, Shohami E: Cerebral ischemia and trauma—different etiologies yet similar mechanisms: Neuroprotective opportunities. *Brain Res Brain Res Rev* 2002; 39: 55–73
46. Hall ED: Free radicals in stroke. In: *Stroke Therapy: Basic, Preclinical, and Clinical Directions*. Miller LP (Ed). New York, Wiley-Liss, 1999, pp 245–270
47. Globus MY-T, Busto R, Lin B, et al: Detection of free radical activity during transient global ischemia and recirculation: Effects of intra-ischemic brain temperature modulation. *J Neurochem* 1995; 65:1250–1256
48. Dempsey RJ, Combs DJ, Maley ME, et al: Moderate hypothermia reduces postischemic edema development and leukotriene production. *Neurosurgery* 1987; 21: 177–181
49. Busto R, Dietrich WD, Globus MY, et al: Small differences in intrischemic brain temperature critically determine the extent of ischemic neuronal injury. *J Cereb Blood Flow Metab* 1987; 7:729–738
50. Globus MY-T, Alonso O, Dietrich WD, et al: Glutamate release and free radical production following brain injury: Effects of post-traumatic hypothermia. *J Neurochem* 1995; 65:1704–1711
51. Baker AJ, Zornow MH, Grafe MR, et al: Hypothermia prevents ischemia-induced increases in hippocampal glycine concentrations in rabbits. *Stroke* 1991; 22: 666–673
52. Busto R, Globus MY, Dietrich WD, et al: Effect of mild hypothermia on ischemia-induced release of neurotransmitters and free fatty acids in rat brain. *Stroke* 1989; 20:904–910
53. Kaibara T, Sutherland GR, Colbourne F, et al: Hypothermia: Depression of tricarboxylic acid cycle flux and evidence for pentose phosphate shunt upregulation. *J Neurosurg* 1999; 90:339–347
54. Kuboyama K, Safar P, Radovsky A, et al: Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: A prospective, randomized study. *Crit Care Med* 1993; 21:1348–1358
55. Baker CJ, Onesti ST, Solomon RA: Reduction by delayed hypothermia of cerebral infarction following middle cerebral artery occlusion in the rat: A time-course study. *J Neurosurg* 1992; 77:438–444
56. Hickey RW, Ferimer H, Alexander HL, et al: Delayed, spontaneous hypothermia reduces neuronal damage after asphyxial cardiac arrest in rats. *Crit Care Med* 2000; 28: 3511–3516
57. Coimbra C, Wieloch T: Hypothermia ameliorates neuronal survival when induced 2 hours after ischaemia in the rat. *Acta Physiol Scand* 1992; 146:543–544
58. Colbourne F, Li H, Buchan AM: Indefatigable CA1 sector neuroprotection with mild hypothermia induced 6 hours after severe forebrain ischemia in rats. *J Cereb Blood Flow Metab* 1999; 19:742–749
59. Coimbra C, Wieloch T: Moderate hypothermia mitigates neuronal damage in the rat brain when initiated several hours following transient cerebral ischemia. *Acta Neuropathol* 1994; 87:325–333
60. Leonov Y, Sterz F, Safar P, et al: Moderate hypothermia after cardiac arrest of 17 minutes in dogs: Effect on cerebral and cardiac outcome: A preliminary study. *Stroke* 1990; 21:1600–1606
61. Leonov Y, Sterz F, Safar P, et al: Mild cerebral hypothermia during and after cardiac arrest improves neurologic outcome in dogs. *J Cereb Blood Flow Metab* 1990; 10: 57–70
62. Sterz F, Safar P, Tisherman S, et al: Mild hypothermic cardiopulmonary resuscitation improves outcome after prolonged cardiac arrest in dogs. *Crit Care Med* 1991; 19:379–389
63. Safar P, Xiao F, Radovsky A, et al: Improved cerebral resuscitation from cardiac arrest in dogs with mild hypothermia plus blood flow promotion. *Stroke* 1996; 27:105–113
64. Kollmar R, Schabitz WR, Heiland S, et al: Neuroprotective effect of delayed moderate hypothermia after focal cerebral ischemia: An MRI study. *Stroke* 2002; 33:1899–1904
65. Weinrauch V, Safar P, Tisherman S, et al: Beneficial effect of mild hypothermia and detrimental effect of deep hypothermia after cardiac arrest in dogs. *Stroke* 1992; 23: 1454–1462
66. Gisvold SE, Safar P, Rao G, et al: Multifac-

- eted therapy after global brain ischemia in monkeys. *Stroke* 1984; 15:803–812
67. Markarian GZ, Lee JH, Stein DJ, et al: Mild hypothermia: Therapeutic window after experimental cerebral ischemia. *Neurosurgery* 1996; 38:542–550
 68. Dietrich WD, Busto R, Globus MY, et al: Brain damage and temperature: Cellular and molecular mechanisms. *Adv Neurol* 1996; 71:177–194
 69. Schmidt OI, Heyde CE, Ertel W, et al: Closed head injury—an inflammatory disease? *Brain Res Brain Res Rev* 2005; 48:388–399
 70. Asensio VC, Campbell IL: Chemokines in the CNS: Plurifunctional mediators in diverse states. *Trends Neurosci* 1999; 22:504–512
 71. Merrill JE, Benveniste EN: Cytokines in inflammatory brain lesions: Helpful and harmful. *Trends Neurosci* 1996; 19:331–338
 72. Morganti-Kossmann MC, Rancan M, Stahel PF, et al: Inflammatory response in acute traumatic brain injury: A double-edged sword. *Curr Opin Crit Care* 2002; 8:101–105
 73. Schoettle RJ, Kochanek PM, Magargee MJ, et al: Early polymorphonuclear leukocyte accumulation correlates with the development of posttraumatic cerebral edema in rats. *J Neurotrauma* 1990; 7:207–217
 74. Patel HC, Boutin H, Allan SM: Interleukin-1 in the brain: Mechanisms of action in acute neurodegeneration. *Ann NY Acad Sci* 2003; 992:39–47
 75. Aibiki M, Maekawa S, Ogura S, et al: Effect of moderate hypothermia on systemic and internal jugular plasma IL-6 levels after traumatic brain injury in humans. *J Neurotrauma* 1999; 16:225–232
 76. Kimura A, Sakurada S, Ohkuni H, et al: Moderate hypothermia delays proinflammatory cytokine production of human peripheral blood mononuclear cells. *Crit Care Med* 2002; 30:1499–1502
 77. Suehiro E, Fujisawa H, Akimura T, et al: Increased matrix metalloproteinase-9 in blood in association with activation of interleukin-6 after traumatic brain injury: Influence of hypothermic therapy. *J Neurotrauma* 2004; 21:1706–1711
 78. Dietrich WD, Chatzipanteli K, Vitarbo E, et al: The role of inflammatory processes in the pathophysiology and treatment of brain and spinal cord trauma. *Acta Neurochir Suppl* 2004; 89:69–74
 79. Novack TA, Dillon MC, Jackson WT: Neurochemical mechanisms in brain injury and treatment: A review. *J Clin Exp Neuropsychol* 1996; 18:685–706
 80. Raghupathi R, McIntosh TK: Pharmacotherapy for traumatic brain injury: A review. *Proc West Pharmacol Soc* 1998; 41:241–246
 81. Huang ZG, Xue D, Preston E, et al: Biphasic opening of the blood–brain barrier following transient focal ischemia: Effects of hypothermia. *Can J Neurol Sci* 1999; 26:298–304
 82. Chi OZ, Liu X, Weiss HR: Effects of mild hypothermia on blood–brain barrier disruption during isoflurane or pentobarbital anesthesia. *Anesthesiology* 2001; 95:933–938
 83. Smith SL, Hall ED: Mild pre- and posttraumatic hypothermia attenuates blood–brain barrier damage following controlled cortical impact injury in the rat. *J Neurotrauma* 1996; 13:1–9
 84. Jurkovich GJ, Pitt RM, Curreri PW, et al: Hypothermia prevents increased capillary permeability following ischemia–reperfusion injury. *J Surg Res* 1988; 44:514–521
 85. Kinoshita K, Chatzipanteli K, Alonso OF, et al: The effect of brain temperature on hemoglobin extravasation after traumatic brain injury. *J Neurosurg* 2002; 97:945–953
 86. Fischer S, Clauss M, Wiesnet M, et al: Hypoxia induces permeability in brain microvessel endothelial cells via VEGF and NO. *Am J Physiol* 1999; 276:C812–C820
 87. Fischer S, Renz D, Wiesnet M, et al: Hypothermia abolishes hypoxia-induced hyperpermeability in brain microvessel endothelial cells. *Brain Res Mol Brain Res* 1999; 74:135–144
 88. Polderman KH, Ely EW, Badr AE, et al: Induced hypothermia in traumatic brain injury: Considering the conflicting results of meta-analyses and moving forward. *Intensive Care Med* 2004; 30:1860–1864
 89. Ghajar J: Traumatic brain injury. *Lancet* 2000; 356:923–929
 90. Reinert MM, Bullock R: Clinical trials in head injury. *Neurol Res* 1999; 21:330–338
 91. Yokota S, Imagawa T, Miyamae T, et al: Hypothetical pathophysiology of acute encephalopathy and encephalitis related to influenza virus infection and hypothermia therapy. *Pediatr Int* 2000; 42:197–203
 92. Morimoto Y, Kemmotsu O, Kitami K, et al: Acute brain swelling after out-of-hospital cardiac arrest: Pathogenesis and outcome. *Crit Care Med* 1993; 21:104–110
 93. Brain Trauma Foundation: Guidelines for the management of severe traumatic brain injury. <http://www.braintrauma.org>. Accessed May 13, 2009.
 94. Stocchetti N, Colombo A, Ortolano F, et al: Time course of intracranial hypertension after traumatic brain injury. *J Neurotrauma* 2007; 24:1339–1346
 95. Clifton GL, Miller ER, Choi SC, et al: Lack of effect of induction of hypothermia after acute brain injury. *N Engl J Med* 2001; 344:556–563
 96. McIntyre LA, Fergusson DA, Hebert PC, et al: Prolonged therapeutic hypothermia after traumatic brain injury in adults: A systematic review. *JAMA* 2003; 289:2992–2999
 97. Schaller B, Graf R: Hypothermia and stroke: The pathophysiological background. *Pathophysiology* 2003; 10:7–35
 98. Ding D, Moskowitz SI, Li R, et al: Acidosis induces necrosis and apoptosis of cultured hippocampal neurons. *Exp Neurol* 2000; 162:1–12
 99. Natale JA, D'Alecy LG: Protection from cerebral ischemia by brain cooling without reduced lactate accumulation in dogs. *Stroke* 1989; 20:770–777
 100. Jiang JY, Liang YM, Luo QZ, et al: Effect of mild hypothermia on brain dialysate lactate after fluid percussion brain injury in rodents. *Neurosurgery* 2004; 54:713–717
 101. Chatauret N, Rose C, Therrien G, et al: Mild hypothermia prevents cerebral edema and CSF lactate accumulation in acute liver failure. *Metab Brain Dis* 2001; 16:95–102
 102. Amess PN, Penrice J, Cady EB, et al: Mild hypothermia after severe transient hypoxia—ischemia reduces the delayed rise in cerebral lactate in the newborn piglet. *Pediatr Res* 1997; 41:803–808
 103. Vejchapipat P, Proctor E, Ramsay A, et al: Intestinal energy metabolism after ischemia–reperfusion: Effects of moderate hypothermia and perfluorocarbons. *J Pediatr Surg* 2002; 37:786–790
 104. Chatauret N, Zwingmann C, Rose C, et al: Effects of hypothermia on brain glucose metabolism in acute liver failure: A H/C-nuclear magnetic resonance study. *Gastroenterology* 2003; 125:815–824
 105. Vaquero J, Blei AT: Mild hypothermia for acute liver failure: A review of mechanisms of action. *J Clin Gastroenterol* 2005; 39:S147–S157
 106. Lanier WL: Cerebral metabolic rate and hypothermia: Their relationship with ischemic neurologic injury. *J Neurosurg Anesthesiol* 1995; 7:216–221
 107. Soukup J, Zauner A, Döppner EM, et al: Relationship between brain temperature, brain chemistry and oxygen delivery after severe human head injury: The effect of mild hypothermia. *Neurol Res* 2002; 24:161–168
 108. Kimura T, Sako K, Tanaka K, et al: Effect of mild hypothermia on energy state recovery following transient forebrain ischemia in the gerbil. *Exp Brain Res* 2002; 145:83–90
 109. Nybo L, Secher NH, Nielsen B: Inadequate heat release from the human brain during prolonged exercise with hyperthermia. *J Physiol* 2002; 545:697–704
 110. Mariak Z, Bondyra Z, Piekarska M: The temperature within the circle of Willis versus tympanic temperature in resting normothermic humans. *Eur J Appl Physiol Occup Physiol* 1993; 66:518–520
 111. Crowder CM, Tempelhoff R, Theard MA, et al: Jugular bulb temperature: Comparison with brain surface and core temperatures in neurosurgical patients during mild hypothermia. *J Neurosurg* 1996; 85:98–103
 112. McIlvoy L: Comparison of brain temperature to core temperature: A review of the literature. *J Neurosci Nurs* 2004; 36:23–31
 113. Henker RA, Brown SD, Marion DW: Comparison of brain temperature with bladder and rectal temperatures in adults with se-

- vere head injury. *Neurosurgery* 1998; 42: 1071–1075
114. Rumana CS, Gopinath SP, Uzura M, et al: Brain temperature exceeds systemic temperature in head-injured patients. *Crit Care Med* 1998; 26:562–567
 115. Verlooy J, Heytens L, Veeckmans G, et al: Intracerebral temperature monitoring in severely head injured patients. *Acta Neurochir (Wien)* 1995; 134:76–78
 116. Møllergaard P, Nordstrom CH: Intracerebral temperature in neurosurgical patients. *Neurosurgery* 1991; 28:709–713
 117. Schwab S, Spranger M, Aschhoff A, et al: Brain temperature monitoring and modulation in patients with severe MCA infarction. *Neurology* 1997; 48:762–767
 118. Childs C, Vail A, Protheroe R, et al: Differences between brain and rectal temperatures during routine critical care of patients with severe traumatic brain injury. *Anaesthesia* 2005; 60:759–765
 119. Albrecht RF, Wass CT, Lanier WL: Occurrence of potentially detrimental temperature alterations in hospitalized patients at risk for brain injury. *Mayo Clin Proc* 1998; 73:629–635
 120. Albrecht RF 2nd, Wass CT, Lanier WL, et al: Uncontrolled hyperthermia in the neurosurgical intensive care unit. *Neurosurgery* 2000; 47:850–855
 121. Schwarz S, Hafner K, Aschoff A, et al: Incidence and prognostic significance of fever following intracerebral hemorrhage. *Neurology* 2000; 54:354–361
 122. Reith J, Jørgensen HS, Pedersen PM, et al: Body temperature in acute stroke: Relations to stroke severity, infarct size, mortality, and outcome. *Lancet* 1996; 347:422–425
 123. Hajat C, Hajat S, Sharma P: Effects of post-stroke pyrexia on stroke outcome: Meta-analysis of studies on patients. *Stroke* 2000; 31:410–414
 124. Gorbach AM, Heiss J, Kufta C, et al: Intraoperative infrared functional imaging of human brain. *Ann Neurol* 2003; 54:297–309
 125. Collins CM, Smith MB, Turner R: Model of local temperature changes in brain upon functional activation. *J Appl Physiol* 2004; 97:2051–2055
 126. Gorbach AM: Infrared imaging of brain function. *Adv Exp Med Biol* 1993; 333: 95–123
 127. Ecker RD, Goerss SJ, Meyer FB, et al: Vision of the future: Initial experience with intraoperative real-time high-resolution dynamic infrared imaging. Technical note. *J Neurosurg* 2002; 97:1460–1471
 128. Hayashi N, Hirayama T, Udagawa A, et al: Systemic management of cerebral edema based on a new concept in severe head injury patients. *Acta Neurochir Suppl (Wien)* 1994; 60:541–543
 129. Piepgras A, Elste V, Frietsch T, et al: Effect of moderate hypothermia on experimental severe subarachnoid hemorrhage, as evaluated by apparent diffusion coefficient changes. *Neurosurgery* 2001; 48:1128–1135
 130. Wass CT, Lanier WL, Hofer RE, et al: Temperature changes of $> \text{or} = 1^\circ\text{C}$ alter functional neurologic outcome and histopathology in a canine model of complete cerebral ischemia. *Anesthesiology* 1995; 83:325–335
 131. Baena RC, Busto R, Dietrich WD, et al: Hyperthermia delayed by 24 hours aggravates neuronal damage in rat hippocampus following global ischemia. *Neurology* 1997; 48:768–773
 132. Kim Y, Busto R, Dietrich WD, et al: Delayed postischemic hyperthermia in awake rats worsens the histopathological outcome of transient focal cerebral ischemia. *Stroke* 1996; 27:2274–2281
 133. Hickey RW, Kochanek PM, Ferimer H, et al: Induced hyperthermia exacerbates neurologic neuronal histologic damage after asphyxial cardiac arrest in rats. *Crit Care Med* 2003; 31:531–535
 134. Chen H, Chopp M, Welch KMA: Effect of mild hyperthermia on the ischemic infarct volume after middle cerebral artery occlusion in the rat. *Neurology* 1991; 41: 1133–1135
 135. Wass CT, Lanier WL, Hofer RE, et al: Temperature changes of $> \text{or} = 1$ degree C alter functional neurologic outcome and histopathology in a canine model of complete cerebral ischemia. *Anesthesiology* 1995; 83: 325–335
 136. Ginsberg MD, Sternau LL, Globus MY-T, et al: Therapeutic modulation of brain temperature: Relevance to ischemic brain injury. *Cerebrovasc Brain Metab Rev* 1992; 4:189–225
 137. Takino M, Okada Y: Hyperthermia following cardiopulmonary resuscitation. *Intensive Care Med* 1991; 17:419–420
 138. Zeiner A, Holzer M, Sterz F, et al: Hyperthermia after cardiac arrest is associated with an unfavorable neurologic outcome. *Arch Intern Med* 2001; 161:2007–2012
 139. Soukup J, Zauner A, Doppenberg EM, et al: The importance of brain temperature in patients after severe head injury: Relationship to intracranial pressure, cerebral perfusion pressure, cerebral blood flow, and outcome. *J Neurotrauma* 2002; 19:559–571
 140. Azzimondi G, Bassein L, Nonino F, et al: Fever in acute stroke worsens prognosis. A prospective study. *Stroke* 1995; 26: 2040–2043
 141. Castillo J, Davalos A, Marrugat J, et al: Timing for fever-related brain damage in acute ischemic stroke. *Stroke* 1998; 29:2455–2460
 142. Kammersgaard LP, Jørgensen HS, Rungby JA, et al: Admission body temperature predicts long-term mortality after acute stroke: The Copenhagen Stroke Study. *Stroke* 2002; 33:1759–1762
 143. Böttiger BW, Motsch J, Bohrer H, et al: Activation of blood coagulation after cardiac arrest is not balanced adequately by activation of endogenous fibrinolysis. *Circulation* 1995; 92:2572–2578
 144. Böttiger BW, Martin E: Thrombolytic therapy during cardiopulmonary resuscitation and the role of coagulation activation after cardiac arrest. *Curr Opin Crit Care* 2001; 7:176–183
 145. Gando S, Kameue T, Nanzaki S, et al: Massive fibrin formation with consecutive impairment of fibrinolysis in patients with out-of-hospital cardiac arrest. *Thromb Haemost* 1997; 77:278–282
 146. Fischer M, Böttiger BW, Popov-Cenic S, et al: Thrombolysis using plasminogen activator and heparin reduces cerebral no-reflow after resuscitation from cardiac arrest: An experimental study in the cat. *Intensive Care Med* 1996; 22:1214–1223
 147. Darius H, Yanagisawa A, Brezinski ME, et al: Beneficial effects of tissue-type plasminogen activator in acute myocardial ischemia in cats. *J Am Coll Cardiol* 1986; 8:125–131
 148. Kim YH, Park JH, Hong SH, et al: Nonproteolytic neuroprotection by human recombinant tissue plasminogen activator. *Science* 1999; 284:647–650
 149. Böttiger BW, Bode C, Kern S, et al: Efficacy and safety of thrombolytic therapy after initially unsuccessful cardiopulmonary resuscitation: A prospective clinical trial. *Lancet* 2001; 357:1583–1585
 150. Spöhr F, Arntz HR, Bluhmki E, et al: International multicentre trial protocol to assess the efficacy and safety of tenecteplase during cardiopulmonary resuscitation in patients with out-of-hospital cardiac arrest: The Thrombolysis in Cardiac Arrest (TROICA) Study. *Eur J Clin Invest* 2005; 35:315–323
 151. Valeri CR, Feingold H, Cassidy G, et al: Hypothermia-induced reversible platelet dysfunction. *Ann Surg* 1987; 205:175–181
 152. Michelson AD, MacGregor H, Barnard MR, et al: Hypothermia-induced reversible platelet dysfunction. *Thromb Haemost* 1994; 71:633–640
 153. Valeri CR, MacGregor H, Cassidy G, et al: Effects of temperature on bleeding time and clotting time in normal male and female volunteers. *Crit Care Med* 1995; 23: 698–704
 154. Watts DD, Trask A, Soeken K, et al: Hypothermic coagulopathy in trauma: Effect of varying levels of hypothermia on enzyme speed, platelet function, and fibrinolytic activity. *J Trauma* 1998; 44:846–854
 155. Leffer CW: Prostanoids: Intrinsic modulation of cerebral circulation. *News Physiol Sci* 1997; 12:72–77
 156. Dogne JM, de Leval X, Hanson J, et al: New developments on thromboxane and prostacyclin modulators part I: Thromboxane modulators. *Curr Med Chem* 2004; 11: 1223–1241
 157. De Leval X, Hanson J, David JL, et al: New developments on thromboxane and prostacyclin modulators part II: Prostacyclin modulators. *Curr Med Chem* 2004; 11: 1243–1252
 158. Westcott JY, Murphy RC, Stenmark K: Eicosanoids in human ventricular cerebrospi-

- nal fluid following severe brain injury. *Prostaglandins* 1987; 34:877–887
159. Hsu CY, Halushka PV, Hogan EL, et al: Alteration of thromboxane and prostacyclin levels in experimental spinal cord injury. *Neurology* 1985; 35:1003–1009
 160. Chen ST, Hsu CY, Hogan EL, et al: Thromboxane, prostacyclin, and leukotrienes in cerebral ischemia. *Neurology* 1986; 36:466–470
 161. Maekawa S, Aibiki M, Ogura S, et al: Mild hypothermia suppresses thromboxane B2 production in brain-injured patients. *In: The Immune Consequences of Trauma, Shock and Sepsis. Mechanisms and Therapeutic Approaches.* Faist E (Ed). Bologna, Italy, Monduzzi Editore, 1997, pp 135–138
 162. Aibiki M, Maekawa S, Yokono S: Moderate hypothermia improves imbalances of thromboxane A2 and prostaglandin I2 production after traumatic brain injury in humans. *Crit Care Med* 2000; 28:3902–3906
 163. Chen L, Piao Y, Zeng F, et al: Moderate hypothermia therapy for patients with severe head injury. *Chin J Traumatol* 2001; 4:164–167
 164. Yuan HB, Huang Y, Zheng S, et al: Hypothermic preconditioning increases survival of purkinje neurons in rat cerebellar slices after an in vitro simulated ischemia. *Anesthesiology* 2004; 100:331–337
 165. Yunoki M, Nishio S, Ukita N, et al: Hypothermic preconditioning induces rapid tolerance to focal ischemic injury in the rat. *Exp Neurol* 2003; 181:291–300
 166. Strauch JT, Lauten A, Spielvogel D, et al: Mild hypothermia protects the spinal cord from ischemic injury in a chronic porcine model. *Eur J Cardiothorac Surg* 2004; 25:708–715
 167. Oku K, Kuboyama K, Safar P, et al: Cerebral and systemic arteriovenous oxygen monitoring after cardiac arrest: Inadequate cerebral oxygen delivery. *Resuscitation* 1994; 27:141–152
 168. Jordan KG: Nonconvulsive status epilepticus in acute brain injury. *J Clin Neurophysiol* 1999; 16:332–340
 169. Krumholz A: Epidemiology and evidence for morbidity of nonconvulsive status epilepticus. *J Clin Neurophysiol* 1999; 16:314–322
 170. Vespa PM, Nuwer MR, Nenov V, et al: Increased incidence and impact of nonconvulsive and convulsive seizures after traumatic brain injury as detected by continuous electroencephalographic monitoring. *J Neurosurg* 1999; 91:750–760
 171. Towne AR, Waterhouse EJ, Boggs JG, et al: Prevalence of nonconvulsive status epilepticus in comatose patients. *Neurology* 2000; 54:340–345
 172. Dennis LJ, Claassen J, Hirsch LJ, et al: Nonconvulsive status epilepticus after subarachnoid hemorrhage. *Neurosurgery* 2002; 51:1136–1143
 173. Maeda T, Hashizume K, Tanaka T: Effect of hypothermia on kainic acid-induced limbic seizures: An electroencephalographic and 14C-deoxyglucose autoradiographic study. *Brain Res* 1999; 818:228–235
 174. Karkar KM, Garcia PA, Bateman LM, et al: Focal cooling suppresses spontaneous epileptiform activity without changing the cortical motor threshold. *Epilepsia* 2002; 43:932–935
 175. Corry JJ, Dhar R, Murphy T, et al: Hypothermia for refractory status epilepticus. *Neurocrit Care* 2008; 9:189–197
 176. Lundgren J, Smith ML, Blennow G, et al: Hypothermia aggravates and hypothermia ameliorates epileptic brain damage. *Exp Brain Res* 1994; 99:43–55
 177. Yager JY, Armstrong EA, Jaharus C, et al: Preventing hyperthermia decreases brain damage following neonatal hypoxic-ischemic seizures. *Brain Res* 2004; 1011:48–57
 178. Liu Z, Gatt A, Mikati M, et al: Effect of temperature on kainic acid-induced seizures. *Brain Res* 1993; 631:51–58
 179. Somjen GG, Aitken PG, Balestrino M, et al: Spreading depression-like depolarization and selective vulnerability of neurons. A brief review. *Stroke* 1990; 21(Suppl):III-179–III-183
 180. Lundgren J, Smith ML, Siesjo BK: Influence of moderate hypothermia on ischemic brain damage incurred under hyperglycemic conditions. *Exp Brain Res* 1991; 84:91–101
 181. Church AJ, Andrew RD: Spreading depression expands traumatic injury in neocortical brain slices. *J Neurotrauma* 2005; 22:277–290
 182. Yenari MA, Onley D, Hedehus M, et al: Diffusion- and perfusion-weighted magnetic resonance imaging of focal cerebral ischemia and cortical spreading depression under conditions of mild hypothermia. *Brain Res* 2000; 885:208–219
 183. Polderman KH, Callaghan J: Equipment review: Cooling catheters to induce therapeutic hypothermia? *Crit Care* 2006; 10:234
 184. Polderman KH: Application of therapeutic hypothermia in the intensive care unit. Opportunities and pitfalls of a promising treatment modality. Part 2: Practical aspects and side effects. *Intensive Care Med* 2004; 30:757–769
 185. Polderman KH, Peerdeman SM, Girbes ARJ: Hypophosphatemia and hypomagnesemia induced by cooling in patients with severe head injury. *J Neurosurg* 2001; 94:697–705
 186. Polderman KH, Rijnsburger ER, Peerdeman SM, et al: Induction of hypothermia in patients with various types of neurologic injury with use of large volumes of ice-cold intravenous fluid. *Crit Care Med* 2005; 33:2744–2751
 187. Maxwell WL, Watson A, Queen R, et al: Slow, medium, or fast re-warming following post-traumatic hypothermia therapy? An ultrastructural perspective. *J Neurotrauma* 2005; 22:873–884
 188. Alam HB, Rhee P, Honma K, et al: Does the rate of rewarming from profound hypothermic arrest influence the outcome in a swine model of lethal hemorrhage? *J Trauma* 2006; 60:134–146
 189. Hildebrand F, van Griensven M, Giannoudis P, et al: Effects of hypothermia and re-warming on the inflammatory response in a murine multiple hit model of trauma. *Cytokine* 2005; 31:382–393
 190. Kawahara F, Kadoi Y, Saito S, et al: Slow rewarming improves jugular venous oxygen saturation during rewarming. *Acta Anaesthesiol Scand* 2003; 47:419–424
 191. Bissonnette B, Holtby HM, Davis AJ, et al: Cerebral hyperthermia in children after cardiopulmonary bypass. *Anesthesiology* 2000; 93:611–618
 192. Lavinio A, Timofeev I, Nortje J, et al: Cerebrovascular reactivity during hypothermia and rewarming. *Br J Anaesth* 2007; 99:237–244
 193. Sessler DI: Thermoregulatory defense mechanisms. *Crit Care Med* 2009; 37:S203–S210
 194. Frank SM, Fleisher LA, Breslow MJ, et al: Perioperative maintenance of normothermia reduces the incidence of morbid cardiac events. A randomized clinical trial. *JAMA* 1997; 277:1127–1134
 195. Frank SM, Beattie C, Christopherson R, et al: Unintentional hypothermia is associated with postoperative myocardial ischemia. The Perioperative Ischemia Randomized Anesthesia Trial Study Group. *Anesthesiology* 1993; 78:468–476
 196. Leslie K, Sessler DI: Perioperative hypothermia in the high-risk surgical patient. *Best Pract Res Clin Anaesthesiol* 2003; 17:485–498
 197. De Witte J, Sessler DI: Perioperative shivering: Physiology and pharmacology. *Anesthesiology* 2002; 96:467–484
 198. Van Zanten ARH, Polderman KH: Skin counterwarming to prevent shivering during therapeutic cooling: Blowing hot and cold? *Crit Care Med* 2009; 37:2106–2108
 199. Badjatia N, Strongilis E, Prescutti M, et al: Metabolic benefits of surface counterwarming during therapeutic temperature modulation. *Crit Care Med* 2009; 37:1893–1897
 200. Badjatia N, Strongilis E, Gordon E, et al: Metabolic impact of shivering during therapeutic temperature modulation. The Bed-side Shivering Assessment Scale. *Stroke* 2008; 39:3242–3247
 201. Thoresen M, Satas S, Loberg EM, et al: Twenty-four hours of mild hypothermia in unsedated newborn pigs starting after a severe global hypoxic-ischemic insult is not neuroprotective. *Pediatr Res* 2001; 50:405–411
 202. Tooley JR, Satas S, Porter H, et al: Head cooling with mild systemic hypothermia in anesthetized piglets is neuroprotective. *Ann Neurol* 2003; 53:65–72
 203. Bacher A: Effects of body temperature on blood gases. *Intensive Care Med* 2005; 31:24–27
 204. Polderman KH, van Zanten ARH, Girbes

- ARJ: The importance of magnesium in critically ill patients: A role in mitigating neurological injury and in the prevention of vasospasms. *Intensive Care Med* 2003; 29: 1202–1203
205. Soliman HM, Mercan D, Lobo SS, et al: Development of ionized hypomagnesemia is associated with higher mortality rates. *Crit Care Med* 2003; 31:1082–1087
206. Weisinger JR, Bellorin-Font E: Magnesium and phosphorus. *Lancet* 1998; 352:391–396
207. Fischer UM, Cox CS Jr, Laine GA, et al: Mild hypothermia impairs left ventricular diastolic but not systolic function. *J Invest Surg* 2005; 18:291–296
208. Goldberg LI: Effects of hypothermia on contractility of the intact dog heart. *Am J Physiol* 1958; 194:92–98
209. Suga H, Goto Y, Igarashi Y, et al: Cardiac cooling increases Emax without affecting relation between O₂ consumption and systolic pressure-volume area in dog left ventricle. *Circ Res* 1988; 63:61–71
210. Mikane T, Araki J, Suzuki S, et al: O₂ cost of contractility but not of mechanical energy increases with temperature in canine left ventricle. *Am J Physiol* 1999; 277:H65–H73
211. Lewis ME, Al-Khalidi AH, Townend JN, et al: The effects of hypothermia on human left ventricular contractile function during cardiac surgery. *J Am Coll Cardiol* 2002; 39:102–108
212. Rose JC, McDermott TF, Lilienfeld LS, et al: Cardiovascular function in hypothermic anesthetized man. *Circulation* 1957; 15: 512–517
213. Thoresen M, Whitelaw A: Cardiovascular changes during mild therapeutic hypothermia and rewarming in infants with hypoxic-ischemic encephalopathy. *Pediatrics* 2000; 106:92–99
214. Mattheussen M, Mubagwa K, Van Aken H, et al: Interaction of heart rate and hypothermia on global myocardial contraction of the isolated rabbit heart. *Anesth Analg* 1996; 82:975–981
215. Tveita T, Ytrehus K, Myhre ES, et al: Left ventricular dysfunction following rewarming from experimental hypothermia. *J Appl Physiol* 1998; 85:2135–2139
216. Polderman KH, Tjong Tjin Joe R, Peerderman SM, et al: Effects of artificially induced hypothermia on intracranial pressure and outcome in patients with severe traumatic head injury. *Intensive Care Med* 2002; 28: 1563–1567
217. Pozos RS, Danzl D: Human physiological responses to cold stress and hypothermia. In: *Medical Aspects of Harsh Environments*. Vol. 1. Textbooks of Military Medicine. Pandolf KB and Burr RE (Eds). Washington, DC, Walter Reed Army Medical Center Borden Institute, 2001, pp 351–382
218. Morgan ML, Anderson RJ, Ellis MA, et al: Mechanism of cold diuresis in the rat. *Am J Physiol* 1983; 244:F210–F216
219. Allen DE, Gellai M: Mechanisms for the diuresis of acute cold exposure: Role for vasopressin?. *Am J Physiol* 1993; 264: R524–R532
220. Sun Z: Genetic AVP deficiency abolishes cold-induced diuresis but does not attenuate cold-induced hypertension. *Am J Physiol Renal Physiol* 2006; 290:F1472–F1477
221. Sun Z, Zhang Z, Cade R: Renal responses to chronic cold exposure. *Can J Physiol Pharmacol* 2003; 81:22–27
222. Moat NE, Lamb RK, Edwards JC, et al: Induced hypothermia in the management of refractory low cardiac output states following cardiac surgery in infants and children. *Eur J Cardiothorac Surg* 1992; 6:579–584
223. Deakin CD, Knight H, Edwards JC, et al: Induced hypothermia in the postoperative management of refractory cardiac failure following paediatric cardiac surgery. *Anaesthesia* 1998; 53:848–853
224. Dalrymple-Hay MJ, Deakin CD, Knight H, et al: Induced hypothermia as salvage treatment for refractory cardiac failure following paediatric cardiac surgery. *Eur J Cardiothorac Surg* 1999; 15:515–518
225. Moriyama Y, Iguro Y, Shimokawa S, et al: Successful application of hypothermia combined with intra-aortic balloon pump support to low-cardiac-output state after open heart surgery. *Angiology* 1996; 47:595–599
226. Yahagi N, Kumon K, Watanabe Y, et al: Value of mild hypothermia in patients who have severe circulatory insufficiency even after intra-aortic balloon pump. *J Clin Anesth* 1998; 10:120–125
227. Hovdenes J, Laake JH, Aaberge L, et al: Therapeutic hypothermia after out-of-hospital cardiac arrest: Experiences with patients treated with percutaneous coronary intervention and cardiogenic shock. *Acta Anaesthesiol Scand* 2007; 51:137–142
228. Skulec R, Kovarnik T, Dostalova G, et al: Induction of mild hypothermia in cardiac arrest survivors presenting with cardiogenic shock syndrome. *Acta Anaesthesiol Scand* 2008; 52:188–194
229. Oddo M, Ribordy V, Feihl F, et al: Early predictors of outcome in comatose survivors of ventricular fibrillation and non-ventricular fibrillation cardiac arrest treated with hypothermia: A prospective study. *Crit Care Med* 2008; 36:2296–2301
230. Boddicker KA, Zhang Y, Zimmerman MB, et al: Hypothermia improves defibrillation success and resuscitation outcomes from ventricular fibrillation. *Circulation* 2005; 111:3195–3201
231. Rhee BJ, Zhang Y, Boddicker KA, et al: Effect of hypothermia on transthoracic defibrillation in a swine model. *Resuscitation* 2005; 65:79–85
232. Harada M, Honjo H, Yamazaki M, et al: Moderate hypothermia increases the chance of spiral wave collision in favor of self-termination of ventricular tachycardia/fibrillation. *Am J Physiol Heart Circ Physiol* 2008; 294:H1896–H1905
233. Bash SE, Shah JJ, Albers WH, et al: Hypothermia for the treatment of postsurgical greatly accelerated junctional ectopic tachycardia. *J Am Coll Cardiol* 1987; 10: 1095–1099
234. Balaji S, Sullivan J, Deanfield J, et al: Moderate hypothermia in the management of resistant automatic tachycardias in children. *Br Heart J* 1991; 66:221–224
235. Pfammatter JP, Paul T, Ziemer G, et al: Successful management of junctional tachycardia by hypothermia after cardiac operations in infants. *Ann Thorac Surg* 1995; 60:556–560
236. Frank SM, Satitpunwaycha P, Bruce SR, et al: Increased myocardial perfusion and sympathoadrenal activation during mild core hypothermia in awake humans. *Clin Sci (Lond)* 2003; 104:503–508
237. Nabel EG, Ganz P, Gordon JB, et al: Dilation of normal and constriction of atherosclerotic coronary arteries caused by the cold pressor test. *Circulation* 1988; 77:43–52
238. Schefold JC, Storm C, Joerres A, et al: Mild therapeutic hypothermia after cardiac arrest and the risk of bleeding in patients with acute myocardial infarction. *Int J Cardiol* 2008; 132:387–391
239. Tortorici MA, Kochanek PM, Poloyac SM: Effects of hypothermia on drug disposition, metabolism, and response: A focus of hypothermia-mediated alterations on the cytochrome P450 enzyme system. *Crit Care Med* 2007; 35:2196–2204
240. Kurz A, Sessler DI, Lenhardt R, et al: Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. *N Engl J Med* 1996; 334:1209–1215
241. Peterson K, Carson S, Carney N: Hypothermia treatment for traumatic brain injury: A systematic review and meta-analysis. *J Neurotrauma* 2008; 25:62–71
242. Schwab S, Georgiadis D, Berrouschot J, et al: High risk of pneumonia in stroke patients under hypothermia. Feasibility and safety of moderate hypothermia after massive hemispheric infarction. *Stroke* 2001; 32:2033–2035
243. De Jonge E, Schultz MJ, Spanjaard L, et al: Effects of selective decontamination of digestive tract on mortality and acquisition of resistant bacteria in intensive care: A randomised controlled trial. *Lancet* 2003; 362: 1011–1016
244. De Smet AM, Kluytmans JA, Cooper BS, et al: Decontamination of the digestive tract and oropharynx in intensive care patients. *New Engl J Med* 2009; 360:20–31