

## EDITORIAL



# Clinical examination: a trigger but not a substitute for hemodynamic evaluation

Daniel De Backer<sup>1\*</sup> and Antoine Vieillard-Baron<sup>2,3</sup>

© 2019 Springer-Verlag GmbH Germany, part of Springer Nature

Clinical examination is an essential part of the assessment of the critically ill patient. Among the different signs to look at, signs of tissue hypoperfusion and hemodynamic instability are particularly relevant. Several clinical signs can be used to detect tissue hypoperfusion. As an example, cerebral impairment, oliguria, capillary refill time, and mottling score are used in shock assessment [1]. Biological signs such as lactate have gained popularity, helping the clinician to suspect anaerobic metabolism, a potential consequence of severe hypoperfusion. Many of these signs are associated with outcome [2] and some may even be used to trigger resuscitation procedures [3].

While it is crucial to detect tissue hypoperfusion and its consequences, it is even more crucial to understand the cause of tissue hypoperfusion, whether cardiac output is sufficient, and what the cause of impaired cardiac output is. An important question is thus whether clinical and/or biological signs can predict cardiac index (CI). Tachycardia and hypotension mainly reflect the impaired cardiovascular status but it is obvious that neither tachycardia nor hypotension can reliably predict CI.

In this issue, Hiemstra et al. [4] prospectively evaluated in 783 critically ill patients whether 19 clinical and biological signs (among which heart rate, blood pressure, respiratory rate, cardiac and lung auscultation, mental state, urine output, and various signs of skin perfusion), evaluated in a protocolized manner, could predict CI. The authors first used linear regression to evaluate variables potentially associated with CI. Next they conducted

multivariate analyses to determine which combined clinical signs were identified with low or high CI.


In this single-center study, CI was measured by critical care echocardiography, using a transthoracic route, by experts in the field. The first interesting result is that among the 1212 eligible patients (unplanned ICU admission, expected to stay for at least 24 h), echocardiography could not be performed for technical reasons (chest drains, proning, wounds, etc.) in 80 patients (6.6%) and, when performed, was of insufficient quality to obtain CI in 292 other patients (24%). In trauma patients, factors such as a weight gain greater than 10%, a PEEP of at least 15 cmH<sub>2</sub>O, and chest tubes may contribute to the high failure rate (38%) of transthoracic echocardiography imaging [5]. Curiously, Hiemstra et al. [4] reported an unexpected similar failure rate in this study that included mostly medical patients. Even more surprising, Hiemstra et al. [6] previously mentioned in their preliminary report of the first 704 included patients that echocardiography was judged by experts to be of sufficient quality in 632 patients (90%), a much more usual failure rate. The second interesting finding is that CI was frequently low in this unselected population of ICU patients. A low CI (below 2.2 l/min/m<sup>2</sup>) was observed in 36% of the patients and 49% of the patients had a CI below 2.5 l/min/m<sup>2</sup>. Identifying these patients who potentially may benefit from hemodynamically targeted interventions is thus relevant.

Twelve clinical signs differed between patients with low and high CI. Seven signs reflecting impaired respiration, central circulation and skin perfusion were independently associated with CI. Five clinical signs (atrial fibrillation, systolic blood pressure below 90 mmHg, altered mental state, capillary refill time greater than 4.5 s at the sternum, and knee skin mottling) had a high specificity to detect a low CI. Interestingly, mean arterial

\*Correspondence: ddebacke@ulb.ac.be

<sup>1</sup> Department of Intensive Care, CHIREC Hospitals, Université Libre de Bruxelles, Brussels, Belgium

Full author information is available at the end of the article



pressure, central venous pressure, and urine output did not differ between the various CI level groups.

Unfortunately, the diagnostic accuracy was insufficient to identify low CI on the basis of these clinical signs. Even though most patients had altered signs of central hemodynamics and of tissue perfusion, the sensitivity of these signs was poor. Indeed, among the 368 patients with cardiac index below 2.2 L/min/m<sup>2</sup>, 34 had a systolic blood pressure below 90 mmHg and 52 a mean arterial pressure below 65 mmHg, 177 had cold extremities by subjective assessment, 38 had prolonged capillary refill time at the sternum, and 74 had severe skin mottling. The five highly specific signs mentioned above were present in only 10% of the patients with low CI. In addition to the very low the positive predictive value (ranging between 30% and 50% for most signs), the negative predictive value was only slightly better, ranging between 60% and 75%. This low positive predictive value is an important result as it is more representative of the physician's way of thinking in trying to detect a low CI rather than specificity. Indeed, it defines the probability that a low CI is present given the observed clinical signs. Furthermore, differences in the different variables between low and high CI were sometimes relatively minimal. For example, only 1 s in capillary refill time separated low and high CI [3.0 (3.0–4.0) s] in low CI patients and 2.0 (2.0–3.0) s in high CI patients.

While Hiemstra et al.'s results were somewhat expected from studies using either the pulmonary artery catheter [7] or transpulmonary thermodilution [8], the authors have to be congratulated as their rigorous analysis definitively gives us key information about the value of usual clinical signs.

What should we do next with these signs? We should not throw the baby with the bathwater. These signs are very useful to identify that there was a hemodynamic/perfusion issue. The included patients were very severely ill, half of them receiving norepinephrine, and very few patients (5%) had all clinical signs normal. Three quarters of the patients had abnormal skin perfusion signs, 68% abnormal signs of renal perfusion and one quarter signs of abnormal cerebral perfusion. Accordingly, these signs performed relatively well in indicating that there was a problem, but did not help so much in indicating what the source of the problem was.

We suggest, in accordance with the ESICM consensus report on hemodynamic management of shock [9], to use clinical signs to identify patients with potential hemodynamic issues, then to rapidly perform critical care echocardiography [10] in order to identify the type of shock and the source, if any, of the hemodynamic problem (Fig. 1).

In conclusion, clinical signs are important warning signals indicating that something is going wrong but these cannot reliably indicate whether cardiac index is low or high, nor indicate the source of the hemodynamic alteration. Physicians should not base their decisions/management solely on these clinical signs. The presence of altered clinical signs should prompt physicians to perform additional hemodynamic evaluations and in particular echocardiography.

#### Author details

<sup>1</sup> Department of Intensive Care, CHIREC Hospitals, Université Libre de Bruxelles, Brussels, Belgium. <sup>2</sup> Medical-surgical Intensive Care Unit, Ambroise Paré

University Hospital, APHP, Boulogne-Billancourt, France. <sup>3</sup> Université Versailles Saint Quentin, INSERM UMR1018, Team Kidney and Heart, CESP, Villejuif, France.

### Compliance with ethical standards

### Conflicts of interest

Daniel De Backer: Consultant to and material for studies by Edwards Lifesciences. Antoine Vieillard-Baron: Received grant from GSK for conducting clinical research and was member of the scientific advisory board for the study.

### Ethical approval

An approval by an ethics committee was not applicable.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 15 January 2019 Accepted: 17 January 2019

Published online: 24 January 2019

### References

1. Vincent JL, De Backer D (2013) Circulatory shock. *N Engl J Med* 369:1726–1734
2. Ait-Oufella H, Bakker J (2016) Understanding clinical signs of poor tissue perfusion during septic shock. *Intensive Care Med* 42:2070–2072
3. Hernandez G, Cavalcanti AB, Ospina-Tascon G, Zampieri FG, Dubin A, Hurtado FJ et al (2018) Early goal-directed therapy using a physiological holistic view: the ANDROMEDA-SHOCK—a randomized controlled trial. *Ann Intensive Care* 8:52
4. Hiemstra B, Koster G, Wiersema R, Hummel Y, van der Harst P, Snieder H et al (2019) The diagnostic accuracy of clinical examination for estimating cardiac output in critically ill patients: the Simple Intensive Care Studies-I. *Intensive Care Med*. <https://doi.org/10.1007/s00134-019-05527-y>
5. Cook CH, Praba AC, Beery PR, Martin LC (2002) Transthoracic echocardiography is not cost-effective in critically ill surgical patients. *J Trauma* 52:280–284
6. Hiemstra B, Eck RJ, Koster G, Wetterslev J, Perner A, Pettila V et al (2017) Clinical examination, critical care ultrasonography and outcomes in the critically ill: cohort profile of the Simple Intensive Care Studies-I. *BMJ Open* 7:e017170
7. Dawson NV, Connors AF, Speroff T, Kemka A, Shaw P, Arkes HR (1993) Hemodynamic assessment in managing the critically ill: is physician confidence warranted? *Med Decis Mak* 13:258–266
8. Perel A, Saugel B, Teboul JL, Malbrain ML, Belda FJ, Fernandez-Mondejar E et al (2016) The effects of advanced monitoring on hemodynamic management in critically ill patients: a pre and post questionnaire study. *J Clin Monit Comput* 30:511–518
9. Cecconi M, De Backer D, Antonelli M, Beale RJ, Bakker J, Hofer C et al (2014) Consensus on circulatory shock and hemodynamic monitoring. Task Force of the European Society of Intensive Care Medicine. *Intensive Care Med* 40:1795–1815
10. Expert round table on echocardiography in ICU (2014) International consensus statement on training standards for advanced critical care echocardiography. *Intensive Care Med* 2014(40):654–666

ORIGINAL



# The diagnostic accuracy of clinical examination for estimating cardiac index in critically ill patients: the Simple Intensive Care Studies-I

Bart Hiemstra<sup>1\*</sup> , Geert Koster<sup>1</sup> , Renske Wiersema<sup>1</sup> , Yoran M. Hummel<sup>2</sup>, Pim van der Harst<sup>2</sup> , Harold Snieder<sup>3</sup> , Ruben J. Eck<sup>1</sup> , Thomas Kaufmann<sup>4</sup> , Thomas W. L. Scheeren<sup>4</sup> , Anders Perner<sup>5,6</sup> , Jørn Wetterslev<sup>6,7</sup> , Anne Marie G. A. de Smet<sup>1</sup> , Frederik Keus<sup>1</sup>, Iwan C. C. van der Horst<sup>1</sup> and SICS Study Group<sup>1</sup>

© 2019 The Author(s)

## Abstract

**Purpose:** Clinical examination is often the first step to diagnose shock and estimate cardiac index. In the Simple Intensive Care Studies-I, we assessed the association and diagnostic performance of clinical signs for estimation of cardiac index in critically ill patients.

**Methods:** In this prospective, single-centre cohort study, we included all acutely ill patients admitted to the ICU and expected to stay > 24 h. We conducted a protocolised clinical examination of 19 clinical signs followed by critical care ultrasonography for cardiac index measurement. Clinical signs were associated with cardiac index and a low cardiac index ( $< 2.2 \text{ L min}^{-1} \text{ m}^2$ ) in multivariable analyses. Diagnostic test accuracies were also assessed.

**Results:** We included 1075 patients, of whom 783 (73%) had a validated cardiac index measurement. In multivariable regression, respiratory rate, heart rate and rhythm, systolic and diastolic blood pressure, central-to-peripheral temperature difference, and capillary refill time were statistically independently associated with cardiac index, with an overall  $R^2$  of 0.30 (98.5% CI 0.25–0.35). A low cardiac index was observed in 280 (36%) patients. Sensitivities and positive and negative predictive values were below 90% for all signs. Specificities above 90% were observed only for 110/280 patients, who had atrial fibrillation, systolic blood pressures  $< 90 \text{ mmHg}$ , altered consciousness, capillary refill times  $> 4.5 \text{ s}$ , or skin mottling over the knee.

**Conclusions:** Seven out of 19 clinical examination findings were independently associated with cardiac index. For estimation of cardiac index, clinical examination was found to be insufficient in multivariable analyses and in diagnostic accuracy tests. Additional measurements such as critical care ultrasonography remain necessary.

**Keywords:** Physical examination, Cardiac index, Haemodynamics, Sensitivity and specificity, Ultrasonography, Critical care, Intensive care unit, Prospective study

\*Correspondence: b.hiemstra01@umcg.nl

<sup>1</sup> Department of Critical Care, University of Groningen, University Medical Center Groningen, P.O. Box 30.001, 9700 RB Groningen, The Netherlands  
Full author information is available at the end of the article

## Introduction

Clinical examination is the first step to estimate cardiac index of critically ill patients to guide interventions and further diagnostic testing. Clinical signs such as altered consciousness, oliguria and a cold, clammy skin indicate organ hypoperfusion and are used to diagnose shock in critically ill patients [1]. If the initial clinical examination is inconclusive, further hemodynamic assessment with critical care ultrasonography (CCUS) is advocated [1, 2]. The evidence base of clinical examination is currently considered 'best practice' as there are scarce data on its diagnostic value, especially in comparison with newer, non-invasive bedside tools such as CCUS [1, 3].

Clinical examination for diagnosing shock is fast, easy to conduct and low in cost, yet its diagnostic accuracy is questioned [3–5]. Particularly, physicians seem insufficiently capable of diagnosing a low cardiac index purely based on their clinical examination [6–9]. Previous studies scarcely specified their methods of clinical examination in terms of variables collected and definitions employed, leaving variability at the physician's discretion and making these studies difficult to reproduce [7, 9–15]. Cardiac index was measured only in small samples of selected patients who failed to respond to initial therapy or in whom clinical examination alone was deemed insufficient, so the accuracy of clinically estimated cardiac index was biased by definition [1, 12–15].

The value of clinical signs for estimating cardiac index remains to be established in a large, consecutively recruited cohort of critically ill patients. Our aim was to study the diagnostic performance of clinical examination in a twostep approach: to establish (1) which combination of clinical examination findings are independently associated with cardiac index and (2) the performance of clinical signs to diagnose a low or high cardiac index. The Simple Intensive Care Studies-I (SICS-I) was designed to assess the ability of clinical examination to estimate cardiac index and to identify patients with a low or high cardiac index [16].

## Methods

### Design, setting and patients

We conducted the prospective, observational, single-centre SICS-I following a pre-published protocol and statistical analysis plan (SAP; [clinicaltrials.gov](https://clinicaltrials.gov); NCT02912624) [17]. All consecutive patients admitted to the intensive care unit (ICU) of the University Medical Center Groningen (UMCG) between 27 March 2015 and 22 July 2017 were considered eligible. We included patients who were aged 18 years or older, had an unplanned ICU admission and were expected to stay for at least 24 h. We

## Take-home message

Abnormal clinical signs and a low cardiac index were often present in acutely admitted ICU patients and seven out of nineteen clinical examination findings were independently associated with cardiac index. To estimate cardiac index, clinical examination performs insufficiently in both multivariable analyses and in diagnostic accuracy tests; additional measurements such as critical care ultrasonography remain necessary.

excluded patients if their ICU admission was planned preoperatively, if acquiring research data interfered with clinical care (e.g., mechanical circulatory support) and if informed consent was not provided (e.g., refusal, serious language barrier). In unresponsive patients, informed consent was first obtained from the legal representatives. Consent for use of the study data was asked at a later time if the patient recovered consciousness. If the patient died before consent was obtained, the study data was used, and the legal representatives were informed on the study. The study was approved by the local institutional review board (METc M15.168207).

All patients were included within the first 24 h of their ICU admission and underwent clinical examination, immediately followed by CCUS. Patients were included by medical research interns and PhD students who had received a focused CCUS training course given by experienced cardiologist-intensivists (protocol in Supplements 1) [16]. These researchers were not involved in patient care, and their findings were not revealed to the patients' caregivers.

### Clinical examination

All clinical examinations were standardised and cut-off values for abnormal clinical signs predefined in the protocol ([clinicaltrials.gov](https://clinicaltrials.gov); NCT02912624). We recorded in total 19 clinical signs per patient (e-Table 1). Respiratory rate, heart rate and rhythm, arterial blood pressures and central venous pressures were recorded from the bedside monitor. Patients were auscultated for the presence of cardiac murmurs and crepitations. Clinical signs reflecting organ perfusion were obtained from the three organs readily accessible to clinical examination: cerebral (mental status), renal (urine output) and skin perfusion (capillary refill time (CRT), central-to-peripheral temperature difference ( $\Delta T_{c-p}$ ) and skin mottling). Mental status was assessed with the AVPU scale, which consists of the categories 'Alert', 'responsive to Voice', 'responsive to Pain' and 'Unresponsive' and was not scored in patients who were receiving sedative drugs or who were admitted after a trauma. Urine output was scored 1 and 6 h prior to the clinical examination, adjusted for body weight, and

considered decreased if  $<0.5 \text{ mL kg}^{-1} \text{ h}^{-1}$ . CRT was the time for skin colour to fully return after applying firm pressure at the sternum, index finger, and knee for 15 s and considered prolonged if  $>4.5 \text{ s}$  [18].  $\Delta T_{c-p}$  was the difference between central temperature measured by a bladder thermistor catheter and peripheral temperature measured by a skin probe on the big toe and dorsum of the foot and considered abnormal if  $>7^\circ \text{C}$  [19]. The degree of skin mottling was rated at the knee according to a score from 0 to 5, where 0–1 was regarded as mild, 2–3 as moderate and 4–5 as severe mottling [20].

### Outcome definition

Cardiac index was measured by transthoracic echocardiography using the Vivid-S6 system (General Electric, Horton, Norway) with cardiac probe M3S or M4S, and with default cardiac imaging setting. The parasternal long axis (PLAX) was used to measure the left ventricular outflow tract (LVOT) diameter. In the apical five chamber (AP5CH) view, a pulse wave Doppler signal in the LVOT was used to measure the velocity time integral (VTI); three VTIs were traced when the heart rhythm was regular and eight VTIs when the heart rhythm was irregular. Cardiac output was calculated using an established formula and was adjusted for body surface area, i.e. cardiac index, because it allowed us to compare patients with different body dimensions [21]. We tested the diagnostic accuracy of clinical signs on four frequently used cut-off values: below 2.2 and 2.5, and above 4.0 and  $4.5 \text{ L min}^{-1} \text{ m}^{-2}$  [10, 11, 22, 23].

All CCUS images were validated and each cardiac index measured by experts from an independent imaging core laboratory (Groningen Imaging Core Laboratory, [www.g-icl.com](http://www.g-icl.com)). These experts were blinded for the clinical examination findings. The most-often used modality for measuring cardiac index in the literature currently is the transpulmonary thermodilution method [24, 25]. We chose CCUS because it is non-invasive, its use is advocated by guidelines, and ensured high-quality measurements throughout validation by experts who followed recent guidelines [26].

### Sample size and missing data

No previous research has studied clinical examination in a cohort of consecutively included critically ill patients. Therefore, we estimated our sample size based on the number of acute ICU admissions annually ( $n=1500$ ) and estimated that half would fulfil the inclusion criteria. When our sample size exceeded 1000 patients, we calculated a potentially detectable difference using CRT as an example; we were able to detect a  $0.10 \text{ L min}^{-1} \text{ m}^{-2}$  increase in cardiac index for each

second of CRT increase with a power of 100% and alpha of 0.015 [17]. Missing clinical examination values were imputed using multiple imputations (20 times) as these were considered missing at random. Regression coefficients of our final models were averaged using Rubin's formula [27]. Central venous pressure was missing in 822 (76%) patients and was therefore excluded from the analyses and imputations. Following our SAP, we also imputed missing cardiac index values based on validated LVOT diameters and VTIs (Supplements 2; sensitivity analyses).

### Analytical approach

The aims of our analyses were twofold: first, we conducted a least-squares linear regression analysis to identify the clinical examination findings that were independently associated with cardiac index as a continuous variable. Second, we calculated diagnostic test accuracies for each clinical sign and conducted multivariable logistic regression analyses to determine which combined clinical signs were independently associated with a low ( $<2.2$  and  $<2.5 \text{ L min}^{-1} \text{ m}^{-2}$ ) or high ( $>4.0$  and  $>4.5 \text{ L min}^{-1} \text{ m}^{-2}$ ) cardiac index.

### Multivariable model development and validation

We conducted a linear regression analysis when using cardiac index as a continuous variable and logistic regression analysis when using a dichotomised cardiac index. We used a  $p<0.25$  threshold for inclusion in the multivariable models, which was constructed using forward stepwise regression by adding blocks of variables. Noradrenaline infusion rate was added as a confounder to our multivariable models on a theory-driven basis. For the linear regression model, normality of the residuals was assessed with kernel density plots and multicollinearity was checked with variance inflation factors. The model was internally validated with bootstrap sampling, in which we tested whether each predictor was significant in at least 80 of the 100 bootstrap replications. For the logistic regression model, we assessed calibration with the Hosmer–Lemeshow test and evaluated discrimination using receiver operating characteristic (ROC)-curves.

### Diagnostic test accuracy

The following diagnostic test accuracies were calculated: sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and overall accuracy. Furthermore, we calculated likelihood ratios (LRs), which represent the increase of probability of a low cardiac index when a clinical sign is abnormal (positive LR) or normal (negative LR) [28]. Following specific Standards for Reporting of Diagnostic Accuracy Studies of medical

history and physical examination [29], we summarised all  $2 \times 2$  tables in an overview table (e-Tables 7a and 8a). Clinical signs reflecting central circulation were also dichotomised to test their diagnostic performance.

### Subgroup analyses

We conducted two planned subgroup analyses on the diagnostic test accuracies. First, we stratified the study population by noradrenaline administration. Second, we divided patients by primary reasons of ICU admittance as these underlying pathologies may have influenced cardiac index: i.e., acute liver failure or post orthotopic liver transplantation (OLT), heart failure, septic shock, cardiac arrest, and central nervous system (CNS) pathologies.

### Statistical significance

We conducted our analyses with Stata version 15.1 software (StataCorp, College Station, TX, USA) and followed a published SAP [17]. Cardiac index was one of the six primary outcomes tested in our cohort and, therefore, we adjusted for multiple hypothesis testing [16, 30]. We refer to our SAP for more details (Supplements 3), but in short a  $p$  value of 0.015 indicated statistical significance and  $p$  values between 0.015 and 0.05 indicated suggestive significance with an increased family-wise error rate [17]. Accordingly, we presented our primary analysis with 98.5% CI, all secondary analyses with 95% CIs, and discussed any suggestively significant findings based on the  $p$  value and results from the bootstrap replications.

## Results

### Study population

A total of 1212 patients were eligible, of which 137 patients were not included because the desired CCUS window was obstructed by chest drains, wounds, emphysema, or prone positioning ( $n=80$ ); routine clinical care such as acute surgery or angiography did not allow patient inclusion ( $n=40$ ); or other reasons such as colonisation with multiresistant bacteria ( $n=17$ ), leaving 1075 patients for inclusion. Of these, 292 (27%) patients were excluded because one or both of the CCUS views were of insufficient quality as scored by the independent echocardiography core laboratory, resulting in a total of 783 (73%) patients included in the current analyses (Fig. 1). One-third of the patients were admitted after acute or complicated surgery, and the most common admission diagnoses were of cardiovascular or respiratory origin (Table 1). The median time from ICU admission to inclusion was 15 h (IQR 8–20 h).

The mean cardiac index was  $2.65 \pm 0.93 \text{ L min}^{-1} \text{ m}^{-2}$ . Table 2 shows that 12 clinical examination findings differed significantly over five cardiac index categories. Clinical signs reflecting normal skin, renal and cerebral perfusion were present in 36 (5%) patients. Abnormal clinical signs were observed in 598 (76%) patients for skin perfusion, 536 (68%) patients for renal perfusion, and 204 (26%) patients for cerebral perfusion. Both cerebral and renal perfusion were abnormal in 141 (18%) patients; skin, renal and cerebral perfusion were abnormal in 103 (13%) patients.


### Clinical examination associated with cardiac index

Univariable analyses in the overall population showed that heart rate, systolic blood pressure and all clinical signs reflecting abnormal organ perfusion were statistically significantly associated with cardiac index (supplements; e-Table 2). The strongest association was found for heart rate with an  $R^2$  of 0.15. At the mean heart rate of 87 beats per minute, the mean cardiac index was  $2.1 \text{ L min}^{-1} \text{ m}^{-2}$  with an individual 95% CI that ranged from 0.87 to  $4.1 \text{ L min}^{-1} \text{ m}^{-2}$  (e-Fig. 1), reflecting the inaccuracy of a univariable prediction. In multivariable linear regression seven clinical signs, i.e., respiratory rate, heart rate and rhythm, systolic and diastolic blood pressure,  $\Delta\text{Tc-p}$ , and CRT were independently associated with cardiac index (e-Table 3). The presence of cardiac murmurs had a  $p$  value  $>0.015$  and was statistically significant in less than 80 of the 100 bootstrap replications. The multivariable model had an  $R^2$  of 0.30 (98.5% CI 0.26–0.37). Other model diagnostics and sensitivity analyses are presented in the supplements (e-Figs. 2–4 and e-Tables 4, 5).

### Diagnostic test performance of clinical signs

#### Accuracy of single clinical signs

A cardiac index below  $2.2 \text{ L min}^{-1} \text{ m}^{-2}$  was observed in 280 (36%) patients. The performance of the 19 clinical signs to diagnose a low cardiac index showed that none had sensitivities, PPVs and NPVs that exceeded 90% (Table 3). A specificity above 90% was found when patients had atrial fibrillation, a systolic blood pressure below 90 mmHg or a diastolic blood pressure below 45 mmHg, an altered consciousness without sedation (i.e., patients who reacted to a pain stimulus only or who were unresponsive), a CRT of  $>4.5 \text{ s}$  at the sternum, or had mottling over the knee (Fig. 2). One of these five clinical signs was abnormal in 110 (39%) of the 280 patients with a cardiac index below  $2.2 \text{ L min}^{-1} \text{ m}^{-2}$ . A



**Fig. 1** Flow diagram of the Simple Intensive Care Studies-I (SICS-I). *ICU* intensive care unit, *CCUS* critical care ultrasonography, *PLAX* parasternal long axis, *AP5CH* apical 5-chamber

low cardiac index was over 1.5 times more likely when patients had atrial fibrillation, subjectively cold feet,

or a CRT at the sternum or index finger of more >4.5 s (Table 3; positive likelihood ratio).

**Table 1 Clinical characteristics**

Variable	All patients N = 783
Age (years)	61 ± 15
Male gender	484 (62%)
Mechanical ventilation	438 (56%)
PEEP (cm H <sub>2</sub> O)	7 (5, 8)
Admission type	
Medical	544 (69%)
Acute surgery	207 (26%)
Complications of surgery	32 (4%)
Admission diagnosis by organ system	
Cardiovascular	228 (29%)
Gastrointestinal	120 (15%)
Genitourinary	20 (3%)
Haematological	14 (2%)
Metabolic	16 (2%)
Musculoskeletal/skin	11 (1%)
Neurological	116 (15%)
Respiratory	173 (22%)
Transplant	29 (4%)
Trauma	55 (7%)
Subgroups	
Acute heart failure	49 (11%)
Cardiac arrest	94 (21%)
CNS failure	115 (25%)
Liver failure	42 (9%)
Sepsis	151 (33%)

PEEP positive end-expiratory pressure

The diagnostic performance was also assessed for cardiac index cut-off values below 2.5 and above 4.0 and 4.5 L min<sup>-1</sup> m<sup>-2</sup>, and results are presented in the supplements (e-Fig. 6, e-Tables 7 and 8). The accuracy of diagnostic test performance was comparable in all 1075 patients with an imputed cardiac index (e-Tables 9, 10).

#### Accuracy of combined clinical signs

The PPV of a cardiac index below 2.2 L min<sup>-1</sup> m<sup>-2</sup> was higher when patients had lower heart rates and lower systolic blood pressures, and when more organs showed signs of hypoperfusion (Fig. 3). Multivariable logistic regression analyses adjusted for noradrenaline infusion rate showed that respiratory rate, heart rate, atrial fibrillation, systolic and diastolic blood pressure, CRT at the sternum and  $\Delta Tc-p$  were independently associated with a cardiac index below 2.2 L min<sup>-1</sup> m<sup>-2</sup>. The model had an area under the ROC of 0.74 (95% CI 0.70–0.78; e-Table 6).

These clinical signs correctly classified 556 (71%) of the 783 patients into a low or normal cardiac index.

#### Subgroup analyses

We conducted two predefined subgroup analyses according to noradrenaline use and primary reason for admission. The subgroups were either small ( $n < 50$ ) or the diagnostic test performance showed little improvement compared to the entire cohort (i.e.,  $< 5\%$ ; e-Fig. 7).

#### Discussion

In the SICS-I, we studied the association of 19 clinical examination findings with cardiac index and assessed their performance in diagnosing a low cardiac index in a consecutive cohort of acutely ill patients admitted to the ICU. We observed that only 5% of the patients had clinical signs indicating normal cerebral, renal and skin perfusion. A little over one-third of all patients had a cardiac index below 2.2 L min<sup>-1</sup> m<sup>-2</sup> and clinical examination findings were more often abnormal when cardiac index was low. Seven clinical examination findings which reflect respiration, central circulation and skin hypoperfusion were independently associated with cardiac index. Five clinical signs with a high specificity may be used to conclude that a low cardiac index is likely. Nevertheless, the performance of clinical examination in both multivariable analyses and diagnostic tests is insufficient for estimating cardiac index.

This study is the first to associate a broad set of clinical examination variables with cardiac index measured by CCUS. The multivariable regression analyses of our primary outcome showed that 30% of the variance was explained by the model, implicating that clinical examination is insufficiently capable of estimating cardiac index. Compared to smaller studies which assessed one, two or three clinical examination variables, the associations for heart rate, respiratory rate and  $\Delta Tc-p$  gradients were similar [31–35]. In contrast, by including all previously clinical signs, we showed that a CRT measured at the sternum and not peripheral CRT is independently associated with cardiac index [20].

Studies that reported on the physician's educated guess of cardiac output lacked the design to assess the diagnostic performance of each clinical sign, obtained according to strict definitions. Using multivariable logistic regression, we correctly classified 556 (71%) patients into a low or normal cardiac index, which contrasts with others who found agreements of 50–60% [8–11, 13–15]. One other study evaluated three standardised clinical signs and also found that an abnormality of all three clinical signs

**Table 2 Clinical examination and cardiac index**

Variable	All patients N = 783	CI < 1.8 N = 138	CI 1.8–2.2 N = 142	CI 2.2–2.5 N = 105	CI 2.5–4.0 N = 332	CI > 4.0 N = 66	p value
Central circulation							
Respiratory rate (per minute)	18 ± 6	18 ± 6	18 ± 5	18 ± 5	18 ± 5	19 ± 7	0.34
Heart rate (beats per minute)	87 ± 21	77 ± 19	84 ± 21	82 ± 18	90 ± 19	101 ± 22	< 0.001
Atrial fibrillation	56 (7%)	11 (8%)	15 (11%)	7 (7%)	18 (5%)	5 (8%)	0.38
Systolic blood pressure (mmHg)	120 ± 24	113 ± 23	119 ± 21	117 ± 24	123 ± 25	122 ± 26	< 0.001
Diastolic blood pressure (mmHg)	60 ± 12	61 ± 11	60 ± 10	59 ± 13	59 ± 11	59 ± 12	0.47
Mean arterial pressure (mmHg)	79 ± 14	78 ± 14	79 ± 12	77 ± 16	80 ± 15	79 ± 15	0.37
Central venous pressure (mmHg)	9 (5, 13)	9 (5, 14)	10 (7, 12)	13 (6, 16)	8 (5, 12)	10 (6, 13)	0.45
Cardiac murmurs	75 (10%)	5 (4%)	13 (9%)	12 (11%)	34 (10%)	11 (17%)	0.039
Crepitations	115 (15%)	26 (19%)	10 (7%)	13 (12%)	52 (16%)	14 (21%)	0.022
Noradrenaline	361 (46%)	73 (53%)	72 (51%)	48 (46%)	136 (41%)	32 (48%)	0.11
Organ perfusion							
Consciousness							
Alert	254 (32%)	33 (24%)	39 (27%)	30 (29%)	127 (38%)	25 (38%)	< 0.001
Reacting to voice	154 (20%)	19 (14%)	24 (17%)	22 (21%)	73 (22%)	16 (24%)	
Reacting to pain	67 (9%)	10 (7%)	20 (14%)	7 (7%)	25 (8%)	5 (8%)	
Unresponsive	308 (39%)	76 (55%)	59 (42%)	46 (44%)	107 (32%)	20 (30%)	
Urine output (mL kg <sup>-1</sup> h <sup>-1</sup> )	0.56 (0.31, 1.11)	0.53 (0.26, 1.15)	0.54 (0.30, 1.00)	0.50 (0.25, 1.03)	0.61 (0.35, 1.14)	0.63 (0.35, 1.43)	0.31
Urine output (mL kg <sup>-1</sup> 6 h <sup>-1</sup> )	0.65 (0.37, 1.16)	0.64 (0.37, 1.06)	0.58 (0.34, 1.08)	0.56 (0.36, 1.20)	0.70 (0.41, 1.19)	0.68 (0.34, 1.33)	0.34
Central temperature (°C)	36.9 ± 0.9	36.6 ± 0.9	37.0 ± 0.9	36.9 ± 0.9	37.0 ± 0.9	37.2 ± 1.0	< 0.001
ΔTc–p, dorsum foot (°C)	7.5 ± 3.1	8.6 ± 3.1	8.2 ± 3.0	7.5 ± 3.0	7.0 ± 3.0	6.4 ± 3.1	< 0.001
ΔTc–p, big toe (°C)	9.1 ± 3.6	10.1 ± 3.2	9.9 ± 3.4	9.3 ± 3.5	8.5 ± 3.6	7.5 ± 3.8	< 0.001
Cold extremities, subjective	285 (36%)	77 (56%)	55 (39%)	43 (41%)	93 (28%)	17 (26%)	< 0.001
Capillary refill time sternum (s)	3.0 (2.0, 3.0)	3.0 (3.0, 4.0)	3.0 (2.0, 3.0)	3.0 (2.0, 3.0)	3.0 (2.0, 3.0)	2.0 (2.0, 3.0)	< 0.001
Capillary refill time finger (s)	3.0 (2.0, 4.0)	4.0 (2.0, 5.0)	3.0 (2.0, 4.0)	3.0 (2.0, 4.0)	2.0 (2.0, 3.0)	2.0 (2.0, 4.0)	< 0.001
Capillary refill time knee (s)	3.0 (2.0, 4.0)	4.0 (3.0, 5.0)	3.0 (3.0, 5.0)	3.0 (2.0, 4.0)	3.0 (2.0, 4.0)	3.0 (2.0, 4.0)	< 0.001
Skin mottling severity <sup>a</sup>							
Mild (0–1)	543 (69%)	80 (58%)	96 (68%)	78 (74%)	243 (73%)	46 (70%)	0.096
Moderate (2–3)	217 (28%)	52 (38%)	43 (30%)	24 (23%)	79 (24%)	19 (29%)	
Severe (4–5)	23 (3%)	6 (4%)	3 (2%)	3 (3%)	10 (3%)	1 (2%)	

CI/ cardiac index in L min<sup>-1</sup> m<sup>-2</sup>, ΔTc–p, central-to-peripheral temperature difference

<sup>a</sup> Mottling was scored according to Ait-Oufella et al. [20]

have high specificities and low sensitivities to diagnose a low cardiac index [36]. In the above mentioned studies, cardiac index was obtained with transpulmonary thermodilution, compared to echocardiography in our study; evidence indicates that these two techniques may not be interchangeable [37].

### Implications for practice

Clinical signs are important warning signals of a deteriorating patient condition, but cannot reliably indicate whether the cardiac index is low, normal or high, or the underlying cause. Our results show both the relevance and limitations of clinical examination: five clinical signs had a specificity above 90%, yet the PPV is much more pertinent than specificity in the clinical process, as it reflects the probability that a low CI is present given

**Table 3 Diagnostic performance of clinical signs for estimating a cardiac index below 2.2 L·min<sup>-1</sup>·m<sup>-2</sup>**

Clinical sign	Abnormal  N (%)	Diagnostic performance in %					Likelihood and odds ratios		
		Sensitivity	Specificity	PPV	NPV	Accuracy	Positive	Negative	Odds
Central circulation									
Respiratory rate > 22 pm	550 (70%)	77 (72–82)	34 (29–38)	39 (37–41)	73 (67–77)	49 (46–53)	1.16 (1.06–1.27)	0.68 (0.53–0.87)	1.71 (1.22–2.38)
Heart rate < 100 bpm	571 (73%)	82 (77–86)	32 (28–36)	40 (36–44)	76 (70–82)	50 (46–53)	1.20 (1.11–1.31)	0.57 (0.43–0.75)	2.11 (1.48–3.02)
Atrial fibrillation	56 (7%)	9 (6–13)	94 (92–96)	46 (33–60)	64 (60–67)	64 (61–67)	1.56 (0.94–2.58)	0.96 (0.92–1.01)	1.61 (0.94–2.77)
SBP < 90 mmHg	56 (7%)	6 (4–10)	92 (90–95)	32 (20–46)	62 (58–65)	63 (60–66)	0.85 (0.50–1.46)	1.01 (0.97–1.05)	0.84 (0.47–1.49)
DBP < 45 mmHg	40 (5%)	5 (3–8)	95 (92–96)	33 (19–49)	64 (61–68)	62 (59–66)	0.86 (0.45–1.65)	1.01 (0.97–1.04)	0.86 (0.44–1.67)
MAP < 65 mmHg	97 (12%)	11 (8–16)	87 (84–90)	33 (24–43)	64 (60–68)	60 (57–63)	0.88 (0.59–1.32)	1.02 (0.96–1.07)	0.87 (0.56–1.36)
Cardiac murmurs	75 (10%)	6 (4–10)	89 (86–91)	24 (15–35)	63 (59–67)	59 (56–63)	0.57 (0.34–0.94)	1.06 (1.01–1.10)	0.54 (0.31–0.93)
Crepitations	115 (15%)	13 (9–17)	84 (81–87)	31 (23–41)	64 (60–67)	59 (55–62)	0.82 (0.57–1.18)	1.03 (0.97–1.10)	0.79 (0.52–1.21)
Organ perfusion									
Consciousness									
Reacting to voice	121 (15%)	11 (8–15)	82 (79–85)	26 (18–34)	62 (59–66)	57 (53–60)	0.62 (0.24–0.91)	1.08 (1.02–1.15)	0.57 (0.37–0.88)
Reacting to pain	33 (4%)	5 (3–8)	96 (94–98)	42 (26–61)	65 (61–68)	64 (60–67)	1.32 (0.67–2.60)	0.99 (0.96–2.60)	1.34 (0.67–2.68)
Unresponsive	50 (6%)	7 (4–10)	94 (91–96)	38 (25–53)	63 (59–66)	63 (60–66)	1.10 (0.63–1.91)	0.99 (0.96–1.03)	1.11 (0.62–1.99)
Urine output < 0.5 ml·kg <sup>−1</sup> ·h <sup>−1</sup>									
Over 1 h	345 (44%)	47 (41–53)	58 (53–62)	38 (33–44)	66 (62–71)	54 (50–57)	1.11 (0.95–1.31)	0.92 (0.80–1.05)	1.21 (0.91–1.63)
Over 6 h	288 (37%)	40 (34–46)	65 (61–69)	39 (33–44)	66 (62–70)	56 (52–59)	1.13 (0.93–1.36)	0.93 (0.83–1.04)	1.21 (0.90–1.63)
Skin temperature cold <sup>a</sup>									
ΔTc–p, dorsum foot	402 (51%)	78 (72–82)	40 (35–44)	42 (37–46)	76 (70–81)	53 (50–57)	1.28 (1.17–1.41)	0.57 (0.45–0.72)	2.25 (1.62–3.14)
ΔTc–p, big toe	521 (67%)	65 (59–71)	56 (52–61)	45 (40–50)	74 (70–79)	59 (56–63)	1.49 (1.30–1.69)	0.62 (0.52–0.74)	2.39 (1.77–3.23)
Cold subjective	285 (36%)	47 (41–53)	70 (65–74)	46 (40–52)	70 (66–74)	62 (58–65)	1.55 (1.29–1.86)	0.76 (0.67–0.86)	2.04 (1.51–2.76)
CRT prolonged <sup>b</sup>									
Sternum	62 (8%)	10 (7–15)	93 (91–95)	47 (34–60)	65 (62–69)	64 (60–67)	1.58 (0.98–2.54)	0.96 (0.92–1.00)	1.65 (0.98–2.76)
Finger	147 (19%)	25 (20–31)	85 (82–88)	48 (40–57)	67 (63–71)	64 (60–67)	1.68 (1.26–2.24)	0.88 (0.81–0.95)	1.91 (1.33–2.74)
Knee	199 (25%)	33 (27–39)	79 (75–82)	46 (39–53)	68 (64–72)	62 (59–66)	1.54 (1.22–1.96)	0.85 (0.78–0.94)	1.81 (1.31–2.51)
Skin mottling severity <sup>c</sup>									
Over the knee (≥ 3)	92 (12%)	16 (12–21)	91 (88–93)	49 (38–60)	66 (62–70)	64 (61–67)	1.72 (1.17–2.52)	0.93 (0.87–0.98)	1.86 (1.20–2.87)
Severe (≥ 4)	23 (3%)	3 (2–6)	97 (95–99)	39 (20–62)	64 (61–68)	64 (60–67)	1.15 (0.51–2.63)	1.00 (0.97–1.02)	1.16 (0.51–2.66)

Values between parentheses present 95% confidence intervals

SBP systolic blood pressure, DBP diastolic blood pressure, MAP mean arterial pressure, PPV positive predictive value, NPV negative predictive value, ΔTc–p central-to-peripheral temperature difference, CRT capillary refill time


<sup>a</sup> ΔTc–p was considered cold if > 7 °C

<sup>b</sup> CRT was considered prolonged if > 4.5 s

<sup>c</sup> Mottling was scored according to Ait-Oufella et al. [19]

that the clinical signs indicate hypoperfusion. The PPV of single clinical signs (Fig. 2) is too low in the majority of patients. PPVs above 90% were observed in a minority of patients with several abnormal clinical signs (Fig. 3). It is unacceptable for a diagnostic test to miss a low cardiac

index in one-quarter of cases, or even a larger proportion when diagnosing a high cardiac index (e-Table 6). If physicians immediately perform CCUS, they could diagnose cardiac failure within a few minutes and perform



the adequate supportive and therapeutic measures. In the acute setting, this is superior to the collection of urine over 1 h.

The results of our study imply that physicians should not rely solely on clinical signs for their decision making. Although the clinical signs atrial fibrillation, systolic blood pressure below 90 mmHg, altered consciousness, CRT of  $>4.5$  s at the sternum and skin mottling over the knee make a low cardiac index very likely, we advocate ultrasonography for additional evaluation in these patients. For patients with normal clinical signs additional evaluation should be performed on indication. Therewith, we verify two important statements of the current circulatory shock guidelines for which no evidence base existed (i.e., ‘best practice’) [1].

### Strengths and limitations

The clinical examination findings measured in the SICS-I were prone to confounding by interventions. The administration of noradrenaline may have influenced both cardiac index and organ perfusion variables, which is why we added it as a confounder in our multivariable models. The observational design of our study does not allow for conclusions if the measured cardiac index is sufficient for the individual patient. CCUS is focused on obtaining few potentially important measurements and therewith valve pathologies can be overseen. Additional CCUS measures such as ejection fraction and end-diastolic volumes are needed to distinguish between a compensating or a failing heart, but this was beyond the scope of our current research question.

The SICS-I included acutely ill patients admitted to the ICU of a tertiary referral hospital. Compared to others who selected patients with sepsis or myocardial

SBP (mmHg)	No. skin signs abnormal				Heart rate (bpm)
	0	1	2	3	
>140	0%	0%	0%	0%	> 120
120-140	0%	0%	0%	0%	
100-120	0%	33%	50%	17%	
<100	33%	25%	50%	0%	
>140	11%	14%	33%	100%	100 - 120
120-140	0%	24%	31%	29%	
100-120	6%	33%	18%	33%	
<100	0%	33%	45%	25%	
>140	23%	13%	40%	0%	80 - 100
120-140	38%	45%	36%	38%	
100-120	22%	37%	36%	40%	
<100	18%	24%	17%	58%	
>140	13%	36%	55%	46%	60 - 80
120-140	23%	44%	58%	38%	
100-120	33%	31%	48%	62%	
<100	0%	42%	83%	91%	
>140	40%	17%	67%	67%	< 60
120-140	40%	60%	100%	100%	
100-120	83%	50%	75%	80%	
<100	100%	100%	100%	100%	
	0	1	2	3	
Number of skin signs abnormal					
Abnormal skin signs were counted as any prolonged capillary refill time, any cold peripheral temperature and moderate skin mottling					

**Fig. 3** Positive predictive values of combined clinical signs to diagnose a cardiac index  $< 2.2 \text{ L min}^{-1} \text{ m}^{-2}$ . The figure displays the positive predictive value for combinations of the clinical signs heart rate, systolic blood pressure, and number of skin signs showing hypoperfusion (i.e., capillary refill time, cold peripheral temperature, and skin mottling). The figure shows that the PPV increases when patients have lower heart rates combined with lower systolic blood pressures and more signs of skin hypoperfusion. SBP systolic blood pressure

infarction, we studied the value of clinical examination in a large, unselected cohort. Since PPV and NPV values are dependent on the baseline prevalence of low cardiac index, we selected all consecutive patients to maximise representativeness of our cohort to the ICU. However, our study requires external validation before its findings can be applied to other ICUs. Furthermore, our study did not test the inter- and intra-observer agreement of cardiac index measured with CCUS; we will address the inter-observer agreement (researcher versus expert) separately.

## Conclusion

Abnormal clinical signs and a low cardiac index were often present in acutely admitted ICU patients, and seven

out of 19 clinical examination findings were independently associated with cardiac index. To estimate cardiac index, clinical examination performs insufficiently in both multivariable analyses and in diagnostic accuracy tests. Additional measurements such as critical care ultrasonography remain necessary.

## Electronic supplementary material

The online version of this article (<https://doi.org/10.1007/s00134-019-05527-y>) contains supplementary material, which is available to authorized users.

## Author details

<sup>1</sup> Department of Critical Care, University of Groningen, University Medical Center Groningen, P.O. Box 30.001, 9700 RB Groningen, The Netherlands.

<sup>2</sup> Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. <sup>3</sup> Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. <sup>4</sup> Department of Anesthesiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. <sup>5</sup> Department of Intensive Care, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark. <sup>6</sup> Centre for Research in Intensive Care, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark. <sup>7</sup> The Copenhagen Trial Unit, Centre for Clinical Intervention Research, Copenhagen, Denmark.

## Acknowledgements

We would like to thank all medical students and coordinators from the SICS Study Group for their devoted involvement with patient inclusions.

SICS Study Group information: The SICS Study Group members include the following: project leaders: Geert Koster, MD; Frederik Keus, MD, PhD; Iwan CC van der Horst, MD, PhD. Research coordinator: Willem Dieperink, PhD. Researchers who conducted patient inclusions: Roos Bleijendaal, MD; Yasmin F. Cawale, MD; Ramon P. Clement, MD; Devon Dijkhuizen, BSc; Ruben J Eck, MD; Bart Hiemstra, MD; Anja Haker, BSc; Casper D.H. Hilbink, MD; Thomas Kaufmann, MD; Martiene Klasen, MD; Manon Klaver, MD; Laura J. Schokking, BSc; Victor W. Sikkens, MD; Madelon Vos, MD; Justin Woerlee, MD and Renske Wiersema, BSc.

## Author contributions

BH, IvdH and EK drafted the manuscript and conducted the analyses. IvdH and EK were involved in the study conception. BH, RJE and GK developed the protocol and implemented the study. RW, YMH and TK contributed substantially to the data collection. JW and HS contributed to the design of the detailed statistical analysis plan. TWLS, AP, and AMdS critically reviewed the manuscript for important intellectual content. All authors critically reviewed the manuscript and agreed with the final version and findings.

## Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## Compliance with ethical standards

## Conflicts of interest

TWLS reports personal fees and other compensation from Edwards Lifesciences, personal fees from Masimo Inc., outside the submitted work. The other authors declare that they have no conflict of interest.

## Ethical approval

The study was approved by the local institutional review board (Medisch Ethische Toetsingscommissie, University Medical Center Groningen; METC M15.168207).

## OpenAccess

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the

original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 23 October 2018 Accepted: 9 January 2019

Published online: 31 January 2019

## References

- Cecconi M, De Backer D, Antonelli M et al (2014) Consensus on circulatory shock and hemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. *Intensive Care Med* 40:1795–1815. <https://doi.org/10.1007/s00134-014-3525-z>
- Vincent JL, De Backer D (2013) Circulatory shock. *N Engl J Med* 369:1726–1734. <https://doi.org/10.1056/NEJMr1208943>
- Narula J, Chandrashekar Y, Braunwald E (2018) Time to add a fifth pillar to bedside physical examination: inspection, palpation, percussion, auscultation, and insonation. *JAMA Cardiol*. <https://doi.org/10.1001/jamacardio.2018.0001>
- Verghese A, Charlton B, Kassirer JP, Ramsey M, Ioannidis JP (2015) Inadequacies of physical examination as a cause of medical errors and adverse events: a collection of vignettes. *Am J Med*. <https://doi.org/10.1016/j.amjmed.2015.06.004>
- Elder A, Japp A, Verghese A (2016) How valuable is physical examination of the cardiovascular system? *BMJ* 354:i3309. <https://doi.org/10.1136/bmj.i3309>
- Hiemstra B, Eck RJ, Keus F, van der Horst ICC (2017) Clinical examination for diagnosing circulatory shock. *Curr Opin Crit Care*. <https://doi.org/10.1097/mcc.0000000000000420>
- Perel A, Saugel B, Teboul JL et al (2016) The effects of advanced monitoring on hemodynamic management in critically ill patients: a pre and post questionnaire study. *J Clin Monit Comput* 30:511–518. <https://doi.org/10.1007/s10877-015-9811-7>
- Duan J, Cong LH, Wang H, Zhang Y, Wu XJ, Li G (2014) Clinical evaluation compared to the pulse indicator continuous cardiac output system in the hemodynamic assessment of critically ill patients. *Am J Emerg Med* 32:629–633. <https://doi.org/10.1016/j.ajem.2014.03.023>
- Nowak RM, Sen A, Garcia AJ, Wilkie H, Yang JJ, Nowak MR, Moyer ML (2012) The inability of emergency physicians to adequately clinically estimate the underlying hemodynamic profiles of acutely ill patients. *Am J Emerg Med* 30:954–960. <https://doi.org/10.1016/j.ajem.2011.05.021>
- Iregui MG, Prentice D, Sherman G, Schallom L, Sona C, Kollef MH (2003) Physicians' estimates of cardiac index and intravascular volume based on clinical assessment versus transesophageal Doppler measurements obtained by critical care nurses. *Am J Crit Care* 12:336–342
- Veale WN Jr, Morgan JH, Beatty JS, Sheppard SW, Dalton ML, Van de Water JM (2005) Hemodynamic and pulmonary fluid status in the trauma patient: are we slipping? *Am Surg* 71:621–626
- Connors AF Jr, Dawson NV, McCaffree R, Gray BA, Siciliano CJ (1987) Assessing hemodynamic status in critically ill patients: do physicians use clinical information optimally? *J Crit Care* 2:174–180
- Staudinger T, Locker GJ, Laczika K et al (1998) Diagnostic validity of pulmonary artery catheterization for residents at an intensive care unit. *J Trauma* 44:902–906
- Mimoz O, Rauss A, Rekik N, Brun-Buisson C, Lemaire F, Brochard L (1994) Pulmonary artery catheterization in critically ill patients: a prospective analysis of outcome changes associated with catheter-prompted changes in therapy. *Crit Care Med* 22:573–579
- Steingrub JS, Celoria G, Vickers-Lahti M, Teres D, Bria W (1991) Therapeutic impact of pulmonary artery catheterization in a medical/surgical ICU. *Chest* 99:1451–1455
- Hiemstra B, Eck RJ, Koster G et al (2017) Clinical examination, critical care ultrasonography and outcomes in the critically ill: cohort profile of the simple intensive care studies-I. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2017-017170>
- Hiemstra B, Nolte IM, Thio CH et al. (2018) Detailed statistical analysis plan of the simple intensive care studies-I (SICS-I). *Clinicaltrials.gov*: NCT02912624. <https://clinicaltrials.gov/ct2/show/NCT02912624>. Accessed 3 Apr 2018
- Schrager DL, Baraff L (1988) Defining normal capillary refill: variation with age, sex, and temperature. *Ann Emerg Med* 17:932
- Curley FJ, Smyrniotis NA (1990) Routine monitoring of critically ill patients. *J Intensive Care Med* 5:153–174. <https://doi.org/10.1177/088506669000500402>
- Ait-Oufella H, Lemoine S, Boelle PY et al (2011) Mottling score predicts survival in septic shock. *Intensive Care Med* 37:801–807. <https://doi.org/10.1007/s00134-011-2163-y>
- Coats AJ (1990) Doppler ultrasonic measurement of cardiac output: reproducibility and validation. *Eur Heart J* 11(Suppl I):49–61
- Rao V, Ivanov J, Weisel RD, Ikonomidis JS, Christakis GT, David TE (1996) Predictors of low cardiac output syndrome after coronary artery bypass. *J Thorac Cardiovasc Surg* 112:38–51
- National Heart L, Wheeler AP, Bernard GR et al (2006) Pulmonary-artery versus central venous catheter to guide treatment of acute lung injury. *N Engl J Med* 354:2213
- Godje O, Hoke K, Goetz AE et al (2002) Reliability of a new algorithm for continuous cardiac output determination by pulse-contour analysis during hemodynamic instability. *Crit Care Med* 30:52–58
- Teboul JL, Saugel B, Cecconi M et al (2016) Less invasive hemodynamic monitoring in critically ill patients. *Intensive Care Med* 42:1350–1359. <https://doi.org/10.1007/s00134-016-4375-7>
- Lang RM, Badano LP, Mor-Avi V et al (2015) Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 16:233–270. <https://doi.org/10.1093/ehjci/jev014>
- White IR, Royston P, Wood AM (2011) Multiple imputation using chained equations: issues and guidance for practice. *Stat Med* 30:377–399. <https://doi.org/10.1002/sim.4067>
- Bossuyt PM, Reitsma JB, Bruns DE et al (2015) STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ*. <https://doi.org/10.1136/bmj.h5527>
- Simel DL, Rennie D, Bossuyt PM (2008) The STARD statement for reporting diagnostic accuracy studies: application to the history and physical examination. *J Gen Intern Med*. <https://doi.org/10.1007/s11606-008-0583-3>
- Jakobsen JC, Wetterslev J, Winkel P, Lange T, Gluud C (2014) Thresholds for statistical and clinical significance in systematic reviews with meta-analytic methods. *BMC Med Res Methodol* 14:120. <https://doi.org/10.1186/1471-2288-14-120>
- Joly HR, Weil MH (1969) Temperature of the great toe as an indication of the severity of shock. *Circulation* 39:131–138
- Bailey JM, Levy JH, Kopel MA, Tobia V, Grabenkort WR (1990) Relationship between clinical evaluation of peripheral perfusion and global hemodynamics in adults after cardiac surgery. *Crit Care Med* 18:1353–1356
- Vincent JL, Moraine JJ, van der Linden P (1988) Toe temperature versus transcutaneous oxygen tension monitoring during acute circulatory failure. *Intensive Care Med* 14:64–68
- Schey BM, Williams DY, Bucknall T (2009) Skin temperature as a noninvasive marker of haemodynamic and perfusion status in adult cardiac surgical patients: an observational study. *Intensive Crit Care Nurs* 25:31–37. <https://doi.org/10.1016/j.iccn.2008.05.003>
- Sasse SA, Chen PA, Mahutte CK (1996) Relationship of changes in cardiac output to changes in heart rate in medical ICU patients. *Intensive Care Med* 22:409–414
- Grissom CK, Morris AH, Lanken PN et al (2009) Association of physical examination with pulmonary artery catheter parameters in acute lung injury. *Crit Care Med* 37:2720–2726
- Wetterslev M, Moller-Sorensen H, Johansen RR, Perner A (2016) Systematic review of cardiac output measurements by echocardiography vs. thermodilution: the techniques are not interchangeable. *Intensive Care Med* 42:1223–1233. <https://doi.org/10.1007/s00134-016-4258-y>