

# Diagnostic errors in the intensive care unit: a systematic review of autopsy studies

Bradford Winters,<sup>1</sup> Jason Custer,<sup>2</sup> Samuel M Galvagno Jr,<sup>1</sup> Elizabeth Colantuoni,<sup>3</sup> Shruti G Kapoor,<sup>1</sup> HeeWon Lee,<sup>4</sup> Victoria Goode,<sup>5</sup> Karen Robinson,<sup>6</sup> Atul Nakhasi,<sup>3</sup> Peter Pronovost,<sup>1</sup> David Newman-Toker<sup>7</sup>

► An additional appendix is published online only. To view this file please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2012-000803/content/early/recent>).

<sup>1</sup>Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, UK

<sup>2</sup>Department of Pediatrics, Division of Pediatric Intensive Care, University of Maryland, Baltimore, Maryland, UK

<sup>3</sup>Department of Biostatistics, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, UK

<sup>4</sup>Johns Hopkins University School of Medicine, Baltimore, Maryland, UK

<sup>5</sup>Welch Medical Library, Johns Hopkins University School of Medicine, Baltimore, Maryland, UK

<sup>6</sup>Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, UK

<sup>7</sup>Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, UK

## Correspondence to

Dr Bradford Winters, Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, 600 N. Wolfe St, Baltimore, MD 21287, UK; [bwinters@jhmi.edu](mailto:bwinters@jhmi.edu)

Accepted 24 June 2012

## ABSTRACT

**Context:** Misdiagnoses may be an underappreciated cause of preventable morbidity and mortality in the intensive care unit (ICU). Their prevalence, nature, and impact remain largely unknown.

**Objectives:** To determine whether potentially fatal ICU misdiagnoses would be more common than in the general inpatient population (~5%), and would involve more infections or vascular events.

**Data sources:** Systematic review of studies identified by electronic (MEDLINE, etc.) and manual searches (references in eligible articles) without language restriction (1966 through 2011).

**Study selection and data abstraction:** Observational studies examining autopsy-confirmed diagnostic errors in the adult ICU were included. Studies analysing misdiagnosis of one specific disease were excluded. Study results (autopsy rate, misdiagnosis prevalence, Goldman error class, diseases misdiagnosed) were abstracted and descriptive statistics calculated. We modelled the prevalence of Class I (potentially lethal) misdiagnoses as a non-linear function of the autopsy rate.

**Results:** Of 276 screened abstracts, 31 studies describing 5863 autopsies (median rate 43%) were analysed. The prevalence of misdiagnoses ranged from 5.5%–100% with 28% of autopsies reporting at least one misdiagnosis and 8% identifying a Class I diagnostic error. The projected prevalence of Class I misdiagnoses for a hypothetical autopsy rate of 100% was 6.3% (95% CI 4.0% to 7.5%). Vascular events and infections were the leading lethal misdiagnoses (41% each). The most common individual Class I misdiagnoses were PE, MI, pneumonia, and aspergillosis.

**Conclusions:** Our data suggest that as many as 40 500 adult patients in an ICU in USA may die with an ICU misdiagnosis annually. Despite this, diagnostic errors receive relatively little attention and research funding. Future studies should seek to prospectively measure the prevalence and impact of diagnostic errors and potential strategies to reduce them.

## INTRODUCTION

Diagnostic errors are an underappreciated cause of preventable mortality and morbidity in hospitalised patients, accounting for an estimated 40 000–80 000 annual deaths in USA.<sup>1</sup> Patients in an intensive care unit (ICU) may be especially prone to suffer harm from diagnostic errors. They have limited reserve, often require fast diagnosis and treatment, are cared for by multiple clinicians and undergo frequent laboratory and imaging evaluations.<sup>2</sup> Harm may result from delay or failure to treat the correct underlying disease, complications of unnecessary diagnostic testing or treating a condition that is not actually present.<sup>1</sup>

Diagnostic errors are missed, wrong or delayed diagnoses that are detected by some subsequent definitive test or finding.<sup>3</sup> Autopsy-detected misdiagnoses are typically classified by the Goldman criteria and based on their clinical relevance and the potential that timely therapy would have changed the outcome. The Goldman criteria for autopsy confirmed discrepancies stratify misdiagnosis errors as major (Class I and II misdiagnosis) and minor (Class III and IV misdiagnosis). A Class I error is a major missed diagnosis with a potential adverse impact on survival (likely was the cause of death or contributed to the patient's death) and had that diagnosis been known would have changed management and outcome. A Class II error is a major missed diagnosis that did not have an impact on survival (was not the cause of death nor directly contributed to the patient's death). A Class III error is a minor missed diagnosis related to a terminal disease but not related to the cause of death, and Class IV errors are other missed minor discrepancies.<sup>4</sup>

## Systematic review

A 2003 systematic review of autopsy-confirmed misdiagnoses found that 9% were Class I errors and 15% were Class II errors.<sup>5</sup> After adjusting for declining autopsy rates over time, the authors estimated that 10% of all hospital deaths had a major diagnostic error and 5% of these were Class I errors. They also estimated that the Class I error prevalence in the ICU population was twice the overall prevalence in US hospitals.

Nevertheless, the prevalence of diagnostic errors, the conditions ultimately diagnosed and the degree of harm from these errors are largely unknown. The methods to measure diagnostic errors are underdeveloped and most studies measuring patient safety ignore diagnostic errors.<sup>1</sup> In a recent study for example, diagnostic error was not a variable in examining temporal changes in patient harm over time.<sup>6</sup> As such, the problem of diagnostic errors remains largely invisible. The purpose of this study was to systematically estimate the prevalence and distribution of autopsy-confirmed diagnostic errors in the ICU population. We hypothesised that diagnostic errors would be more common in the ICU population than previously reported for the general hospital population. Additionally, based on a preliminary assessment of the literature, we hypothesised that most errors identified related to infections or vascular events.

## METHODS

### Search strategy


We conducted a systematic review of the published literature from 1966 through 2011 for observational studies that examined the prevalence of potentially lethal diagnostic errors in the ICU, as diagnosed by the reference standard of autopsy. Databases searched included PubMed, Embase, Cumulative Index to Nursing and Allied Health Literature, and the Cochrane Database of Systematic Reviews using the Johns Hopkins Welch Medical Library online electronic resources. There were no language limitations. The full search strategies (see online appendix) were designed by librarians from the Johns Hopkins Welch Medical Library (KR, VG) together with clinical investigators (BW, DNT). All of the clinical investigators that served as reviewers have expertise in critical care or evaluation of diagnostic errors. Two reviewers (JC and SGK) independently screened each abstract and all studies considered eligible by either reviewer were passed on to two reviewers (JC and BW) for a full article review to determine inclusion in the final analysis. Disagreements between these reviewers were resolved by group consensus. Bibliographies from retrieved articles were screened for other relevant studies; none were identified. We did not attempt to contact authors or identify meeting abstracts, unpublished studies, or other grey literature.

### Study selection

We included studies if they provided autopsy-confirmed misdiagnosis data on adult patients who died while in an intensive care setting. Studies without original patient data were excluded. We did not set standards for autopsy performance or for conduct and reporting of individual studies (eg, prospective vs retrospective). Two types of studies were defined: (1) general misdiagnosis studies (reporting on all misdiagnoses or all misdiagnoses of a minimum severity),<sup>7–37</sup> and (2) disease-specific studies (reporting only on misdiagnoses of particular conditions, such as pulmonary embolus).<sup>38–46</sup> We excluded disease-specific studies from our analysis. A flowchart of our study selection is in [figure 1](#).

### Analysis

Four authors (JC, BW, AN, SMG) reviewed the general misdiagnosis studies and abstracted the data for pooled analysis. These data included number of deaths, autopsies and errors, error class, underlying diagnoses, country and ICU type. We did not assign a quality rating. Errors were defined by the Goldman Classification scheme. Class I are missed major diagnoses with potential adverse impact on survival and that would have changed management. Class II is a missed major diagnosis that would not have impacted survival and that



**Figure 1** This figure shows the study selection flowchart.

would not have changed therapy. Class III and Class IV are minor errors related to the terminal disease but not death and completely unrelated to either, respectively. The majority of studies reported Class I<sup>7–16 18–32 34–37</sup> and II<sup>7 9 10 12 13 15 16 18–32 34–36</sup> error data, and less reported Class III<sup>9 10 12 13 15 16 18–20 22 26 30 31</sup> and IV<sup>9 12 13 16 18 20 23</sup> error data.

We defined four broad disease categories to assess the distribution of types of misdiagnoses. These categories included: (1) vascular (myocardial infarction, pulmonary embolus, stroke, any organ infarct, any vessel rupture or dissection, and any haemorrhage), (2) infectious (bacterial, aspergillosis, other fungal, viral, and tuberculosis), (3) mechanical pathophysiological states (cardiac tamponade, bowel obstruction, perforated viscus, congestive heart failure, and cardiac valve failure), and (4) neoplastic/other. Some studies treated pneumonia and aspergillosis separately from other infections.<sup>10 14 19 20 25 34</sup> We aggregated these results during the analytic phase.

Data were tabulated using Microsoft Excel 2008 for Mac (Microsoft Corp., Redmond, Washington, USA). Descriptive statistics were calculated using Stata V.11 (Stata Corp., College Station). A *p* value <0.05 was considered statistically significant. A logistic regression model was performed to create a hypothetical prediction of the prevalence of misdiagnoses at higher autopsy rates to predict what the actual error rate might be if every patient who died in an ICU always had an autopsy. A random intercept logistic regression model was fit to the class I misdiagnosis rate (number of class I errors vs number of autopsies performed) as a non-linear function of the per cent of autopsies performed (natural spline with two degrees of freedom). A likelihood ratio test was performed with R V.2.11.1 (R Foundation for Statistical Computing, Vienna, Austria) to determine if the non-linear function was justified over a linear fit. The random intercept was included to account for natural variation across studies as is standard in meta-analysis. We controlled for size of study (number of autopsies) and the study time period in this model.

This report was prepared according to Meta-analysis of Observational Studies in epidemiology (MOOSE) guidelines for systematic reviews of observational studies.<sup>47</sup>

## RESULTS

Of 276 abstracts screened, 78 were advanced to full article review ( $\kappa$  for concordance was 0.97; SE =0.11). We identified 41 studies and subsequently excluded one by group consensus because the unit studied could not be clearly identified as an ICU. Of the remaining 40 studies, 30 reported general misdiagnosis (28 retrospective) representing a total of 5863 autopsies and were included in our

analysis, and nine were disease-specific misdiagnosis and excluded from our analysis (tables 1 and 2). The study periods ranged from 11 months to 300 months and included a wide range of ICU types. Autopsy rates ranged from 6% to 100% (weighted mean 41%, median 43%).

The prevalence of autopsy-detected misdiagnoses ranged from 5.5% to 100%. In aggregate, 28% (n=1632/5863) of autopsies reported at least one misdiagnosis (figure 2). Of the 31 studies, 27 reported class I error data, 25 class II, 13 class III, and 7 class IV error data. Class I diagnostic errors were identified in 8% (n=343/4514); Class II errors in 15% (n=637/4403); Class III errors in 15% (n=264/1711); and Class IV errors in 21% (n=175/816) of autopsies reporting (figure 2). Class I misdiagnoses were less prevalent at higher autopsy rates (figure 3) although total misdiagnoses were not (data not shown). The predicted prevalence of potentially lethal misdiagnoses, when extrapolated to a hypothetical 'perfect' 100% autopsy rate was 6.3% (95% CI=4.0% to 7.5%). Our model estimates the Class I error prevalence as a function of autopsy rate; the model result (6.3%) differs from the measured prevalence (8%), which is simply the average misdiagnosis prevalence from all autopsies reporting Class I errors. A likelihood ratio test was used to determine if the non-linear function was justified over a linear fit (*p*=0.021). In addition, a likelihood ratio test was used to determine if the autopsy rate (based on natural spline fit) provided any information for the class I misdiagnosis rate (*p*=0.002). After accounting for the autopsy rate, the estimated SD of the log odds of a class I misdiagnosis across studies was 0.58 (SE: 0.11), translating to study to study variation accounting for roughly 10% of the total variation.

Final diagnoses were provided in 25 of 31 studies for Class I misdiagnoses and in 16 of 31 studies for Class II misdiagnoses (figure 4 and box 1). Among all Class I errors, vascular events (41%) and infections (41%) were the most common diagnosis, followed by mechanical pathophysiological states (10%), and neoplasms and other conditions (9%). For all Class II errors, vascular events (47%) were the most common diagnosis, followed by infections (29%), neoplasms and other conditions (21%), and mechanical obstructive pathophysiological conditions (4%). Within these broad categories, pulmonary embolus, myocardial infarction, pneumonia, and aspergillosis were the most common diagnoses in the class I errors. Cumulatively, they accounted for about one-third of all potentially lethal diagnostic errors (box 1).

## DISCUSSION

In this study, 28% of autopsied ICU patients had at least one misdiagnosis. Of the autopsies reporting misdiagnosis error classifications, 8% were Class I (major and

## Systematic review

**Table 1** Included studies of intensive care unit (ICU) misdiagnoses in general

Author	Year	Country	Type of study	Length of study period (months)	ICU type	Total # of deaths	No of autopsies used for analysis
Berlot <sup>7</sup>	1999	Italy	Retrospective	36	Mixed	346	159
Blosser <sup>8</sup>	1998	USA	Retrospective	12	Med/ coronary	132	41
Combes <sup>9</sup>	2004	France	Prospective	36	Med/surg	315	167
Dimopolous <sup>10</sup>	2004	Belgium	Retrospective	12	Med/surg	489	222
Duke <sup>11</sup>	1999	Australia	Retrospective	24	Not given	238	Not given
Fernandez-Segoviano <sup>12</sup>	1988	Spain	Prospective	30	Mixed	Not given	100
Fish <sup>13</sup>	2000	USA	Retrospective	72	Burn	94	88
Gerain <sup>14</sup>	1990	Belgium	Retrospective	11	Oncology	48	34
Gut <sup>15</sup>	1999	Brazil	Retrospective	36	Med/surg	152	30
Kallinen <sup>16</sup>	2008	Finland	Retrospective	72	Burn	74	71
Koch <sup>17</sup>	2008	Germany	Retrospective	120	Medical	1205	1205
Magret <sup>18</sup>	2006	Spain	Retrospective	46	Mixed	525	80
Maris <sup>19</sup>	2007	Belgium	Retrospective	24	Med/surg	786	289
Mort <sup>20</sup>	1999	USA	Retrospective	72	Surg	560	149
Nadrous <sup>21</sup>	2003	USA	Retrospective	24	Multiple	1597	455
Ong <sup>22</sup>	2002	USA	Retrospective	24	Trauma/burn	158	153
Palazon-Sanchez <sup>23</sup>	1999	Spain	Retrospective	12	Med/surg	67	24
Papadakis <sup>24</sup>	1991	USA	Retrospective	24	Medical	401	172
Pastores <sup>25</sup>	2007	USA	Retrospective	69	Med/surg	658	86
Perkins <sup>26</sup>	2003	Great Britain	Retrospective	42	Med/surg	636	38
Podbregar <sup>27</sup>	2001	Slovenia	Retrospective	24	Medical	270	126
Podbregar <sup>28</sup>	2011	Slovenia	Retrospective	20	Medical	373	170
Roosen <sup>29</sup>	2000	Belgium	Retrospective	24	Medical	108	100
Saad <sup>30</sup>	2007	Brazil	Retrospective	24	Coronary	Unclear	161
Sharma <sup>31</sup>	2005	India	Retrospective	36	Trauma/burn	249	163
Silfvast <sup>32</sup>	2003	Finland	Retrospective	48	Mixed	388	346
Simon <sup>33</sup>	2001	Hungary	Retrospective	12	Medical	163	110
Tai <sup>34</sup>	2001	USA	Retrospective	24	Medical	401	91
Tejerina	2010	Spain	Retrospective	300	Mixed	2857	866
Twigg <sup>36</sup>	2001	Great Britain	Retrospective	36	Med/surg	252	97
Yalamarthy <sup>37</sup>	1998	Great Britain	Retrospective	24	Not given	233	70

ICU, Intensive Care Unit.

potentially lethal), and 15% were Class II (major but not lethal) despite presumably aggressive diagnostic assessment in the critical care environment. Given the 540 000 deaths annually during or immediately following an ICU stay,<sup>48</sup> and extrapolating to a 100% autopsy rate, our findings suggest that 34 000 (95% CI =22 600 to 40 500) ICU patients in the US may die as a result of a Class I error annually, assuming that the error was the cause of death. This predicted number is comparable to the estimated deaths from central line-associated bloodstream infections in the ICU<sup>49</sup> and from breast cancer<sup>50</sup> in the US, and does not consider misdiagnosis-related morbidity among intensive care patients. These results also suggest that despite the dropping autopsy rate over time, autopsy remains a crucial asset especially in ICU patients.

We proved our hypothesis that infections and vascular events would culminate a large majority of the diagnostic errors. Vascular events and infections were the leading

potentially lethal misdiagnoses accounting for almost 82% of potentially lethal Class I errors and almost 76% of the serious but not lethal Class II errors. Neoplastic disease was the third most common overall misdiagnosis category but was rarely deemed lethal. Mechanical pathophysiological states were uncommon. Importantly, pulmonary embolus, myocardial infarction, pneumonia, and aspergillosis accounted for over one-third of all Class I misdiagnoses. These four particular diagnostic errors may be useful as outcome measure for designing early interventions to reduce diagnostic errors as a whole.

Our results add new information to the field of diagnostic error research. Shojania *et al*<sup>5</sup> conducted a systematic review of autopsy-detected diagnostic error prevalence rates over time, but they did not describe specific misdiagnoses in the ICU or elsewhere. We included the seven ICU studies from their review in our analysis to offer refined estimates of the prevalence of

**Table 2** Condition/disease specific misdiagnosis studies not analysed

Author	Year	Country	Type of study	Study period (mos)	ICU type	Total no of deaths	No of autopsies analysed	No of missed diagnoses	Disease or diagnosis
deHemptinne <sup>38</sup>	2009	Belgium	Retrospective	36	Med/surg	169	64	15	ARDS only
Du <sup>39</sup>	1996	China	Retrospective	492	Surg	Not given	3447	80	Fungal only
Hasani <sup>40</sup>	2005	Iran	Prospective	12	Mixed	Not given	43	2	Meningitis only
Janson <sup>41</sup>	1976	Germany	Retrospective	24	Surg	100	100	48	CXR misDx only
Petersen <sup>43</sup>	1999	Denmark	Prospective	12	Multiple	659	141	70	Pneumonia only
Pinheiro <sup>44</sup>	2007	Brazil	Retrospective	96	Not given	592	22	7	ARDS only
Al-Saidi <sup>42</sup>	2002	Canada	Retrospective	67	Med/surg	59	28	10	Bone marrow transplant only
Tejerina <sup>45</sup>	2009	Spain	Retrospective	150	Not given	1633	253	111	VAP only
Widimsky <sup>46</sup>	1985	Czech-Slovakia	Retrospective	120	Coronary	Not given	31	12	PE only

ARDS, Acute Respiratory Distress syndrome; CXR, chest X-ray; ICU, Intensive Care Unit; misDx, misdiagnosis; mos, months; PE, pulmonary embolus; VAP, ventilator-associated pneumonia.


misdiagnoses in the ICU by error class and autopsy rate. Shojania *et al* projected that, despite a reduction of 33% in the rate of class I errors over 4 decades, a US hospital would likely continue to observe a class I error at a rate of 4.1%–6.7% across all patients. While a few of the studies in Shojania's review included ICU patients in addition to general ward and emergency department patients, our results suggest that the rate of class I errors are 16%–48% higher in the ICU patient population than the general hospital patient population as a whole. Methodologically, we found that autopsy rates of 30%–40% or higher are likely to produce fairly accurate estimates of the overall prevalence of potentially lethal misdiagnoses in the ICU population, despite potential biases inherent in autopsy case selection.<sup>51</sup>

Observed diagnostic error frequencies were highest in studies with the lowest autopsy rates. Clinical selection of the most complex cases to undergo autopsy is believed to contribute to this effect.<sup>51</sup> To counter this effect, we have modelled the measured error frequency as a function of autopsy rate, controlled for study size and time and extrapolated out to conditions of 100% autopsy. Nevertheless, it is possible that other factors may contribute to the difference seen in studies with low autopsy rates such as a more detailed and rigorous approach to autopsy and error finding. Additionally, the level of diagnostic certainty as to the cause of death is likely to influence the autopsy rate. Podbregar *et al*<sup>27</sup> attempted to measure this influence finding that diagnostic certainty significantly influenced the decision to pursue an autopsy though there were no differences in the class I error rate. As a consequence, our extrapolated estimate for the total error frequency likely represents a conservative, minimum estimate of the true frequency as aggregated from the present data.

Although we could not measure morbidity associated with misdiagnosis in survivors, the consequences and costs are likely substantial. ICU-based studies have reported complications of misdiagnosis, such as re-operation,<sup>52</sup> and increased length of stay.<sup>53</sup> The most frequently reported non-lethal long-term consequences of misdiagnosis are neurological. Preventable or treatable neurological complications of critical illness such as stroke may go unrecognised with devastating consequences.<sup>54–56</sup> Robust estimates of the morbid consequences of diagnostic errors will likely require prospective studies.

Risk predictors for misdiagnosis in the ICU at the patient, organisational or provider level are incompletely defined and there is sparse data on root causes of misdiagnosis in the ICU. Off-hours occurrence appears to be a risk factor for at least some types of ICU misdiagnosis.<sup>53</sup> Patient demographics and institutional factors have been cited as possible influences.<sup>5 57 58</sup> There is



## Systematic review



**Figure 2** This figure shows the proportion of all autopsies that identified a misdiagnosis, and the proportion of autopsies that identified a misdiagnosis stratified by the Goldman criterion (Class I, II, III, and IV). The grey bars represent identified misdiagnosis and the black bars represent all autopsies reported for total and class. Of the 30 studies included in this analysis, 26 reported class I error data, 24 class II, 13 class III, and 7 class IV error data. Thus, the sample sizes by Goldman classification varied. Stratified by class, 8% of autopsies detected a Class I error, 15% detected a Class II error, 15% detected a Class III error, and 21% detected a Class IV error.

evidence that the patient's clinical presentation such as greater co-morbidity or illness complexity increases the risk of misdiagnosis.<sup>52 53 56 59</sup> The diagnostic process is complex<sup>60</sup> and errors may occur at any stage during the process, often at multiple stages.<sup>61</sup> Some misdiagnoses may be directly linked to limited sensitivity and specificity of individual tests in the critical care setting,<sup>62</sup> but remediable causes are most often associated with failures

**Figure 4** This figure shows the most common misdiagnoses by broad disease categories for Class I and II errors. These categories are vascular (myocardial infarction, pulmonary embolus, stroke, any organ infarct, any vessel rupture or dissection, and any haemorrhage), infectious (bacterial, aspergillosis, other fungal, viral and tuberculosis), mechanical pathophysiological states (cardiac tamponade, bowel obstruction, perforated viscus, congestive heart failure and cardiac valve failure), and neoplasm and other (includes acute respiratory distress syndrome (ARDS)). The number of Class I (black bar) and Class II (grey bar) errors are compared for each diagnostic category. The data for Class I errors represents 24 of the 30 studies in the analysis and data for Class II errors represents 16 of 30 studies.



**Figure 3** This figure illustrates the prevalence of Class I misdiagnoses (as a per cent of autopsies) and the autopsy rate. Each point on the figure represents data from one study. The plotting symbols are proportional to the number of autopsies performed in each study (with larger plotting symbols indicating larger number of autopsies). The solid line represents the logistic regression models estimate for reported values. The dotted lines represent the 95% CIs. The curve was generated using a natural spline with two degrees of freedom.

in information processing. Cognitive errors are also a major contributor to misdiagnoses and a recent review by Graber *et al*<sup>63</sup> identified a large number of possible approaches (some tested and some not) that may help reduce cognitive errors and hence, misdiagnoses.

**Box 1** Breakdown of misdiagnoses by final diagnosis reported

Vascular events
Haemorrhage
Vascular rupture
Aortic aneurysm/dissection (11, 1)
Other (2, 1)
Abdominal bleed (21, 41)
Other (5, 12)
Thrombosis
Pulmonary embolus (37, 64)
Other (1, 0)
Ischemia
Myocardial infarction (31, 33)
Bowel infarction (13, 10)
Stroke (1, 0)
Infections
Mechanical obstructive pathophysiology
Cardiac tamponade (10, 0)
Perforated viscus not designated peritonitis (9, 4)
Bowel obstruction (1, 4)
Tension pneumothorax (1, 0)
Congestive heart failure (2, 2)
Cardiac valvular lesion (not endocarditis) (5, 2)
Other
Malignancy (6, 47)
Pulmonary oedema (not acute respiratory distress syndrome) (2, 2)
Acute respiratory distress syndrome (3, 1)
Pancreatitis (4, 1)
Cirrhosis (0, 1)
Airway obstruction (2, 0)
Other (9, 22)

Values in parentheses are the actual total number of class I and class II identified in studies reporting this data.

A major impediment to better understanding of diagnostic errors is the declining autopsy rate.<sup>5</sup> The reasons for this are multi-factorial and likely include medical culture and societal attitude shifts. One alternative that might improve the rate of autopsy is the ‘virtual autopsy’ using sophisticated radiological techniques as opposed to dissection. Wichmann *et al.*<sup>64</sup> recently reported a comparison of ‘virtual autopsy’ to traditional autopsy in ICU patients suggesting that it compares favourably and had a higher acceptance rate in terms of per cent of deaths referred to post-mortem analysis. With further refinement this approach may offer a way to continue to obtain the post-mortem information required.

Our systematic review has significant policy implications. Diagnostic errors have not been the focus of patient safety campaigns or systematic studies.<sup>1 65</sup> A 2003 report of 93 Agency for Health Research and Quality (AHRQ)-funded safety projects found only one focused on misdiagnosis.<sup>65</sup> A recent Special Emphasis Notice has sought proposals for these errors in ambulatory and

emergency department settings,<sup>66</sup> but it has not been linked to a funded request for applications.

Our review has limitations. We did not search the grey literature nor did we follow-up with authors, pursue secondary references or consult with experts and publication bias could, in theory, have inflated our measured prevalence. Virtually all studies were retrospective in nature and there is potential for selection bias in those autopsied though Podbregar *et al.*<sup>27</sup> found that while diagnostic certainty did influence the rate of autopsy, it did not influence the rate of class I errors. Unfortunately, the reliability of determining whether an error occurred and its severity could not be determined to any meaningful extent in the reviewed studies. Thus our estimate of the class I (lethal) diagnostic error frequency may either be an over- or under-estimate of the true frequency. Debate continues on the extent to which this is a potential problem.<sup>11</sup> There was some heterogeneity in the study populations included in our review and the measured misdiagnosis prevalence varied even at high autopsy rates, so misdiagnosis prevalence might not be evenly distributed. We did not assign a quality rating to the studies analysed based on autopsy confirmed diagnosis being a ‘gold standard’. Autopsy, as a ‘gold standard’, is potentially flawed as the decision to perform an autopsy or the results of the autopsy may be influenced by the degree of involvement of clinicians in the judgement of diagnostic errors, the degree to which pathologists knew the clinical history and working diagnoses at the time that they performed autopsy and the format/context in which requests for autopsy were transmitted. The original study authors may have misclassified the types of diagnostic errors, resulting in inaccurate estimates of lethality or preventability and disease frequency estimates might be distorted. Nevertheless, given the consistency with which studies reported missed infections and vascular events as a common source of lethal diagnostic errors, these are likely important.

In conclusion, this systematic review and analysis suggests that between 22 600–40 500 ICU patients die each year in the US with and potentially from a diagnostic error and many more suffer a clinically relevant diagnostic error. These data suggest that diagnostic errors are a significant cause of preventable harm in hospitalised patients. To this point, diagnostic errors have received relatively little attention and research funding, leaving the methods to measure them immature; this must change. Future studies should seek to prospectively measure the prevalence and direct impact of diagnostic errors, including those that are non-lethal, and potential strategies to reduce them.

**Acknowledgements** The authors wish to thank Christine G. Holzmueller, BLA for her assistance in editing the manuscript.

## Systematic review

**Contributors** Conception and design: BW, KR, DNT, PP. Acquisition of data: BW, VG, SMG, KR, JC, SGK, AN. Analysis and interpretation of data: BW, JC, HWL, EC, SMG, AN, DNT, PP. Drafting of the manuscript: BW, JC, DNT, PP. Critical revision of manuscript: BW, JC, HWL, SMG, AN, DNT, PP. Statistical analysis: BW, EC, SMG, AN. Administrative, technical, or material support: VG, KR, PP. Supervision: BW, DNT, PP.

**Funding** SMG was supported by an NIH training grant awarded to the Johns Hopkins University School of Medicine; DN-T was supported in part by a grant from the Agency for Healthcare Research and Quality (HS017755-01).

**Competing interests** BWinters reports fees for expert testimony at several defense and plaintiff law firms and speaking honoraria from 3M corporation and various health systems to speak on patient safety and Rapid Response Systems. PP reports receiving grant or contract support from the Agency for Healthcare Research and Quality, the National Institutes of Health, the Robert Wood Johnson Foundation and The Commonwealth Fund for research related to measuring and improving patient safety; honoraria from various hospitals and healthcare systems and the Leigh Bureau to speak on quality and safety; consultancy with the Association for Professionals in Infection Control and Epidemiology, Inc.; and book royalties for authoring "Safe Patients, Smart Hospitals: How One Doctor's Checklist Can Help Us Change Health Care from the Inside Out." The following authors report no conflicts of interest: JC, EC, SGK, HWL, VG, KR, AN; SMG and DN-T have no further conflicts beyond what is reported under funding.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- Newman-Toker DE, Pronovost PJ. Diagnostic errors—the next frontier for patient safety. *JAMA* 2009;301:1060–2.
- Newman-Toker DE, Pham JC, Winters BD, *et al*. Diagnostic errors in critical care settings—managing information overload. *ICU Management* 2009;9:6–11.
- Graber ML, Franklin N, Ruthanna G. Diagnostic error in internal medicine. *Arch Intern Med* 2005;165:1493–9.
- Goldman L, Sayson R, Robbins S, *et al*. The value of autopsy in three medical eras. *N Engl J Med* 1983;308:1000–5.
- Shojania KG, Burton EC, McDonald KM, *et al*. Changes in rates of autopsy-detected diagnostic errors over time: a systematic review. *JAMA* 2003;289:2849–56.
- Landrigan CP, Parry GJ, Bones CB, *et al*. Temporal trends in rates of patient harm resulting from medical care. *N Engl J Med* 2010;363:2124–34.
- Berlot G, Dezzoni R, Viviani M, *et al*. Does the length of stay in the intensive care unit influence the diagnostic accuracy? A clinical-pathological study. *Eur J Emerg Med* 1999;6:227–31.
- Blosser SA, Zimmerman HE, Stauffer JL. Do autopsies of critically ill patients reveal important findings that were clinically undetected? *Crit Care Med* 1998;26:1332–6.
- Combes A, Mokhtari M, Couvelard A, *et al*. Clinical and autopsy diagnoses in the intensive care unit: a prospective study. *Arch Intern Med* 2004;164:389–92.
- Dimopoulos G, Piagnerelli M, Berré J, *et al*. Post mortem examination in the intensive care unit: still useful? *Intensive Care Med* 2004;30:2080–5.
- Duke GJ, Morley PT, Cooper DJ, *et al*. Management of severe trauma in intensive care units and surgical wards. *Med J Aust* 1999;170:416–19.
- Fernandez-Segoviano P, Lázaro A, Esteban A, *et al*. Autopsy as quality assurance in the intensive care unit. *Crit Care Med* 1988;16:683–5.
- Fish J, Hartshorne N, Reay D, *et al*. The role of autopsy on patients with burns. *J Burn Care Rehabil* 2000;21:339–44.
- Gerain J, Sculier JP, Malengreaux A, *et al*. Causes of deaths in an oncologic intensive care unit: a clinical and pathological study of 34 autopsies. *Eur J Cancer* 1990;26:377–81.
- Gut AL, Ferreira AL, Montenegro MR. Autopsy: quality assurance in the ICU. *Intensive Care Med* 1999;25:360–3.
- Kallinen O, Partanen TA, Maisniemi K, *et al*. Comparison of premortem clinical diagnosis and autopsy findings in patients with burns. *Burns* 2008;34:595–602.
- Koch S, Bredahl J, Wirtz SP, *et al*. How often is bronchopneumonia overlooked as the cause of death in intensive care unit patients? (In German) *Anaesthetist* 2008;57:37–42.
- Magret Iglesias M, Vidaur Tello L, Fernández Olsina S, *et al*. Discrepancies between clinical and pathological diagnosis in a polyvalent intensive care service (In Spanish). *Med Intensiva* 2006;30:95–100.
- Maris C, Martin B, Creteur J, *et al*. Comparison of clinical and post-mortem findings in intensive care unit patients. *Virchows Arch* 2007;450:329–33.
- Mort TC, Yeston NS. The relationship of pre mortem diagnoses and post mortem findings in a surgical intensive care unit. *Crit Care Med* 1999;27:299–303.
- Nadrous HF, Afessa B, Pfeifer EA, *et al*. The role of autopsy in the intensive care unit. *Mayo Clin Proc* 2003;78:947–50.
- Ong AW, Cohn SM, Cohn KA, *et al*. Unexpected findings in trauma patients dying in the intensive care unit: results of 153 consecutive autopsies. *J Am Coll Surg* 2002;194:401–6.
- Palazon-Sanchez C, Palazon-Sanchez E, Pardo Talavera JC, *et al*. Findings on necropsy studies in critically ill patients. *Med Intensiva* 1999;23:6–9.
- Papadakis MA, Mangione CM, Lee KK, *et al*. Treatable abdominal pathologic conditions and unsuspected malignant neoplasms at autopsy in veterans who received mechanical ventilation. *JAMA* 1991;265:885–7.
- Pastores SM, Dulu A, Voigt L, *et al*. Premortem clinical diagnoses and postmortem autopsy findings: discrepancies in critically ill cancer patients. *Crit Care* 2007;11:R48–54.
- Perkins GD, McAuley DF, Davies S, *et al*. Discrepancies between clinical and postmortem diagnoses in critically ill patients: an observational study. *Crit Care* 2003;7:R129–32.
- Podbregar M, Voga G, Krivec B, *et al*. Should we confirm our clinical diagnostic certainty by autopsies? *Intensive Care Med* 2001;27:1750–5.
- Podbregar M, Kralj E, Čičak R, *et al*. A triad algorithm for analysing individual ante- and post-mortem findings to improve the quality of intensive care. *Anaesth Intensive Care* 2011;39:1086–92.
- Roosen J, Frans E, Wilmer A, *et al*. Comparison of premortem clinical diagnoses in critically ill patients and subsequent autopsy findings. *Mayo Clin Proc* 2000;75:562–7.
- Saad R, Yamada AT, Pereira da Rosa FH, *et al*. Comparison between clinical and autopsy diagnoses in a cardiology hospital. *Heart* 2007;93:1414–19.
- Sharma BR, Gupta M, Harish D, *et al*. Missed diagnoses in trauma patients vis-à-vis significance of autopsy. *Injury* 2005;36:976–83.
- Silfvast T, Takkunen O, Kolho E, *et al*. Characteristics of discrepancies between clinical and autopsy diagnoses in the intensive care unit: a 5-year review. *Intensive Care Med* 2003;29:321–4.
- Simon J, Rudas L, Iványi B. The autopsy in verifying diagnostic accuracy at the intensive care unit (In Hungarian). *Orv Hetil* 2001;142:2373–6.
- Tai DY, El-Bilbeisi H, Tewari S, *et al*. A study of consecutive autopsies in a medical ICU: a comparison of clinical cause of death and autopsy diagnosis. *Chest* 2001;119:530–6.
- Tejerina E, Esteban A, Fernández-Segoviano P, *et al*. Clinical diagnoses and autopsy findings; discrepancies in critically ill patients. *Crit Care Med* 2012;40:842–6.
- Twigg SJ, McCrerrick A, Sanderson PM. A comparison of post mortem findings with post hoc estimated clinical diagnoses of patients who die in a United Kingdom intensive care unit. *Intensive Care Med* 2001;27:706–10.
- Yalamarthi S, Ridley S, Barker T. Agreement between ante-mortem diagnoses, death certificates and post-mortem causes of death in critically ill patients. *Clin Intensive Care* 1998;9:100–4.
- deHemptinne Q, Rimmelink M, Brimiouille S, *et al*. ARDS: a clinicopathological confrontation. *Chest* 2009;135:944–9.
- Du B, Zhang H, Chen D. Invasive fungal infection in 3447 autopsy cases. *Natl Med J China* 1996;76:352–4.
- Hasani M, Sahraian MA, Motamedi M, *et al*. Postmortem cerebrospinal fluid analysis in a general intensive care unit. *Indian J Crit Care* 2005;9:176–8.
- Janson R, Thelen M, Rommelsheim K, *et al*. A review of the findings on chest examinations and at autopsy in surgical patients under intensive care. *Fortschr Röntgenstr* 1976;124:427–33.
- Al-Saidi F, Diaz-Granados N, Messner H, *et al*. Relationship between premortem and postmortem diagnosis in critically ill bone marrow transplantation patients. *Crit Care Med* 2002;30:570–3.
- Petersen IS, Aru A, Skødt V, *et al*. Evaluation of pneumonia diagnosis in intensive care patients. *Scand J Infect Dis* 1999;31:299–303.
- Pinheiro BV, Mruaoka FS, Assis RV, *et al*. Accuracy of clinical diagnosis of acute respiratory distress syndrome in comparison with autopsy findings. *J Bras Pneumol* 2007;33:423–8.

45. Tejerina E, Esteban A, Fernández-Segoviano P, *et al.* Accuracy of clinical definitions of ventilator associated pneumonia: comparison with autopsy findings. *J Crit Care* 2009;25:62–8.
46. Widimský J, Staněk V. Clinical diagnosis of haemodynamically significant pulmonary embolism in a coronary care unit. *Cor Vasa* 1985;27:337–45.
47. Stroup DF, Berlin JA, Morton SC, *et al.* Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008–12.
48. Angus DC, Black N. Improving care of the critically ill: institutional and health-care system approaches. *Lancet* 2004;363:1314–20.
49. Pronovost P, Needham D, Berenholtz S, *et al.* An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med* 2006;355:2725–32.
50. *Cancer Facts and Figures, 2010*. Atlanta: American Cancer Society, 2010. <http://www.cancer.org/acs/groups/content/@epidemiologysurveillance/documents/document/acspc-026238.pdf>. (accessed 10 Mar 2011).
51. Shojania KG, Burton EC, McDonald KM, *et al.* Overestimation of clinical diagnostic performance caused by low necropsy rates. *Qual Saf Health Care* 2005;14:408–13.
52. Janjua KJ, Sugrue M, Deane SA. Prospective evaluation of early missed injuries and the role of tertiary trauma survey. *J Trauma* 1998;44:1000–6.
53. Kollef MH. Risk factors for the misdiagnosis of pneumothorax in the intensive care unit. *Crit Care Med* 1991;19:906–10.
54. Fabian TC, Patton JH Jr, Croce MA, *et al.* Blunt carotid injury. Importance of early diagnosis and anticoagulant therapy. *Ann Surg* 1996;223:513–22.
55. Levi AD, Hurlbert RJ, Anderson P, *et al.* Neurologic deterioration secondary to unrecognized spinal instability following trauma—a multicenter study. *Spine (Phila Pa 1976)* 2006;31:451–8.
56. Drislane FW, Lopez MR, Blum AS, *et al.* Detection and treatment of refractory status epilepticus in the intensive care unit. *J Clin Neurophysiol* 2008;25:181–6.
57. Battle RM, Pathak D, Humble CG, *et al.* Factors influencing discrepancies between premortem and postmortem diagnoses. *JAMA* 1987;258:339–44.
58. Tavora F, Crowder CD, Sun CC, *et al.* Discrepancies between clinical and autopsy diagnoses: a comparison of university, community, and private autopsy practices. *Am J Clin Pathol* 2008;129:102–9.
59. Goldhaber SZ, Hennekens CH, Evans DA, *et al.* Factors associated with correct antemortem diagnosis of major pulmonary embolism. *Am J Med* 1982;73:822–6.
60. Singh H, Peterson LA, Thomas EJ. Understanding diagnostic errors in medicine: a lesson from aviation. *Qual Saf Health Care* 2006;15:159–64.
61. Schiff GD, Hasan O, Kim S, *et al.* Diagnostic error in medicine: analysis of 583 physician-reported errors. *Arch Intern Med* 2009;169:1881–7.
62. Crowther MA, Cook DJ, Griffith LE, *et al.* Deep venous thrombosis: clinically silent in the intensive care unit. *J Crit Care* 2005;20:334–40.
63. Graber ML, Kissam S, Payne VL, *et al.* Cognitive interventions to reduce diagnostic error: a narrative review. *BMJ Qual Saf* 2012;21:535–57.
64. Wichmann D, Obbelode F, Vogel H, *et al.* Virtual autopsy as an alternative to traditional medical autopsy in the intensive care unit: a prospective cohort study. *Ann Intern Med* 2012;156:123–30.
65. Schiff GD, Kim S, Abrams R, *et al.* Agency for Healthcare Research and Quality (AHRQ). Diagnosing diagnosis errors: lessons from a multi-institutional collaborative project. *Adv Patient Saf* 2005;2:255–278. <http://www.ahrq.gov/downloads/pub/advances/vol2/Schiff.pdf>. (accessed 10 Mar 2011).
66. *Special Emphasis Notice (SEN): AHRQ Announces Interest in Research on Diagnostic Errors in Ambulatory Care Settings*. <http://grants.nih.gov/grants/guide/notice-files/NOT-HS-08-002.html>. (accessed 10 Mar 2011).



## Diagnostic errors in the intensive care unit: a systematic review of autopsy studies

Bradford Winters, Jason Custer, Samuel M Galvagno, Jr, et al.

*BMJ Qual Saf* published online July 21, 2012

doi: 10.1136/bmjqs-2012-000803

---

Updated information and services can be found at:

<http://qualitysafety.bmj.com/content/early/2012/07/23/bmjqs-2012-000803.full.html>

---

*These include:*

### References

This article cites 64 articles, 5 of which can be accessed free at:  
<http://qualitysafety.bmj.com/content/early/2012/07/23/bmjqs-2012-000803.full.html#ref-list-1>

### P<P

Published online July 21, 2012 in advance of the print journal.

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

Advance online articles have been peer reviewed, accepted for publication, edited and typeset, but have not yet appeared in the paper journal. Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>