

REVIEW

Open Access

# Bowel obstruction: a narrative review for all physicians



Fausto Caterina<sup>1</sup>, Belinda De Simone<sup>1\*</sup>, Federico Coccolini<sup>1</sup>, Salomone Di Saverio<sup>1</sup>, Massimo Sartelli<sup>1</sup> and Luca Ansaloni<sup>1</sup>

## Abstract

Small and large bowel obstructions are responsible for approximately 15% of hospital admissions for acute abdominal pain in the USA and ~ 20% of cases needing acute surgical care. Starting from the analysis of a common clinical problem, we want to guide primary care physicians in the initial management of a patient presenting with acute abdominal pain associated with intestinal obstruction.

**Keywords:** Bowel obstructions, Emergency surgery, WSES guidelines, Primary care physician education

## Backgrounds

Bowel obstruction is an important cause of morbidity and mortality accounting for nearly 30,000 deaths and laparotomic appendectomy when she was 12 years old pre more than \$3 billion per year in direct medical costs; it sends to the emergency department with intermittent acute abdominal pain and vomiting. The last defecation was 2 days ago, and the bowel is closed to gas. Current medications cases needing acute surgical care [1].

Bowel obstruction etiology is based on a mechanical intrinsic luminal obstruction or extrinsic compression of the examination demonstrates pain and localized peritonism in the lower right quadrant. Laboratory tests are normal except white blood cells at 14,000 per microliter. Questions include the following: How should this patient be evaluated and treated? What is the working diagnosis? Options include stump appendicitis, right colon diverticulitis, pelvic inflammatory disease, bowel obstruction, gastroenteritis, right renal colic, right colon cancer, bowel ischemia, or inflammatory bowel disease (Fig. 1).

In 90% of cases, small bowel obstruction is caused by adhesions, hernias, and neoplasms. [Adhesive small bowel obstruction represents 55-75% of small bowel obstruction cases [2] while hernias and small bowel tumors account for the remainder [2]. Large bowel obstruction is provoked by cancer in about 60% of cases [3]. Volvulus and diverticular disease are responsible of other 30% [3]. Other various causes (carcinomatosis, endometriosis, inflammatory bowel disease stenosis, etc.) account for the remaining 10-15% of bowel obstructions. This review focuses on the management of bowel obstruction excluding duodenal mechanical obstruction to be better included in gastric outlet obstruction entity [4].

## Clinical case presentation

An 81-year-old woman with arterial hypertension and only include valsartan 80 mg daily. The body temperature is 37.5 °C, and all vital parameters are normal. The remainder of the examination demonstrates pain and localized peritonism in the lower right quadrant. Laboratory tests are normal except white blood cells at 14,000 per microliter. Questions include the following: How should this patient be evaluated and treated? What is the working diagnosis? Options include stump appendicitis, right colon diverticulitis, pelvic inflammatory disease, bowel obstruction, gastroenteritis, right renal colic, right colon cancer, bowel ischemia, or inflammatory bowel disease (Fig. 1).

## Discussion: strategies and evidence

### Initial patient assessment

A complete history along with physical exam and laboratory tests should be performed upon presentation to the emergency unit. Patients should be asked about their last defecation/bowel gas passage. Having a history of previous abdominal surgery has 85% sensitivity and 78% specificity to predict adhesive small bowel obstruction [5]. Previous diverticulitis episodes or chronic constipation history (dolichosigmoid) may suggest diverticular stenosis and volvulus, respectively. Previous events of rectal bleeding and unexplained weight loss are suggestive of colorectal cancer. Coexisting cardiopulmonary, renal, or hepatic comorbidities

\* Correspondence: [desimoneb@hotmail.it](mailto:desimoneb@hotmail.it)

<sup>1</sup>Emergency and Trauma Surgery Department, Parma University Hospital, Gramsci 14, 43126 Parma, Italy

Full list of author information is available at the end of the article



Table 1 Causes of bowel obstruction in adults

Small bowel obstruction cause	Percentages of cases	Large bowel obstruction cause	Percentages of cases
Adhesions	55	Cancer	60
Hernias	15	Volvulus	15
Malignancies	10	Diverticular	10
Others*	15	Others*	10

Others\*: carcinomatosis, endometriosis, inflammatory bowel disease stenosis, intussusception, ischemic stenosis, radiation stenosis, colonic stenosis, gallstones, foreign bodies, bezoars

also require caution because they are associated with an increased surgical risk and may influence management strategies. Medications that could affect peristalsis are important for differential diagnosis because they are associated with bowel/large bowel volvulus) or perforation. pseudo-obstruction and adynamic ileus.

In bowel obstruction, abdominal pain is classically a colic onset due to an increase in motility to overcome occlusion. detect a strong predictive sign such as abdominal



Fig. 1 Management strategy of bowel obstruction (for about 90% of causes)

distension (sudden onset for volvulus or progressive for colorectal cancer) with a positive likelihood ratio of 16.8 and negative likelihood ratio of 0.27 [10]; peritonism signs are conversely associated with ischemia and/or perforation. Each hernia orifice (umbilical, inguinal, femoral) and all laparotomic/laparoscopic incision scars should be carefully examined. Digital rectal examination and rectoscopy can be useful in patients to detect blood or a rectal mass suggestive of colorectal malignancy.

Anamnesis and clinical examination can be very difficult in elderly or unconscious patients. In these patients is fundamental the evaluation of vital signs with a cardiopulmonary examination: severe bowel obstruction can cause hypovolemic shock and in case of perforation, septic shock. Abnormal vital signs or the general appearance of the patient including facial expression, skin color and temperature, and altered mental activity should alert the clinician that a patient may be in critical conditions. The most common signs of shock include tachycardia, tachypnea, cool extremities, mottled or cyanotic skin, slow capillary refill, and oliguria.

A complete blood count, renal function and electrolytes (to exclude pre-renal acute renal failure), and liver function tests are suggested as the first laboratory tests. Low serum bicarbonate levels, low arterial blood pH, high lactic acid level, marked leukocytosis, and hyperamylasemia may be useful in the diagnosis of intestinal ischemia. A coagulation profile should be also tested because of the potential need for emergency surgery.

Initial management

Supportive treatment must begin as soon as possible with intravenous crystalloids, anti-emetics, and bowel rest.

Isotonic dextrose-saline crystalloid and balanced isotonic crystalloid replacement fluids containing supplemental potassium in an equivalent volume to the patient's losses are recommended. Nasogastric suction can be diagnostically useful to analyze gastric contents (a feculent gastric aspirate is a characteristic of distal small bowel obstruction or large bowel obstruction). Nasogastric suction can be also therapeutically important to prevent aspiration pneumonia decompressing the proximal bowel [1]. A Foley catheter should be also inserted to monitor urine output.

Diagnostic evaluation (Tables 2 and 3)

Abdominal plain X-ray

Abdominal plain X-ray is the first level radiologic study. In small bowel obstruction, plain abdominal radiographic findings are diagnostic in 56-60%, inconclusive in 20-30%, and misleading in 1-20% of patients [2, 13].

In one study after radiography, the sensitivity of bowel obstruction was significantly higher than after clinical

Table 2 Procedures for the evaluation and treatment of bowel obstruction

	Advantages	Disadvantages
Plain abdominal X-ray	Availability	No etiologic diagnosis
Computed tomography scan	Etiologic diagnosis	Ionizing radiations exposure
Endoscopy (large bowel obstruction only)	Endoscopic treatment	Perforation risk
Conservative treatment	Fast recovery	Failure, complications, recurrence
Surgery	Etiologic treatment	Complications, stoma risk, recurrence

evaluation only: 74% versus 57%, respectively (P < 0.01). However, the positive predictive value did not differ significantly between clinical assessment only and with plain radiographs [4].

In a review of 140 cases of suspected large bowel obstruction, the abdominal X-ray had 84% sensitivity and 72% specificity [5].

Water-soluble contrast administration X-ray

A water-soluble contrast enema has 96% sensitivity and 98% specificity in diagnosing large bowel obstruction but cannot distinguish different large bowel obstruction causes.

A small bowel follow-through with water-soluble contrast is widely used in patients for adhesive small bowel obstruction non-operative management. Several systematic reviews and meta-analyses have established the utility of water-soluble contrast agents in the diagnostic work-up of adhesive small bowel obstruction [6-18]. If the contrast has not reached the colon on an abdominal X-ray 24 h after administration, then this is highly indicative of non-operative management failure [9]. Multiple studies have shown that the use of water-soluble contrast agents accurately predicts the need for surgery with an active therapeutic role [6, 17, 20, 21].

The administration of water-soluble contrast agents in adhesive small bowel obstruction is safe in terms of morbidity and mortality, but adverse effects due to their use have been reported. Potential life-threatening complications are aspiration pneumonia and pulmonary edema. To avoid these complications, the contrast medium should be administered when the stomach has been adequately decompressed through a nasogastric tube. Another potential adverse effect is that water-soluble contrast agents, because of higher osmolarity, may further dehydrate a patient with small bowel obstruction, shifting fluids into bowel lumen; in some children and elderly adults, the loss of plasma fluid may be sufficient to cause a shock-like state [2].

The contrast medium may be administered on the dosage of 50-150 ml, either orally or via nasogastric

tube, and can be given both at immediate admission and after an attempt of initial traditional conservative treatment of 48 h. The practice of giving water-soluble contrast at 48 h may reduce both the risk of aspiration pneumonia and of dehydration because the patient should have been adequately rehydrated. In this situation, the contrast material can be diluted with water [22, 23].

Rare anaphylactoid reactions following the use of oral contrast media have been reported [22].

Caution in their administration may be warranted in patients at high risk of gastropathy [24].

#### Ultrasound

Small bowel obstruction can be diagnosed with ultrasound if there are >2.5-cm dilated loops of the bowel that are proximal to collapsed loops of bowel and if there is decreased or absent peristalsis activity [25]. Using ultrasound for small bowel obstruction diagnosis has 90% sensitivity and 96% specificity [26].

Visualization of large bowel obstruction with ultrasound is as good as computed tomography. Computed tomography is clearly superior to ultrasound in terms of the etiologic definition for both small bowel obstruction and large bowel obstruction [26, 27]. Ultrasound performs better than planar abdominal X-ray in large bowel obstruction [28].

#### Computed tomography scan

The diagnostic accuracy of computed tomography with small bowel obstruction and sigmoid volvulus. Computed tomography gives an excellent evaluation of the abdominal radiography and ultrasound (Fig 2). In addition to its higher sensitivity and specificity, an important advantage of computed tomography is its ability to provide information about the underlying cause of obstruction or to provide information about an alternative diagnosis if no signs of bowel obstruction are present. Computed tomography leads to more accurate management and assistance in preoperative planning [29].

Positive oral contrast material is not needed in the diagnosis of small bowel obstruction with computed tomography because the intraluminal fluid and gas are already present within the obstructed bowel and excellent contrast agents. If positive oral contrast material has been given in patients with small bowel obstruction, then a delayed abdominal radiograph during non-operative management can assess if the contrast material has progressed to the colon.

When doubts about the large bowel obstruction diagnosis persist, a water-soluble rectal contrast agent can be administered to better visualize obstruction [29–31]. Computed tomography can also accurately stage neoplastic bowel obstruction and identify superimposed



Fig. 2 ASBO caused by single band adhesion: CT scan evidence

complications such as intestinal perforation and peritonitis [32].

In the case of bowel obstruction, it is critical to identify ischemia and necrosis especially in adhesive small bowel obstruction and sigmoid volvulus. Computed tomography gives an excellent evaluation of the bowel wall, its vessels, and mesentery-mesocolon. Pneumatosis can identify coexistent ischemia and/or infarction.

The sensitivity varies between 75 and 100%, and specificities range from 61 to 93% [33].

The diagnosis of internal hernias is very difficult because of their nonspecific clinical picture [34]. Many different types of internal hernias have been described: paraduodenal, mesentery-related, greater omentum-related, lesser sac, mesocolon-related, pericecal, falciform ligament, pelvic internal, and Roux-en-Y anastomosis-related. An internal hernia can evolve into intestinal strangulation: accurate preoperative diagnosis is possible only with computed tomography [35].

#### Magnetic resonance imaging

To minimize the burden of ionizing radiation in children and pregnant women, magnetic resonance imaging is a valid alternative examination to computed tomography for bowel obstruction [36]: prospective study demonstrated a sensitivity of 95% and a specificity of 100% [37].

### Colonoscopy

The role of colonoscopy is limited to the diagnosis of large bowel obstruction. The goal is to exclude other causes of obstruction. Biopsy should be performed in cases of suspected malignancy when emergency surgery has not been indicated or endoscopic stent placement can be expected. Endoscopic detorsion is a simple and minimally invasive procedure with a success rate of 70 to 95% and a 4% morbidity. However, mortality is about 3% in a recent study with a recurrence rate of up to 71% [38]. In this case, carbon dioxide insufflation may be an acceptable alternative to conventional air insufflation to avoid prolonged abdominal bloating, excessive abdominal pain, and discomfort during colonoscopy. Besides, CO<sub>2</sub> self-expanding metallic stents are preferred to colostomy because they are associated with similar mortality/morbidity and reduces the risk of the bowel ischemia resulting from reduced spasm and pain [39].

### Therapy

#### Conservative (non-surgical) therapy

Conservative treatment is the cornerstone of non-operative management in all patients with adhesive small bowel obstruction unless there are signs of intestinal ischemia/perforation. Evidence for the ideal duration of it may be lacking, but most authors consider a 72-h cutoff safe and appropriate [19]. The mainstay of non-operative management is nil per os and decompression with naso-gastric suction or long intestinal tube. Surgery

There has been some discussion in the literature about the use in adhesive small bowel obstruction of long intestinal tubes: long trilumen naso-intestinal tubes are more effective than naso-gastric tubes, but they require endoscopic insertion [1].

Water-soluble contrast administration is a valid and safe treatment that correlates with a significant reduction in the need for surgery in patients with adhesive small bowel obstruction with also a significant reduction in the time to resolution and length of stay. The administration of water-soluble contrast is a secure treatment with no significant differences in complications or mortality [16, 17]. Adhesive small bowel obstruction recurrence is possible after non-operative management: 12% of non-operatively treated patients are readmitted within 1 year, and this value increases to 20% after 5 years [40]. In case of complicated hernia, a prompt manual reduction has to be attempted. Emergency surgery is needed for unsuccessful reduction [41, 42]. The same admission elective surgery is indicated for all patients submitted to successful manual reduction.

Diverticular obstruction follows multiple attacks of diverticulitis with marked fibrosis of the colon wall leading to narrowing and stricture formation; in some other cases, colonic obstruction can complicate acute diverticulitis due to edema narrowing. The site of obstruction is usually in the sigmoid colon; occlusion is normally incomplete and resolves with conservative treatment [43].

Sigmoid volvulus allows one to assess the viability of the sigmoid but to also achieve

detorsion. If colonic necrosis is present, then the patient undergoes immediate surgery. In the absence of colonic necrosis, endoscopy can convert an urgent situation into elective situation in the same admission. Colonoscopic detorsion is a simple and minimally invasive procedure with a success rate of 70 to 95% and a 4% morbidity. However, mortality is about 3% in a recent study with a recurrence rate of up to 71% [44, 45]. For palliation of obstructing left colon cancer, self-expanding metallic stents are preferred to colostomy because they are associated with similar mortality/morbidity rates but a shorter hospital stay [46]. Self-expanding metallic stents can be also a bridge to elective surgery for obstructing left colon cancer. It offers a better short-term outcome than emergency surgery because the rate of stomas is lower; long-term outcomes seem comparable, but there is still insufficient oncological evidence. Thus, self-expanding metallic stents should not be considered the treatment of choice for obstructing left colon cancer: a valid option in selected cases and in centers with significant expertise [47–50].

Prosthetic repair is the treatment of choice for most abdominal wall complicated hernias (inguinal, femoral, inguinal, umbilical, epigastric, parastomal, spigelian, etc.). In case of perforation/bowel resection with contaminated surgical fields, suture repair is preferred due to the risk of mesh infection. Diagnostic laparoscopy may be a useful tool to assess bowel viability after reduction of complicated hernias [51]. Repair of complicated hernia can be performed with a laparoscopic approach when no bowel resection anastomosis is needed, which normally requires a mini-open approach (small laparotomy) [51]. Internal hernias are treated with prompt reduction, suture repair, and bowel resection anastomosis in case of intestinal necrosis.

Historically, abdominal adhesiolysis through laparotomy has been the standard therapy for adhesive small bowel obstruction. In the case of emergent surgical exploration (i.e., perforation or bowel ischemia) or for conservative treatment failure, operative laparotomic surgery is the treatment of choice [6, 19, 52]. Laparoscopic adhesiolysis has been introduced in recent decades and can decrease morbidity in subgroups of patients undergoing surgery for adhesive small bowel obstruction. The risk of intestinal injury is higher in laparoscopic surgery for adhesive small bowel obstruction. Therefore, careful selection of patients for laparoscopic surgery is mandatory. Results of randomized trials will soon be published [19, 53, 54]. The risk of recurrence is slightly lower after operative treatment compared to non-operative treatment: 8% after 1 year and 16% after 5 years [40, 55].

Small bowel obstruction caused by small bowel tumors failed conservative treatments or as elective therapy to (adenocarcinoma, neuroendocrine tumors, gastrointestinal stromal tumors, and lymphomas) is treated with resection and anastomosis. Oncologic management of these tumors must be obviously considered following the same schemes of tumors that arise outside the small bowel [56–58].

For large bowel obstruction caused by sigmoid volvulus without ischemia or perforation, the best strategy is an endoscopic detorsion procedure followed by same admission surgery that includes a sigmoid colectomy with primary anastomosis. Exclusively endoscopic therapy without subsequent surgery must be reserved for high-surgical-risk patients. In case of ischemic volvulus or failed derotation, surgery has to be performed as soon as possible. In cecocolic volvulus, endoscopy has no role, and surgery (right hemicolectomy) is the only option [44]. The role of laparoscopic surgery for volvulus is limited: the absence of fixation of the sigmoid colon and its excessive length often make laparoscopic exposure and dissection difficult.

Resection and primary anastomosis are the desired procedure for diverticular large bowel obstruction, and it should be attempted regardless of bowel preparation after a successful conservative treatment in the same patient [59]. Exclusively conservative therapy or Hartmann procedure could be more appropriate for high-risk patients.

Resection and primary anastomosis are the best options for malignant large bowel obstruction in the absence of significant risk factors or perforations. Patients with high surgical risk or perforations are better managed with staged procedure (e.g., Hartmann procedure). Many prospective and retrospective studies into resection and primary anastomosis in malignant large bowel obstruction have reported percentages of anastomotic leaks ranging from 2.2 to 12% [60, 61] comparable to the 2–8% rate after elective surgical procedures. In the case of large bowel obstruction caused by extraperitoneal rectal cancer, resection of the primary tumor should be postponed and a stoma should be fashioned to permit a correct staging and a more appropriate oncological neoadjuvant treatment. Laparoscopy in emergency treatment of malignant large bowel obstruction should be reserved to selected cases in specialized centers [62–64].

Other uncommon bowel obstructions  
 Carcinomatosis, endometriosis, inflammatory bowel disease stenosis, intussusception, post-ischemic stenosis, radiation stenosis, postanastomotic stenosis, gallstones, foreign bodies, bezoars, and tuberculosis can cause bowel obstruction in a minority of cases (globally 10–15%). Here, a computed tomography scan is critical for diagnosis. Conservative treatment should be started in the absence of ischemia or perforation, but surgery is needed as a rescue therapy for

precisely the role of conservative treatment in adhesive small bowel obstruction and its length. Moreover, a great debate is underway about laparoscopic adhesiolysis po- endoscopic detorsion procedure followed by same admis- tentialities against possible severe technical complica- tion surgery that includes a sigmoid colectomy with pri- tions. Both issues are critical to decrease the well-known mary anastomosis. Exclusively endoscopic therapy without adhesive small bowel obstruction recurrence risk. Stents will probably become the mainstay of cancer large bowel obstruction transforming emergency operations in elect- surgery has to be performed as soon as possible. In cecal cases decreasing complications and stomas.

Guidelines  
 The recommendations in this article are concordant with guidelines published by the World Society of Emer- gency Surgery [9] (Table 3).

**Conclusions**

Bowel obstruction is the most likely diagnosis in the pa- tient described in the vignette. High white blood cells and peritonism could suggest adhesive small bowel obstruc- tion with possible ischemia. A computed tomography scan with intravenous contrast must be performed as soon as possible. If computed tomography scan confirms adhesive small bowel obstruction with ischemia or perforation, then the patient must go to surgery as soon as possible: a laparoscopic approach should be attempted.

If computed tomography shows adhesive small bowel obstruction without ischemia or perforation, then a con- servative treatment should be initiated: nasogastric suc- tion and fluid replacement therapy have to be performed with a scrupulous wait-and-see strategy. After gastric contents are cleared, a water-soluble contrast adminis- tration challenge should be performed. The patient has

**Table 3 Key clinical points**

- Multidetector computed tomography has emerged as the best imaging test for the diagnosis of mechanical bowel obstruction and its complications and can help patients management to either conservative or operative management
- Conservative adhesive small bowel obstruction treatment is the mainstay non-operative management in all patients with adhesive small bowel obstruction without signs of perforations or bowel ischemia
- Self-expanding metallic stents as bridge to elective surgery
- Constructing left colon cancer offers a better short-term outcome than direct emergency surgery because morbidity is comparable, but rate of stomas is significantly lower; long-term outcomes seem also comparable, but there is still insufficient oncologic scientific evidence to prove it
- Colonoscopic sigmoid volvulus detorsion is a simple and mini-invasive procedure associated with a success rate of 70 to 95% with a 4% morbidity: it can convert an urgent situation into an elective one but with high recurrence rate.

to be monitored with regard to ischemia (peritonism, white blood cells, lactate). Surgery has to be performed immediately in the case of clinical deterioration, preferably beginning with a laparoscopic technique.

After 24 h, a plain abdominal X-ray should be performed to determine if oral contrast reached the large bowel: if positive, then oral nutrition can be started. If negative, non-operative management could be continued for another 48 h: surgery should be performed after this limit, preferably starting with a laparoscopic approach.

**Abbreviations**

ASBO Adhesive small bowel obstruction; CT: Computed tomography; SBO: Small bowel obstructions

**Acknowledgements**

Not applicable

**Funding**

No authors received any funding resource. The paper received a WSES Institutional waiver for this publication.

**Availability of data and materials**

Not applicable

**Authors' contributions**

FC conceived the study, collected the data, and wrote the manuscript; all the authors reviewed and approved the final manuscript.

**Ethics approval and consent to participate**

Not applicable

**Consent for publication**

Not applicable

**Competing interests**

The authors declare that they have no competing interests.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Author details**

<sup>1</sup>Emergency and Trauma Surgery Department, Parma University Hospital, Via Gramsci 14, 43126 Parma, Italy; Emergency and Trauma Surgery Department, Cesena Hospital, Cesena, Italy; General Surgery Department, Cambridge University Hospital, Cambridge, UK; General Surgery Department, Macerata Hospital, Macerata, Italy.

Received: 3 January 2019 Accepted: 10 April 2019

**References**

1. Cappell MS, Batke M. Mechanical obstruction of the small bowel and colon. *Med Clin North Am.* 2008;92(3):575
2. Gore RM, Silvers RI, Thakrar KH, et al. Bowel obstruction. *Radiol Clin N Am.* 2015;53(6):1225
3. Batke M, Cappell MS. Adynamic ileus and acute colonic pseudo-obstruction. *Med Clin North Am.* 2008;92(3):679
4. Saunders MD, Kimmey MB. Systematic review: acute colonic pseudo-obstruction. *Aliment Pharmacol Ther.* 2005;22:2517
5. Miller G, Boman J, Shrier I, Gordon PH. Etiology of small bowel obstruction. *Am J Surg.* 2000;180(1):53
6. Ten Broek RP, Issa Y, van Santbrink EJ, et al. Burden of adhesions in abdominal and pelvic surgery: systematic review and meta-analysis. *BMJ.* 2013;347:f5588.
7. Frago R, Ramirez E, Millan M, Kreisler E, del Valle E, Biondo S. Current management of acute malignant large bowel obstruction: a systematic review. *Am J Surg.* 2014;207(1):1238.

8. Roses RE, Folkert IW, Krouse RS. Malignant bowel obstruction: reappraising the value of surgery. *Surg Oncol Clin N Am.* 2018;27(4):5705
9. Taylor MR, Lalani N. Adult small bowel obstruction. *Acad Emerg Med.* 2013;20:52844.
10. Bohner H, Yang Q, Franke C, Verreet PR, Ohmann C. Simple data from history and physical examination help to exclude bowel obstruction and to avoid radiographic studies in patients with acute abdominal pain. *Eur J Surg.* 1998;164:737.
11. Chen XL, Ji F, Lin Q, Chen YP, et al. A prospective randomized trial of transnasal ileus tube vs nasogastric tube for adhesive small bowel obstruction. *World J Gastroenterol.* 2012;18(16):7468
12. Gans SL, Stoker J, Boermeester MA. Plain abdominal radiography in acute abdominal pain; past, present, and future. *Int J Gen Med.* 2012;3:525
13. Frager D. Intestinal obstruction role of CT. *Gastroenterol Clin N Am.* 2002;31(3):7799.
14. van Randen A, Laméris W, Luitse JS et al.; OPTIMA study group. The role of plain radiographs in patients with acute abdominal pain at the ED. *Am J Emerg Med* 2011;29(6):529.5825e2.
15. Chapman AH, McNamara M, Porter G. The acute contrast enema in suspected large bowel obstruction: value and technique. *Clin Radiol.* 1992;46(4):273.
16. Ceresoli M, Coccolini F, Catena F, et al. Water-soluble contrast agent in adhesive small bowel obstruction: a systematic review and meta-analysis of diagnostic and therapeutic value. *Am J Surg.* 2016;211(6):5114
17. Branco BC, Barmparas G, Schnuriger B, Inaba K, Chan LS, Demetriades D. Systematic review and meta-analysis of the diagnostic and therapeutic role of water-soluble contrast agent in adhesive small bowel obstruction. *Br J Surg.* 2010;97(4):430
18. Abbas S, Bissett IP, Parry BR. Oral water soluble contrast for the management of adhesive small bowel obstruction. *Cochrane Database Syst Rev.* 2007;(3):CD004651. <https://doi.org/10.1002/14651858.CD004651.pub3>
19. Ten Broek RPG, Krielen P, Di Saverio S, et al. Bologna guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2017 update of the evidence-based guidelines from the world society of emergency surgery ASBO working group. *World J Emerg Surg.* 2018;19(13):24.
20. Di Saverio S, Catena F, Ansaloni L, Gavioli M, Valentino M, Pinna AD. Water-soluble contrast medium (gastrografin) value in adhesive small intestine obstruction (ASIO): a prospective, randomized, controlled, clinical trial. *World J Surg.* 2008;32(10):2334.
21. Keenan JE, Turley RS, McCoy CC, Migaly J, Shapiro ML, Scarborough JE. Trials of nonoperative management exceeding 3 days are associated with increased morbidity in patients undergoing surgery for uncomplicated adhesive small bowel obstruction. *J Trauma Acute Care Surg.* 2014;76(6):1367
22. <https://www.acr.org/Clinical-Resources/Contrast-Manual>
23. Di Saverio S, Coccolini F, Galati M, et al. Bologna guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2013 update of the evidence-based guidelines from the world society of emergency surgery ASBO working group. *World J Emerg Surg.* 2013;8(1):42. Published 2013 Oct 1. <https://doi.org/10.1186/1749-7922-8-42>
24. Guy S, Al Askari M. Haemorrhagic gastritis following Gastrografin administration for adhesive small bowel obstruction: a case report of a rare outcome. *Int J Surg Case Rep.* 2017;34:51
25. Ogata M, Mateer JR, Condon RE. Prospective evaluation of abdominal sonography for the diagnosis of bowel obstruction. *Ann Surg.* 1996;223:237
26. Suri S, Gupta S, Sudhakar PJ, Venkataramu NK, Sood B, Wig JD. Comparative evaluation of plain films, ultrasound and CT in the diagnosis of intestinal obstruction. *Acta Radiol.* 1999;40(4):322
27. Hollerweger A, Wüstner M, Dirks K. Bowel obstruction: sonographic evaluation. *Ultraschall Med.* 2015;36(3):3516
28. Chen SC, Yen ZS, Wang HP, Lin FY, Hsu CY, Chen WJ. Ultrasonography is superior to plain radiography in the diagnosis of pneumoperitoneum. *Br J Surg.* 2002;89(3):381
29. Jaffe T, Thompson WM. Large-bowel obstruction in the adult: classic radiographic and CT findings, etiology, and mimics. *Radiology.* 2015;275(3):651
30. O'Malley RG, Al-Hawary MM, Kaza RK, Wasnik AP, Platt JF, Francis IR. MDCT findings in small bowel obstruction: implications of the cause and presence of complications on treatment decisions. *Abdom Imaging.* 2015;40(7):2248
31. Millet I, Taourel P, Ruyet A, Molinari N. Value of CT findings to predict surgical ischemia in small bowel obstruction: a systematic review and meta-analysis. *Eur Radiol.* 2015;25(6):1623

32. Brown CV. Small bowel and colon perforation. *Surg Clin North Am.* 2014; 94(2):471–5.
33. Paulson EK, Thompson WM. Review of small-bowel obstruction: the diagnosis and when to worry. *Radiology.* 2015;275(2):332–42.
34. Turenkov A, Kelley KM, Boustany M. Internal hernia as cause of small bowel obstruction without history of gastrointestinal surgery. *Am Surg.* 2018;84(7):e231–3.
35. Doishita S, Takeshita T, Uchima Y, et al. Internal hernias in the era of multidetector CT: correlation imaging and surgical findings. *Radiographics.* 2016;36(1):88–106.
36. Beddy P, Keogan MT, Sala E, Griffin N. Magnetic resonance imaging for the evaluation of acute abdominal pain in pregnancy. *Semin Ultrasound CT MR.* 2010;31(5):433–41.
37. Beall DP, Fortman BJ, Lawler BC, Regan F. Imaging bowel obstruction: a comparison between fast magnetic resonance imaging and helical computed tomography. *Clin Radiol.* 2002;57(8):719–24.
38. Markogiannakis H, Messaris E, Dardamanis D, et al. Acute mechanical bowel obstruction: clinical presentation, etiology, management and outcome. *World J Gastroenterol.* 2007;13(3):432–7.
39. Sajid MS, Caswell J, Bhatti MI, Sains P, Baig MK, Miles WF. Carbon dioxide insufflation vs conventional air insufflation for colonoscopy: a systematic review and meta-analysis of published randomized controlled trials. *Color Dis.* 2015;17:111–23. <https://doi.org/10.1111/codi.12837>.
40. Foster NM, McGory ML, Zingmond DS, Ko CY. Small bowel obstruction: a population-based appraisal. *J Am Coll Surg.* 2006;203(2):170–6.
41. Fitzgibbons RJ Jr, Forse RA. Clinical practice. Groin hernias in adults. *N Engl J Med.* 2015;372(8):756–63.
42. Bower KL, Lollar DI, Williams SL, Adkins FC, Luyimbazi DT, Bower CE. Small bowel obstruction. *Surg Clin North Am.* 2018;98(5):945–71.
43. Psaras K, Symeonidis NG, Pavlidis ET, et al. *Tech Coloproctol.* 2011;15(Suppl 1):S9–12.
44. Perrot L, Fohlen A, Alves A, Lubrano J. Management of the colonic volvulus in 2016. *J Visc Surg.* 2016;153(3):183–92.
45. Bauman ZM, Evans CH. Volvulus. *Surg Clin North Am.* 2018;98(5):973–93.
46. Takahashi H, Okabayashi K, Tsuruta M, Hasegawa H, Yahagi M, Kitagawa Y. Self-expanding metallic stents versus surgical intervention as palliative therapy for obstructive colorectal cancer: a meta-analysis. *World J Surg.* 2015;39(8):2037–44.
47. Ceresoli M, Allievi N, Coccolini F, et al. Long term oncologic outcomes of stents as a bridge to surgery vs. emergency surgery in malignant left side colonic obstruction: a meta-analysis. *J Surg Oncol.* 2017;8:867–76.
48. Young CJ, De-Loyde KJ, Young JM, et al. Improving quality of life for people with incurable large-bowel obstruction: randomized control trial of colonic stent insertion. *Dis Colon Rectum.* 2015;58(9):838–49.
49. Atukorale YN, Church JL, Hoggan BL, et al. Self-expanding metallic stents for the management of emergency malignant large bowel obstruction: a systematic review. *J Gastrointest Surg.* 2016;20(2):455–62.
50. Cennamo V, Luigiano C, Manes G, et al. Colorectal stenting as a bridge to surgery reduces morbidity and mortality in left-sided malignant obstruction: a predictive risk score-based comparative study. *Dig Liver Dis.* 2012;44(6):508–14.
51. Birindelli A, Sartelli M, Di Saverio S, et al. 2017 update of the WSES guidelines for emergency repair of complicated abdominal wall hernias. *World J Emerg Surg.* 2017;7(12):37.
52. Rami Reddy SR, Cappell MS. A systematic review of the clinical presentation, diagnosis, and treatment of small bowel obstruction. *Curr Gastroenterol Rep.* 2017;19(6):28.
53. Sajid MS, Khawaja AH, Sains P, Singh KK, Baig MK. A systematic review comparing laparoscopic vs open adhesiolysis in patients with adhesional small bowel obstruction. *Am J Surg.* 2016;212(1):138–50.
54. Wiggins T, Markar SR, Harris A. Laparoscopic adhesiolysis for acute small bowel obstruction: systematic review and pooled analysis. *Surg Endosc.* 2015;29(12):3432–42.
55. Catena F, Ansaloni L, Di Saverio S, Pinna AD; World Society of Emergency Surgery. P.O.P.A. study: prevention of postoperative abdominal adhesions by icodextrin 4% solution after laparotomy for adhesive small bowel obstruction. A prospective randomized controlled trial. *J Gastrointest Surg.* 2012;16(2):382–388.
56. Lepage C, Bouvier AM, Manfredi S, Dancourt V, Faivre J. Incidence and management of primary malignant small bowel cancers: a well-defined French population study. *Am J Gastroenterol.* 2006;101(12):2826–32.
57. Catena F, Ansaloni L, Gazzotti F, et al. Small bowel tumours in emergency surgery: specificity of clinical presentation. *ANZ J Surg.* 2005;75(11):997–9.
58. Vallicelli C, Coccolini F, Catena F, et al. Small bowel emergency surgery: literature's review. *World J Emerg Surg.* 2011;7(6):1:1.
59. Dozois EJ. Operative treatment of recurrent or complicated diverticulitis. *J Gastrointest Surg.* 2008;12(8):1321–3.
60. Kim J, Mittal R, Konyalian V, King J, Stamos MJ, Kumar RR. Outcome analysis of patients undergoing colectorectal resection for emergent and elective indications. *Am Surg.* 2007;73(10):991–3.
61. Bellows CF, Webber LS, Albo D, Awad S, Berger DH. Early predictors of anastomotic leaks after colectomy. *Tech Coloproctol.* 2009;13(1):41–7.
62. Yeo HL, Lee SW. Colorectal emergencies: review and controversies in the management of large bowel obstruction. *J Gastrointest Surg.* 2013;17(11):2007–12.
63. Pisano M, Zorcolo L, Merli C, et al. 2017 WSES guidelines on colon and rectal cancer emergencies: obstruction and perforation. *World J Emerg Surg.* 2018;13(13):36.
64. Catena F, Pasqualini E, Tonini V, Avanzolini A, Campione O. Emergency surgery for patients with colorectal cancer over 90 years of age. *Hepatogastroenterology.* 2002;49(48):1538–9.
65. Paul Olson TJ, Pinkerton C, Brasel KJ, Schwarze ML. Palliative surgery for malignant bowel obstruction from carcinomatosis: a systematic review. *JAMA Surg.* 2014;149(4):383–92.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

