

*Jerrold H. Levy, M.D., F.A.H.A., F.C.C.M., Editor*

## Early Hemodynamic Management of Critically Ill Burn Patients

Sabri Soussi, M.D., François Dépret, M.D., Mourad Benyamina, M.D., Matthieu Legrand, M.D., Ph.D.

**B**URN injury is associated with early profound hypovolemia followed by a systemic inflammatory response with a subsequent hyperdynamic state.<sup>1</sup> Hemodynamic management has long been identified as a key factor impacting burn patients' prognosis.<sup>2</sup> Because both under- and over-resuscitation may potentially negatively impact outcome, anesthesiologists and intensivists caring for burn patients will have to face the challenge of fluid balance in these patients.<sup>3,4</sup> The aim of this review is to provide an overview of the hemodynamic consequences of burn injury and to propose strategies for the initial hemodynamic management of severe burn patients using the available evidence-based medicine combined with a physiologic approach.

### Cardiovascular Consequences of Burn Injury

#### Burn Edema Process

Severe burn injury leads to tissue destruction with capillary leak. Edema starts within the first hour after injury in burned tissues. Thereafter, a slower increase in fluid extravasation in both injured and noninjured tissues occurs during the first 24 to 48 h after burn injury when systemic inflammatory response occurs.<sup>5</sup>

The pathogenesis of burn edema involves all the physical forces described by the conventional Starling law. Fluid movement into the interstitium results from the combination of an imbalance between hydrostatic and oncotic forces, together with increase of vascular permeability. The development of gaps in endothelial cell junction with an increase in capillary permeability favors fluid and protein loss into the interstitium in burn tissue and only fluid in the nonburned tissue (fig. 1).<sup>5</sup>

Capillary filtration and reabsorption are also strongly affected by the integrity of the glycocalyx.<sup>6</sup> The endothelial glycocalyx is a dynamic structure composed of cell-bound proteoglycans and sialoproteins (1 to 3  $\mu\text{m}$  in thickness) that envelop endothelial cells on their luminal side and inside the endothelial paracellular clefts. It plays a central role in vascular permeability (mainly paracellular permeability) by maintaining the oncotic gradient across the endothelial barrier. Accordingly, the oncotic gradient between the plasma and interstitial spaces is set between plasma and glycocalyx (subglycocalyx oncotic pressure) rather than being transendothelial as described in the traditional Starling model.<sup>7</sup> Inflammatory injury (sepsis, trauma, thermal injury) to the glycocalyx increases paracellular permeability, which is associated with fluid and albumin leak into the interstitial space (fig. 1).<sup>6</sup>

Urinary microalbuminuria was described as a marker of systemic endothelial permeability.<sup>8</sup> In an experimental model of sepsis, albumin leakage into urine increased in association with glomerular barrier glycocalyx alteration.<sup>9</sup> Syndecan-1 and endocan have been proposed as other biomarkers of glycocalyx degradation and endothelial injury in septic and trauma patients.<sup>10</sup> In a recent study using syndecan-1 as a surrogate marker of glycocalyx shedding in a population of burn patients, Osuka *et al.* found that after adjustment for age, sex, percent total body surface area, and inhalation injury, syndecan-1 was an independent marker of the increase in fluid requirements ( $P = 0.04$ ).<sup>11</sup>


#### Hemodynamic Patterns of Burn Shock

Burn injury leads to initial systemic and pulmonary vasoconstriction, (related to catecholaminergic release and hemolysis), low cardiac output (CO), and low oxygen delivery and


Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are available in both the HTML and PDF versions of this article. Links to the digital files are provided in the HTML text of this article on the Journal's Web site ([www.anesthesiology.org](http://www.anesthesiology.org)).

Submitted for publication December 1, 2017. Accepted for publication May 7, 2018. From the Assistance Publique-Hôpitaux de Paris, Groupe Hospitalier, Saint Louis-Lariboisière, Department of Anesthesiology and Critical Care and Burn Unit, Paris, France (S.S., F.D., M.B., M.L.); University Paris Diderot, Paris, France (F.D., M.L.); and Unité Mixte de Recherche, Institut National de la Santé et de la Recherche Médicale 942, Paris, France (F.D., M.L.).

Copyright © 2018, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2018; XXX:00–00



**Fig. 1.** Pathophysiology of capillary leak syndrome in critically ill burns. Fluid accumulation into the interstitium results from the combination of an imbalance between hydrostatic and oncotic forces favoring the fluid movement into the interstitium together with increase of vascular permeability and **glycocalyx degradation**. **Red arrows** indicate intravascular hydrostatic pressure, **blue arrows** indicate oncotic pressure, and **green broken lines** indicate glycocalyx.  $\Pi$  = Oncotic pressure.



**Fig. 2.** Burn shock hemodynamic pattern in the first 48h. In the first hours, burn injury leads to initial low cardiac output (CO; A) and systemic and pulmonary vasoconstriction (B). Within 24 to 48h, a hyperdynamic and vasoplegic state develops, which is characterized by a high CO (A) and decreased systemic and pulmonary vascular resistances (B). This pattern was observed with or without fluid resuscitation. Fluid resuscitation was, however, associated with a faster restoration of CO as well as higher CO during the hyperdynamic phase, especially when albumin is infused with crystalloids. Figure adapted from Asch *et al.*<sup>1</sup>

consumption.<sup>1</sup> Within 24 to 48h, a hyperdynamic and vasoplegic state develops, which is characterized by a high CO, increased oxygen consumption, and decreased systemic vascular resistance.<sup>12,13</sup> Interestingly, this pattern was observed with or without fluid resuscitation in animals. Fluid resuscitation was, however, associated with a faster restoration of CO as well as a higher CO during the hyperdynamic phase (fig. 2).<sup>1</sup>

### Burn-associated Cardiac Injury

Severe thermal injury can also induce cardiac dysfunction with impaired left ventricular systolic function, slowed isovolemic relaxation, and decreased diastolic compliance.<sup>14</sup> Cardiac stress can result in left heart failure and increased left cardiac filling pressures, which may further promote extravascular fluid transfer into the lungs. An increase in right ventricular workload may also lead to right heart failure, increase in right ventricular filling pressure, and venous congestion.<sup>15</sup>

### Microcirculatory Alterations

Alterations of microcirculatory blood flow can further compromise organ perfusion. These microcirculatory derangements

were described mainly in the splanchnic territory and the skin (in both injured and noninjured tissues) using gastric tonometry and skin laser Doppler and microdialysis, respectively.<sup>16,17</sup> Microcirculatory derangements were associated with organ failure and poor outcome.<sup>13</sup> The development of tissue edema with fluid resuscitation may lead to venous occlusion and increased vascular resistance as well as disseminated intravascular coagulation in the microvasculature and altered erythrocyte deformability.<sup>18</sup> Furthermore, burn-related hemolysis and the use of hydroxocobalamin after inhalation injury may lead to vasoconstriction and compromise the microperfusion through vasoconstriction due to the scavenging effect of nitric oxide.<sup>19,20</sup>

### Risks of Fluid Under- and Over-resuscitation in Burn Patients


The main challenge in the initial fluid administration strategy is to avoid significant hypovolemia, which may induce hypoperfusion, but without over-resuscitating the patient. Under-resuscitation may lead to acute kidney injury and nonocclusive mesenteric ischemia.<sup>3,21</sup> On the other hand,

the risk of abdominal compartmental syndrome, acute kidney injury, and acute respiratory distress syndrome (ARDS) are the most easily identified consequences of fluid over-resuscitation.<sup>4,22</sup> Abdominal compartmental syndrome and ARDS result from the combination of large fluid resuscitation volumes and systemic capillary leak syndrome. In a pediatric study, patients who received large volumes of fluid resuscitation represented an at-risk group for respiratory failure after a scald injury (7.66 vs. 4.07 ml/kg per percent total body surface area,  $P < 0.001$ ).<sup>23</sup> Another study confirmed that higher crude rates of ARDS were observed in patients who received standard or excessive resuscitation (greater than 4 ml/kg per percent total body surface area) than those patients who received restricted resuscitation.<sup>4</sup> Furthermore, the association between a large positive fluid balance and the onset of acute kidney injury was reported in severe burns and suggests a role of venous congestion in the development of acute kidney injury.<sup>24</sup> Interestingly, hypervolemia

promotes fluid extravasation into the interstitial space. A rapid rise in intravascular hydrostatic pressure will increase the hydrostatic gradient pressure and therefore increase transcapillary fluid extravasation according to the Starling forces. A decrease in intravascular oncotic pressure due to hemodilution after crystalloids infusion will further promote extravascular fluid leakage. Finally, hypervolemia has been shown to alter the glycocalyx and therefore increase vascular permeability.<sup>11,25</sup> Therefore, except for critical hypovolemic hypotension, rapid large-volume fluid boluses might be undesirable in order to limit these three factors promoting extravascular fluid leakage (fig. 3).

### Hemodynamic Targets in Early Resuscitation of Critically Ill Burn Patients

The Parkland formula is most commonly used to predict the required fluid volumes. Although the Parkland formula



**Fig. 3.** Saint Louis Hospital hemodynamic management algorithm. Cardiac output, invasive arterial pressure, and central venous pressure (CVP) are monitored in patients with greater than 30% total body surface area burned (TBSA), or greater than 20% TBSA in patients with comorbidities, or severe smoke inhalation injury or patients with hemodynamic instability. Start resuscitation with  $2 \text{ ml} \cdot \text{kg}^{-1} \cdot \% \text{ TBSA}^{-1} \cdot 8 \text{ h}^{-1}$  of crystalloids with four main goals (urinary output [UO] greater than  $0.5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ , cardiac index greater than  $2.2 \text{ l} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ , central venous oximetry [ScvO<sub>2</sub>] greater than 70%, lactate decrease). If one of these targets is not reached with CVP less than 6 mmHg or positive dynamic tests (passive leg raising [PLR], pulse pressure variation [PPV], stroke volume variation [SVV]) or mean arterial pressure (MAP) less than 65 to 70 mmHg, test fluid responsiveness or perform a fluid challenge. If the patient is not a fluid responder, then introduce norepinephrine, perform an echocardiography, and search for a cardiac dysfunction. If the patient is a fluid responder, then step up and increase IV crystalloids rate by 20% and then recheck each target hourly. If all targets are reached, then step down IV crystalloids rate by 10% every hour and recheck each target hourly. \*If the patient has acute respiratory distress syndrome, use inhaled nitric oxide? Perform prone position? PAC = pulmonary artery catheter; RAP = right atrial pressure; RL = Ringer's Lactate.

has long been considered as the reference, it remains rough and inaccurate for several reasons. First, there are obvious interindividual variations between fluid requirements due to different host responses to burn injury and/or smoke inhalation injuries and extreme ages and comorbidities. Many factors other than percent total body surface area burn and body weight may influence fluid requirements in severely burned patients: burn depth, burn mechanism, associated trauma, early escharotomy or fasciotomy, and inhalation injury. Higher fluid requirements may result from the intensive inflammatory response with capillary leak after inhalation injury. Finally, estimating the body weight and the burn total body surface area can be challenging, and errors in such evaluation can lead to errors in the fluid requirements estimation.

Of note, using the Parkland formula, half of the volume should be given within 8 h of the burn injury and half during the next 16 h. This only approximates the fluid needs during the very first hours. This approach also overlooks the dynamic nature of the circulation during the first 24 h (fig. 2). In a study including critically ill burn patients, stroke volume and cardiac index within the first hour after admission were independently associated with 90-day mortality regardless of total body surface area.<sup>26</sup> For these reasons, we strongly recommend hourly assessment of hemodynamics during the first 24 h of burn shock with hourly adjustment of fluid needs (fig. 3). Fluid volume requirements are routinely based on mean arterial pressure and urine output. However, these physiologic targets have been recognized as poor resuscitation endpoints, not accurately reflecting CO or the adequacy of oxygen delivery.<sup>27</sup> A decrease in urine output may be associated with a prerenal cause (e.g., a decrease in renal blood flow or renal perfusion pressure due to fluid depletion), but other causes of acute kidney injury in burn patients include hemolysis, rhabdomyolysis, inflammation and immune-related response, and intrarenal vasoconstriction due to neurohormonal activation, which may lead to an uncoupling between intravascular volume and urine output. In contrast, venous congestion related to an increased venous pressure due to increased cardiac filling pressure and/or increased abdominal pressure can lead to oliguria. In these settings, fluid loading would obviously not represent the right response to low urine output. On the other hand, an increase in urine output and glomerular filtration rate may be observed after a rapid drop in plasma albumin concentration, increasing the glomerular transcapillary fluid rate fraction or hyperglycemia-related hyperosmolarity.<sup>28</sup> In a retrospective study, variations in urine output and vital signs were poorly correlated with oxygen parameter changes (oxygen delivery and consumption) after fluid loading.<sup>27</sup> Sánchez *et al.*, in a prospective cohort study, reported that oliguric patients did not show differences in cardiac index, intrathoracic blood volume, or lactate concentrations compared to nonoligurics.<sup>29</sup> To summarize, low urine output (e.g., less than  $0.5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ) represents an alarm sign

that should prompt rapid investigation of the hemodynamic status. However, urine output *per se* cannot be considered as a reliable parameter of hypovolemia.<sup>30</sup>

### Goal-directed Resuscitation Therapy

Goal-directed fluid resuscitation therapy based on more advanced hemodynamic monitoring to reach alternative targets has been suggested to better tailor fluid resuscitation.<sup>31,32</sup> Serum lactate and base deficit have been found to be associated with outcomes in major burns, and proposed as endpoints of initial resuscitation.<sup>29,33</sup> Low CO and oxygen delivery were also associated with poor outcome in critically ill burn patients.<sup>13,26</sup>

Only a few randomized controlled trials have compared fluid resuscitation based on the Parkland formula with a goal-directed therapy strategy. Csontos *et al.* compared the effect of two resuscitation regimens (based on urine output *vs.* intrathoracic blood volume) on the multiple organ dysfunction syndrome and central venous oximetry in the first 72 h.<sup>34</sup> The mean central venous oximetry was significantly lower in the urine output group than in the intrathoracic blood volume group (68% [64 to 71] and 74% [71 to 78], respectively;  $P = 0.024$ ) for the first 24 h. The multiple organ dysfunction syndrome score was significantly higher in the urine output group than in the intrathoracic blood volume group at 48 h (5 [4.3 to 5.8] *vs.* 4 [3 to 4.3], respectively;  $P = 0.024$ ) and 72 h after injury (5 [4.3 to 6] *vs.* 3 [3.3 to 3.8], respectively;  $P = 0.014$ ) without any impact on mortality.

In a randomized controlled trial using continuous real-time CO monitoring with arterial pulse counter analysis and an algorithm testing fluid responsiveness with dynamic preload parameters, crystalloid volume administration was found to be lower than in the control group ( $5,090 \pm 680 \text{ ml}$  *vs.*  $7,820 \pm 1,050 \text{ ml}$ ,  $P = 0.04$ ) while no difference was found in relation to organ dysfunction and mortality.<sup>35</sup> Arlati *et al.* observed that fluid administration guided by a hemodynamics-oriented approach in the first 24-h period limited to a urine output of  $0.5$  to  $1 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$  and a cardiac index of at least  $2.2 \text{ l} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  was safe, resulting in lower fluid volume administration and less organ dysfunction.<sup>2</sup> In a retrospective case series including burn patients, a goal-directed therapy strategy was associated with better outcome than conventional management.<sup>32</sup> In a systematic review and meta-analysis including 20 studies, a decrease in mortality was found with the use of certain hemodynamic alternative endpoints (mainly cardiac index and intrathoracic blood volume) rather than hourly urine output (risk ratio, 0.77; 95% CI, 0.42 to 0.85;  $P < 0.004$ ).<sup>31</sup>

In contrast, studies that have used a fluid resuscitation strategy mainly based on normalization of static preload parameters (intrathoracic blood volume greater than  $800 \text{ ml/m}^2$ ) found a significant increase in fluid administration in comparison with using the Parkland formula. In a study including 50 patients with a total body surface area greater than

20%, Holm *et al.* compared the results of fluid administration strategy according to the Parkland formula *versus* a goal-directed therapy based on intrathoracic blood volume as an endpoint. Fluid volume administration was higher in the intrathoracic blood volume group compared to the Parkland group without a significant difference in CO and outcome.<sup>36</sup> In a randomized controlled trial by Aboelatta and Abdelsalam, fluid administration in the initial 72 h after thermal injury was significantly higher in the intrathoracic blood volume group.<sup>37</sup>

To summarize, it appears that fluid resuscitation targeting static preload parameters (intrathoracic blood volume, central venous pressure [CVP]) would lead to over-resuscitation without outcome benefit. The goal of hemodynamic monitoring is to ensure adequate oxygen delivery, ensuring a minimum CO in the absence of poor perfusion. An algorithm based on these observations is proposed in figure 3. Parameters of potential inappropriate organs perfusion (*e.g.*, low urine output, mean arterial pressure and CO, high lactate plasma concentration) should lead to evaluation of fluid responsiveness. The use of dynamic parameters should be preferred (absolute value of pulse pressure variation and stroke volume variation, intraindividual changes in pulse pressure variation, passive leg raising). Static preload parameters (*e.g.*, intrathoracic blood volume, CVP) should not be viewed as hemodynamic targets but as safety parameters (*e.g.*, CVP upper limit of 12 to 15 mmHg). In fluid unresponsive patients with cardiovascular or renal failure, other mechanisms of shock should be considered and lead to assessment of cardiac function.

## Types of Fluids

### Crystalloids Solutions

Ringer's Lactate (Baxter Healthcare, USA) is the most used crystalloid in burns. Emerging evidence suggests that the administration of chloride-rich solutions (*e.g.*, sodium chloride 0.9%) in critically ill patients is associated with a higher incidence of hyperchloremic acidosis and acute kidney injury.<sup>38</sup> Given the large volumes employed during burn shock resuscitation, the use of balanced solutions as first-line solutions is preferred in severe burn patients. A randomized controlled trial comparing two balanced solutions (Plasmalyte [Baxter Healthcare, Australia] *vs.* Ringer's Lactate) in major burns is ongoing (ClinicalTrials.gov identifier: NCT03118362).

### Colloids Solutions

The use of colloids in the first 24 h of burn resuscitation is controversial since the capillary leak may cause a transcapillary passage of large molecules into the interstitial space. Nevertheless, the onset of burn-related endothelial dysfunction and capillary leak was shown to be within 2 h after thermal injury with a median duration of 5 h.<sup>39</sup> Human albumin has multiple physiologic effects, including regulation of colloid osmotic pressure, antioxidant properties, nitric oxide

modulation and buffer capabilities, plasmatic binding, and transportation of various substances, which may be of particular relevance in severe burn injury.<sup>40</sup>

In a recent meta-analysis, use of albumin in the first 24 h was not associated with increased survival in severely burned patients. Nonetheless, significant statistical heterogeneity was present, and after exclusion of two studies at high risk of bias, albumin infusion was associated with reduced mortality (odds ratio [95% CI], 0.34 [0.19 to 0.58];  $P < 0.001$ ) and with decreased occurrence of compartment syndrome (pooled odds ratio [95% CI], 0.19 [0.07 to 0.50];  $P < 0.001$ ).<sup>41</sup> In a multicenter unblinded controlled trial in burn patients with total body surface area greater than 20%, Cooper *et al.* observed no significant difference between the treatment (5% albumin) and control groups (Ringer's Lactate) in organ failures.<sup>42</sup> In a before-after study, Park *et al.* observed that the use of 5% albumin in the first 24 h (*vs.* Ringer's Lactate and a synthetic colloid) was associated with lower mortality, use of vasopressors, duration of mechanical ventilation, and ventilator-associated pneumonia incidence.<sup>43</sup> Lawrence *et al.* performed a retrospective observational study where they observed that 5% albumin administration was associated with a reduction in crystalloid requirements.<sup>44</sup>

The current available evidence suggests that exogenous albumin administration in the first 24 h of burn shock resuscitation might be associated with better outcomes.<sup>40</sup> However, the optimal timing, dose, concentration, targeted albumin concentration, and patient population for albumin use remain unclear. Adequately powered, multicenter randomized controlled trials should be undertaken in burn patients.

The European Medicines Agency (London, United Kingdom) contraindicated the use of hydroxyethyl starch in burn patients. This decision was mainly based on the results of studies and meta-analyses, carried out mainly in populations of nonburn patients, showing that the use of these solutions was associated with a higher incidence of mortality and acute kidney injury.<sup>45</sup>

## Catecholamine Use in Early Burn Shock

Data regarding vasopressor use in the early phase of burn shock are scarce. In our experience, vasopressors are usually necessary in the hyperkinetic and vasoplegic phase (sepsis-like phase). Norepinephrine can increase stressed blood volume, venous return, and CO and may therefore limit the amount of administered fluids. Indeed, in hemodynamic physiology, vascular volume has two components. One part simply fills vessels but does not stretch the walls, and is called unstressed volume. The other part stretches the elastic walls and accounts for venous pressure and is called stressed volume. Only stressed volume determines venous return. Sufficient intravascular volume without very profound hypovolemia, however, is needed to allow recruitment of stressed volume.

## Adjunctive Therapies

Major thermal injury causes an important inflammation and production of free oxygen radicals. Antioxidants have been proposed as adjunctive therapies to initial resuscitation. In a randomized controlled trial, adjuvant administration of high-dose ascorbic acid ( $66 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ) during the first 24 h after burn injury significantly reduces fluid volume requirements (5.5 vs. 3.0 ml/kg per percent total body surface area,  $P < 0.01$ ).<sup>46</sup>

Furthermore, low-dose hydrocortisone could reduce shock duration in vasopressor-dependent severely burned patients, possibly by a reduction in capillary leakage and a correction of adrenal insufficiency. In a randomized controlled trial, median norepinephrine treatment duration was shorter in the corticosteroid-treated versus the placebo group (57 vs. 120 h,  $P = 0.035$ ) without a difference in mortality.<sup>47</sup> Nevertheless, the impact of hydrocortisone administration remains controversial with respect to the immune system and the risk of secondary infection in burn patients.

## Conclusions

Initial hemodynamic resuscitation of critically ill burn patients is a challenge with the dual purpose of avoiding under- and over-resuscitation. Balanced crystalloids solutions together with albumin represent the cornerstone of the resuscitation strategy, which we believe should be guided by hemodynamic monitoring. There are, however, many knowledge gaps regarding the hemodynamic targets, role of colloids, adjuvant therapies, vasopressors, and their impact on outcome and the immune system that should be addressed in the near future (see table with research agenda, Supplemental Digital Content, <http://links.lww.com/ALN/B737>).

## Acknowledgments

Many thanks to Dr. James Anstey, M.D., (Intensive Care Unit, Royal Melbourne Hospital, Parkville, Melbourne, Australia) for helping with the English revision of the manuscript.

## Research Support

Support was provided solely from institutional and/or departmental sources.

## Competing Interests

Dr. Legrand discloses consulting income from Sphingotec (Hennigsdorf, Germany), Novartis (Basel, Switzerland), and Baxter Healthcare (Deerfield, Illinois). The other authors declare no competing interests.

## Correspondence

Address correspondence to Dr. Legrand: Department of Anesthesiology and Critical Care and Burn Unit, Saint Louis Hospital, Assistance Publique-Hopitaux de Paris, Paris, France. [matthieu.legrand@aphp.fr](mailto:matthieu.legrand@aphp.fr). Information on purchasing reprints may be found at [www.anesthesiology.org](http://www.anesthesiology.org) or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

## References

1. Asch MJ, Feldman RJ, Walker HL, Foley FD, Popp RL, Mason AD Jr, Pruitt BA Jr: Systemic and pulmonary hemodynamic changes accompanying thermal injury. *Ann Surg* 1973; 178:218–21
2. Arlati S, Storti E, Pradella V, Bucci L, Vitolo A, Pulici M: Decreased fluid volume to reduce organ damage: A new approach to burn shock resuscitation? A preliminary study. *Resuscitation* 2007; 72:371–8
3. Mason SA, Nathens AB, Finnerty CC, Gamelli RL, Gibran NS, Arnoldo BD, Tompkins RG, Herndon DN, Jeschke MG: Inflammation and the Host Response to Injury Collaborative Research Program: Hold the pendulum: Rates of acute kidney injury are increased in patients who receive resuscitation volumes less than predicted by the Parkland equation. *Ann Surg* 2016; 264:1142–7
4. Klein MB, Hayden D, Elson C, Nathens AB, Gamelli RL, Gibran NS, Herndon DN, Arnoldo B, Silver G, Schoenfeld D, Tompkins RG: The association between fluid administration and outcome following major burn: A multicenter study. *Ann Surg* 2007; 245:622–8
5. Demling RH: The burn edema process: Current concepts. *J Burn Care Rehabil* 2005; 26:207–27
6. Kottke MA, Walters TJ: Where's the leak in vascular barriers? A review. *Shock* 2016; 46(3 suppl 1):20–36
7. Woodcock TE, Woodcock TM: Revised Starling equation and the glycocalyx model of transvascular fluid exchange: An improved paradigm for prescribing intravenous fluid therapy. *Br J Anaesth* 2012; 108:384–94
8. Vlachou E, Gosling P, Moiemmen NS: Microalbuminuria: A marker of systemic endothelial dysfunction during burn excision. *Burns* 2008; 34:241–6
9. Ademברי C, Sgambati E, Vitali L, Selmi V, Margheri M, Tani A, Bonaccini L, Nosi D, Caldini AL, Formigli L, De Gaudio AR: Sepsis induces albuminuria and alterations in the glomerular filtration barrier: A morphofunctional study in the rat. *Crit Care* 2011; 15:R277
10. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR: A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann Surg* 2011; 254:194–200
11. Osuka A, Kusuki H, Yoneda K, Matsuura H, Matsumoto H, Ogura H, Ueyama M: Glycocalyx shedding is enhanced by age and correlates with increased fluid requirement in patients with major burns. *Shock* 2017 Oct 11 [Epub ahead of print]
12. Baxter CR, Shires T: Physiological response to crystalloid resuscitation of severe burns. *Ann N Y Acad Sci* 1968; 150:874–94
13. Lorente JA, Ezpeleta A, Esteban A, Gordo F, de la Cal MA, Díaz C, Arévalo JM, Tejedor C, Pascual T: Systemic hemodynamics, gastric intramucosal PCO<sub>2</sub> changes, and outcome in critically ill burn patients. *Crit Care Med* 2000; 28:1728–35
14. Abu-Sittah GS, Sarhane KA, Dibo SA, Ibrahim A: Cardiovascular dysfunction in burns: Review of the literature. *Ann Burns Fire Disasters* 2012; 25:26–37
15. Maybauer MO, Asmussen S, Platts DG, Fraser JF, Sanfilippo F, Maybauer DM: Transesophageal echocardiography in the management of burn patients. *Burns* 2014; 40:630–5
16. Venkatesh B, Meacher R, Muller MJ, Morgan TJ, Fraser J: Monitoring tissue oxygenation during resuscitation of major burns. *J Trauma* 2001; 50:485–94
17. Samuelsson A, Steinvall I, Sjöberg F: Microdialysis shows metabolic effects in skin during fluid resuscitation in burn-injured patients. *Crit Care* 2006; 10:R172
18. Soussi S, Legrand M: Hemodynamic coherence in patients with burns. *Best Pract Res Clin Anaesthesiol* 2016; 30:437–43

19. Dépret F, Dunyach C, De Tymowski C, Chaussard M, Bataille A, Ferry A, Moreno N, Cupaciu A, Soussi S, Benyamina M, Mebazaa A, Serror K, Chaouat M, Garnier JP, Pirracchio R, Legrand M; PRONOBURN Group: Undetectable haptoglobin is associated with major adverse kidney events in critically ill burn patients. *Crit Care* 2017; 21:245
20. Legrand M, Michel T, Daudon M, Benyamina M, Ferry A, Soussi S, Maurel V, Chaussard M, Chaouat M, Mimoun M, Verine J, Mallet V, Mebazaa A; PRONOBURN Study Group: Risk of oxalate nephropathy with the use of cyanide antidote hydroxocobalamin in critically ill burn patients. *Intensive Care Med* 2016; 42:1080–1
21. Soussi S, Taccori M, De Tymowski C, Depret F, Chaussard M, Fratani A, Jully M, Cupaciu A, Ferry A, Benyamina M, Serror K, Boccara D, Chaouat M, Mimoun M, Cattan P, Zagdanski AM, Anstey J, Mebazaa A, Legrand M; PRONOBURN Group: Risk factors for acute mesenteric ischemia in critically ill burns patients—A matched case-control study. *Shock* 2018 Mar 20 [Epub ahead of print]
22. Markell KW, Renz EM, White CE, Albrecht ME, Blackbourne LH, Park MS, Barillo DA, Chung KK, Kozar RA, Minei JP, Cohn SM, Herndon DN, Cancio LC, Holcomb JB, Wolf SE: Abdominal complications after severe burns. *J Am Coll Surg* 2009; 208:940–7; discussion 947–9
23. Blot S, Hoste E, Colardyn F: Acute respiratory failure that complicates the resuscitation of pediatric patients with scald injuries. *J Burn Care Rehabil* 2000; 21:289–90
24. Kumar AB, Andrews W, Shi Y, Shotwell MS, Dennis S, Wanderer J, Summitt B: Fluid resuscitation mediates the association between inhalational burn injury and acute kidney injury in the major burn population. *J Crit Care* 2017; 38:62–7
25. Chappell D, Bruegger D, Potzel J, Jacob M, Brettner F, Vogeser M, Conzen P, Becker BF, Rehm M: Hypervolemia increases release of atrial natriuretic peptide and shedding of the endothelial glycocalyx. *Crit Care* 2014; 18:538
26. Soussi S, Deniau B, Ferry A, Levé C, Benyamina M, Maurel V, Chaussard M, Le Cam B, Blet A, Mimoun M, Lambert J, Chaouat M, Mebazaa A, Legrand M; PRONOBURN Group: Low cardiac index and stroke volume on admission are associated with poor outcome in critically ill burn patients: A retrospective cohort study. *Ann Intensive Care* 2016; 6:87
27. Dries DJ, Waxman K: Adequate resuscitation of burn patients may not be measured by urine output and vital signs. *Crit Care Med* 1991; 19:327–9
28. Fleming SJ, Dallemagne CR, Endre ZH, Yesberg NE, Cross RB: Acute lowering of plasma oncotic pressure increases filtration fraction and sodium excretion in conscious sheep. *Ren Physiol Biochem* 1992; 15:334–40
29. Sánchez M, García-de-Lorenzo A, Herrero E, Lopez T, Galvan B, Asensio M, Cachafeiro L, Casado C: A protocol for resuscitation of severe burn patients guided by transpulmonary thermodilution and lactate levels: A 3-year prospective cohort study. *Crit Care* 2013; 17:R176
30. Legrand M, Le Cam B, Perbet S, Roger C, Darmon M, Guerci P, Ferry A, Maurel V, Soussi S, Constantin JM, Gayat E, Lefrant JY, Leone M; support of the AZUREA Network: Urine sodium concentration to predict fluid responsiveness in oliguric ICU patients: A prospective multicenter observational study. *Crit Care* 2016; 20:165
31. Paratz JD, Stockton K, Paratz ED, Blot S, Muller M, Lipman J, Boots RJ: Burn resuscitation—hourly urine output versus alternative endpoints: A systematic review. *Shock* 2014; 42:295–306
32. Chen ZH, Jin CD, Chen S, Chen XS, Wang ZE, Liu W, Lin JC: The application of early goal directed therapy in patients during burn shock stage. *Int J Burns Trauma* 2017; 7:27–33
33. Anđel D, Kamolz LP, Roka J, Schramm W, Zimpfer M, Frey M, Anđel H: Base deficit and lactate: Early predictors of morbidity and mortality in patients with burns. *Burns* 2007; 33:973–8
34. Csontos C, Foldi V, Fischer T, Bogar L: Arterial thermodilution in burn patients suggests a more rapid fluid administration during early resuscitation. *Acta Anaesthesiol Scand* 2008; 52:742–9
35. Tokarik M, Sjöberg F, Balik M, Pafcuga I, Broz L: Fluid therapy LiDCO controlled trial—optimization of volume resuscitation of extensively burned patients through noninvasive continuous real-time hemodynamic monitoring LiDCO. *J Burn Care Res* 2013; 34:537–42
36. Holm C, Mayr M, Tegeler J, Hörbrand F, Henckel von Donnersmarck G, Mühlbauer W, Pfeiffer UJ: A clinical randomized study on the effects of invasive monitoring on burn shock resuscitation. *Burns* 2004; 30:798–807
37. Aboelatta Y, Abdelsalam A: Volume overload of fluid resuscitation in acutely burned patients using transpulmonary thermodilution technique. *J Burn Care Res* 2013; 34:349–54
38. Soussi S, Ferry A, Chaussard M, Legrand M: Chloride toxicity in critically ill patients: What's the evidence? *Anaesth Crit Care Pain Med* 2017; 36:125–30
39. Vlachou E, Gosling P, Moiemmen NS: Microalbuminuria: A marker of endothelial dysfunction in thermal injury. *Burns* 2006; 32:1009–16
40. Vincent JL, De Backer D, Wiedermann CJ: Fluid management in sepsis: The potential beneficial effects of albumin. *J Crit Care* 2016; 35:161–7
41. Eljaiek R, Heylbroeck C, Dubois MJ: Albumin administration for fluid resuscitation in burn patients: A systematic review and meta-analysis. *Burns* 2017; 43:17–24
42. Cooper AB, Cohn SM, Zhang HS, Hanna K, Stewart TE, Slutsky AS; ALBUR Investigators: Five percent albumin for adult burn shock resuscitation: Lack of effect on daily multiple organ dysfunction score. *Transfusion* 2006; 46:80–9
43. Park SH, Hemmila MR, Wahl WL: Early albumin use improves mortality in difficult to resuscitate burn patients. *J Trauma Acute Care Surg* 2012; 73:1294–7
44. Lawrence A, Faraklas I, Watkins H, Allen A, Cochran A, Morris S, Saffle J: Colloid administration normalizes resuscitation ratio and ameliorates “fluid creep.” *J Burn Care Res* 2010; 31:40–7
45. Zarychanski R, Abou-Setta AM, Turgeon AF, Houston BL, McIntyre L, Marshall JC, Fergusson DA: Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation: A systematic review and meta-analysis. *JAMA* 2013; 309:678–88
46. Tanaka H, Matsuda T, Miyagantani Y, Yukioka T, Matsuda H, Shimazaki S: Reduction of resuscitation fluid volumes in severely burned patients using ascorbic acid administration: A randomized, prospective study. *Arch Surg* 2000; 135:326–31
47. Venet F, Plassais J, Textoris J, Cazalis MA, Pachot A, Bertin-Maghit M, Magnin C, Rimmelé T, Monneret G, Tissot S: Low-dose hydrocortisone reduces norepinephrine duration in severe burn patients: A randomized clinical trial. *Crit Care* 2015; 19:21