EDITORIALS



General Anesthesia — Minding the Mind during Surgery

Gregory Crosby, M.D.

William Morton's demonstration of the use of ether in 1846 was powerful in part because the patient had no memory of the procedure; nowadays, patients expect to have amnesia with general anesthesia. But conscious awareness - the ability to remember and explicitly recall events that transpire during surgery — still occurs on occasion, sometimes with devastating psychological consequences. The easy explanation is that awareness is due to underdosing of the anesthetic agent. This explanation provides a sense of control and a ready fix (administer more anesthesia) but conveniently overlooks a secret: the state of consciousness is typically not monitored directly during general anesthesia. There simply is no accepted way to do it.

In this issue of the Journal, Avidan et al.¹ have addressed this in a large, prospective, multicenter study involving patients at high risk for intraoperative awareness owing to preexisting illness or the nature of the surgery. Subjects were randomly assigned to the management of anesthesia by means of the monitoring of endtidal anesthetic-agent concentration (ETAC) or with the use of an electroencephalogram (EEG)derived, commercially available depth-of-anesthesia monitor. The former, which is the standard method, measures exhaled volatile agent as a correlate of brain concentration; the latter reduces raw EEG data to a unitless bispectral index (BIS) that ranges from 100 (awake) to 0 (isoelectric EEG). Alerts were triggered when the ETAC or BIS values were outside the ranges that were considered to be adequate for surgical anesthesia (0.7 to 1.3 age-adjusted minimum alveolar concentration in the case of ETAC and 40 to 60 in the case of the BIS). Depth-of-anesthesia monitors are controversial, largely because they perform poorly under some conditions, and data on their ability to prevent awareness are conflicting.^{2,3} The results of the study by Avidan et al. add important new information to the debate. Despite arguably optimal management, definite or possible awareness occurred in 0.47% of the patients, and as compared with ETAC monitoring, the depth-of-anesthesia monitor conferred no benefit. Moreover, 41% of the cases of awareness occurred when the ETAC or BIS values were in the target ranges. Thus, although a combined ETAC and BIS protocol was not tested, intraoperative awareness was not entirely preventable with either monitoring method.

These findings are disappointing but not surprising. The key variables - consciousness, memory, and general anesthesia — are obscure, and tools to assess them intraoperatively are rudimentary. The nature of consciousness is a metaphysical problem that has challenged philosophers for centuries and neuroscientists for decades.⁴ Memory, a prerequisite for explicit recall, is similarly complicated,⁵ and agents that are used to produce general anesthesia have varied receptor profiles, actions on neural networks, and ways of producing amnesia.6,7 As such, general anesthesia is not a single phenotype, and there is little wonder that the neurobiology of the state is poorly understood. Yet amid this complexity, brain functioning is judged clinically much as it was 165 years ago, with the use of bodily signs and responses (e.g., blood pressure, heart rate, and movement) that are, at best, loosely related to higher brain function. The astonishing thing is not that awareness occurs but that it occurs so infrequently.

Having found no benefit from processed EEG monitoring, the investigators imply that an ETAC-

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

based protocol, complete with alarms and a checklist, is the way forward.¹ This inference is premature. Without an unmonitored control group, there is no proof that ETAC is superior to doing nothing special. Simply drawing attention to the possibility of awareness, as was done in both groups with a sign on the anesthesia machine, might be equivalent. Nor is ETAC feasible when only intravenous medications are used for anesthesia, which is now common practice. ETAC targets may also be unwise. Even low doses of volatile agents can produce hypotension, potentially exposing the sickest patients to unnecessary cardiovascular and cerebrovascular risk. In addition, ETAC puts the focus on the dose that is administered, not on the response of the brain. Applying similar logic to the cardiovascular system would mean administering predetermined doses of anesthetic agents without measuring blood pressure. Finally, basing ETAC ranges on age disregards the fact that chronologic age is a poor proxy for cognitive function, particularly among seniors, who constitute a large percentage of surgical patients and often have subclinical cerebral pathologic conditions or preexisting cognitive impairment.8 Thus, an ETAC protocol may inadvertently result in overdosing of the brain in cognitively vulnerable persons. This is worrisome because deep sedation is associated with a higher incidence of postoperative delirium, other adverse cognitive outcomes, and increased mortality in elderly surgical and critically ill patients.9.10 Association does not prove causation, but in some cases, too much anesthesia or sedation may be as undesirable as too little.

Monitors are meant to supplement, not supplant, clinical decision making, and depth-of-anesthesia monitors that reduce complex neurobiology to simple numbers are no exception. It is unreasonable to expect any such monitor to unfailingly detect conscious awareness — a specific and still mysterious property of the brain and mind — and neither patients nor physicians should think otherwise. Notwithstanding this and other weaknesses of current devices, a window into the anesthetized brain, albeit a foggy one, may still be useful, in conjunction with information from other monitors, in operating rooms, endoscopy suites, and critical care units as a generic, all-purpose index of the brain's response to powerfully sedating drugs. Whether these devices add value in this way remains to be seen, but when minding the mind during sedation and general anesthesia, a little insight into how the brain is reacting is apt to be better than none, especially if it challenges historical ways of gauging anesthetic depth and catalyzes the search for something better.

Disclosure forms provided by the author are available with the full text of this article at NEJM.org.

From the Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Boston.

1. Avidan MS, Jacobsohn E, Glick D, et al. Prevention of intraoperative awareness in a high-risk surgical population. N Engl J Med 2011;365:591-600.

 Myles PS, Leslie K, McNeil J, Forbes A, Chan MTV. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. Lancet 2004;363:1757-63.
 Avidan MS, Zhang L, Burnside BA, et al. Anesthesia aware-

ness and the bispectral index. N Engl J Med 2008;358:1097-108.4. Dehaene S, Changeux J-P. Experimental and theoretical ap-

proaches to conscious processing. Neuron 2011;70:200-27.5. Kandel ER. The biology of memory: a forty-year perspective. J Neurosci 2009;29:12748-56.

6. Alkire MT, Hudetz AG, Tononi G. Consciousness and anesthesia. Science 2008;322:876-80.

7. Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. N Engl J Med 2010;363:2638-50.

8. Evered LA, Silbert BS, Scott DA, Maruff P, Ames D, Choong PF. Preexisting cognitive impairment and mild cognitive impairment in subjects presenting for total hip joint replacement. Anesthesiology 2011;114:1297-304.

 Sieber FE, Zakriya KJ, Gottschalk A, et al. Sedation depth during spinal anesthesia and the development of postoperative delirium in elderly patients undergoing hip fracture repair. Mayo Clin Proc 2010;85:18-26. [Erratum, Mayo Clin Proc 2010;85:400.]
 Kalkman CJ, Peelen LM, Moons KG. Pick up the pieces: depth of anesthesia and long-term mortality. Anesthesiology 2011:114:485-7.

Copyright © 2011 Massachusetts Medical Society

Hamartoma Syndromes, Exome Sequencing, and a Protean Puzzle

John M. Opitz, M.D., and Lynn B. Jorde, Ph.D.

Gross malformations have been well known for hamartoma. In 1934, the concept entered the centuries, even millennia. However, it was not field of developmental pathology generally in until 1904 that Albrecht coined the concept of reference to tissue malformations. Hamartomata

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

The NEW ENGLAND JOURNAL of MEDICINE

ESTABLISHED IN 1812

AUGUST 18, 2011

VOL. 365 NO. 7

Prevention of Intraoperative Awareness in a High-Risk Surgical Population

Michael S. Avidan, M.B., B.Ch., Eric Jacobsohn, M.B., Ch.B., David Glick, M.D., M.B.A., Beth A. Burnside, B.A., Lini Zhang, M.D., Alex Villafranca, M.S., Leah Karl, B.A., Saima Kamal, M.D., Brian Torres, B.S.N., Michael O'Connor, M.D., Alex S. Evers, M.D., Stephen Gradwohl, B.S., Nan Lin, Ph.D., Ben J. Palanca, M.D., Ph.D., and George A. Mashour, M.D., Ph.D., for the BAG-RECALL Research Group.*

ABSTRACT

BACKGROUND

Unintended intraoperative <u>awareness</u>, which occurs when general anesthesia is not achieved or maintained, affects <u>up to 1%</u> of patients at <u>high risk</u> for this complication. We tested the hypothesis that a protocol incorporating the electroencephalogramderived bispectral index (BIS) is superior to a protocol incorporating standard monitoring of end-tidal anesthetic-agent concentration (ETAC) for the prevention of awareness.

METHODS

We conducted a prospective, randomized, evaluator-blinded trial at three medical centers. We randomly assigned 6041 patients at high risk for awareness to BIS-guided anesthesia (with an audible alert if the BIS value was <40 or >60, on a scale of 0 to 100, with 0 indicating the suppression of detectable brain electrical activity and 100 indicating the awake state) or ETAC-guided anesthesia (with an audible alert if the ETAC was <0.7 or >1.3 minimum alveolar concentration). In addition to audible alerts, the protocols included structured education and checklists. Superiority of the BIS protocol was assessed with the use of a one-sided Fisher's exact test.

RESULTS

A total of 7 of 2861 patients (0.24%) in the BIS group, as compared with 2 of 2852 (0.07%) in the ETAC group, who were interviewed postoperatively had definite intraoperative awareness (a difference of 0.17 percentage points; 95% confidence interval [CI], -0.03 to 0.38; P=0.98). Thus, the superiority of the BIS protocol was not demonstrated. A total of 19 cases of definite or possible intraoperative awareness (0.66%) occurred in the BIS group, as compared with 8 (0.28%) in the ETAC group (a difference of 0.38 percentage points; 95% CI, 0.03 to 0.74; P=0.99), with the superiority of the BIS protocol again not demonstrated. There was no difference between the groups with respect to the amount of anesthesia administered or the rate of major postoperative adverse outcomes.

CONCLUSIONS

The superiority of the BIS protocol was not established; contrary to expectations, fewer patients in the ETAC group than in the BIS group experienced awareness. (Funded by the Foundation for Anesthesia Education and Research and others; BAG-RECALL ClinicalTrials.gov number, NCT00682825.)

From the Departments of Anesthesiology (M.S.A., B.A.B., L.Z., S.K., B.T., A.S.E., S.G., B.J.P.) and Mathematics (N.L.), Washington University School of Medicine, St. Louis; the Department of Anesthesiology, University of Manitoba, Winnipeg, Canada (E.J., A.V.); the Department of Anesthesiology, University of Chicago, Chicago (D.G., L.K., M.O.); and the Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI (G.A.M.). Address reprint requests to Dr. Avidan at the Department of Anesthesiology, Washington University School of Medicine, Campus Box 8054, 660 S. Euclid Ave., St. Louis, MO 63110, or at avidanm@anest .wustl.edu.

*Members of the BIS or Anesthetic Gas to Reduce Explicit Recall (BAG-RECALL) Research Group are listed in the Supplementary Appendix, available at NEJM.org.

N Engl J Med 2011;365:591-600. Copyright © 2011 Massachusetts Medical Society.

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

591

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

WINTENDED INTRAOPERATIVE AWAREness is defined as the experience and explicit recall of sensory perceptions during surgery,¹ which can lead to post-traumatic stress disorder in as many as 70% of those who experience it.² In patients at high risk for intraoperative awareness, the incidence of awareness may approach 1%.³⁻⁵ An estimated 20,000 to 40,000 patients experience awareness yearly in the United States alone.¹ Some cases of awareness might occur as a result of inadequate anesthetic dosing⁶ and therefore constitute potentially preventable medical errors.⁷

A potent inhaled anesthetic agent is incorporated in the majority of general anesthetics, and concentrations of exhaled anesthetic are routinely measured.⁸ The minimum alveolar concentration (MAC) is the concentration of anesthesia required to prevent 50% of subjects from moving in response to a noxious surgical stimulus.⁹ When the end-tidal anesthetic-agent concentration (ETAC) is approximately 0.33 MAC, 50% of subjects do not respond appropriately to oral commands,⁹ and maintaining the ETAC at greater than 0.7 MAC during surgery is thought to decrease the incidence of awareness.^{10,11}

Candidate depth-of-anesthesia monitors based on electroencephalographic data have been developed, partly with the aim of preventing awareness. The most widely used is the bispectral index (BIS) monitor (Covidien), which processes a single frontal electroencephalographic signal with the use of a proprietary algorithm to calculate a dimensionless number that is intended to indicate the patient's level of consciousness. BIS values range from 0, indicating the suppression of detectable brain electrical activity, to 100, indicating the awake state. A target range between 40 and 60 has been advocated both to prevent awareness and to reduce the dose of anesthetic agent that has to be administered.¹²

Simple, but rigorous, protocol-based interventions can decrease perioperative complications and death.¹³⁻¹⁵ The incorporation of information technology into structured protocols to provide support for clinical decision making could improve patient safety and prevent medical errors.¹⁶ The adoption of technologic advances should be driven first by a compelling demonstration of clinical benefit and second by evidence of cost-effectiveness.¹⁷⁻²⁰

The B-Aware trial, involving 2500 patients, showed that administration of anesthesia with the use of a BIS protocol, as compared with standard anesthetic practice, decreased the incidence of intraoperative awareness by 0.74 percentage points (95% confidence interval [CI], 0.14 to 1.40) among patients at high risk for this complication.⁵ The B-Unaware trial (ClinicalTrials.gov number, NCT00281489) showed a reduction of 0 percentage points (95% CI, -0.56 to 0.57) in the incidence of awareness with a BIS protocol, as compared with an ETAC protocol.11 The B-Unaware trial also showed that the BIS protocol was not associated with a reduction in the amount of anesthesia administered or in postoperative mortality.^{11,21,22} Both protocols in the B-Unaware trial resulted in an incidence of awareness in high-risk patients that was lower than the predicted incidence (0.2% with both protocols vs. 1.0% predicted). However, the B-Unaware trial had important limitations. Most important, with 1941 patients, the confidence interval for the reduction in awareness was wide and did not rule out a clinically significant benefit (0.56 percentage points) attributable to BIS monitoring. Furthermore, the trial was conducted at a single center. We therefore designed the threecenter, international BIS or Anesthetic Gas to Reduce Explicit Recall (BAG-RECALL) trial²³ to investigate whether a structured BIS protocol was superior to a structured ETAC protocol in decreasing the incidence of intraoperative awareness among patients at high risk for this complication.

METHODS

STUDY OVERSIGHT

We conducted the study at three medical centers. A detailed description of the experimental protocol for the BAG-RECALL study has been published previously.23 The human studies committees at Washington University in St. Louis, University of Chicago, and University of Manitoba approved the study. The guidelines of the Consolidated Standards of Reporting Trials were followed in the conduct of the study and the reporting of results.²⁴⁻²⁶ The study was designed, the data were collected and analyzed, and the manuscript was prepared exclusively by the study investigators, all of whom made the decision to submit the manuscript for publication and vouch for the accuracy and completeness of the data and the analyses, as well as the adherence of this report to the study protocol.

PATIENTS

Patients 18 years of age or older who were undergoing elective surgery were evaluated for eligibil-

The New England Journal of Medicine

ity. Patients were eligible if they were at high risk for intraoperative awareness and were undergoing general anesthesia with isoflurane, sevoflurane, or desflurane. The criteria for identifying patients at high risk for awareness were based on previous studies, reviews, and guidelines.^{5,6,11,27} Patients at high risk were defined as those with at least one risk factor (Table 1). Patients who had dementia, who were unable to provide written informed consent, or who had a history of stroke with residual neurologic deficits were excluded.

STUDY DESIGN

In this prospective study, 6100 prerandomization designations were generated electronically in blocks of 100, divided equally between the groups. Labels indicating BIS group or ETAC group were sealed in opaque, numbered envelopes. Eligible patients underwent randomization after providing written informed consent. The anesthesia practitioners were aware of the patients' group assignments, but the patients, the postoperative interviewers, the expert reviewers, and the statistician were not.

PROCEDURES

A BIS Quatro (Covidien) sensor was applied to the forehead of each patient. Patients in the ETAC group had monitors configured to conceal the BIS number, so the anesthesia practitioners were unaware of the BIS values. The practitioners in both groups could view the ETAC. Summaries of the BIS and ETAC protocols²⁴ were given to the practitioners to provide education and to increase adherence to the protocol.

In the BIS group, an audible alarm was set to indicate when the BIS value exceeded 60 or fell below 40; no ETAC alarms were set in the BIS group, nor was there any recommendation to maintain the ETAC within any specific range. In the ETAC group, an audible alarm was set to indicate when the ETAC fell below 0.7 or exceeded 1.3 age-adjusted MAC.^{9,28} In the event that alarm settings were unavailable for ETAC, alarms were set for inspired anesthetic agents. During cardiopulmonary bypass, the concentration of anesthetic agent was measured from the effluent of the cardiopulmonary-bypass machine.²⁹ A sign was affixed to the anesthesia machines reminding practitioners to check the BIS value or the ETAC and to consider whether the patient might be aware. It was not the intention of the BIS and ETAC protocols to prescribe or restrict the use of anesthetic agents.²³ For example, practitioners

could decrease anesthetic administration at their discretion if a patient's condition was hemodynamically unstable, which itself can increase the risk of awareness. The protocols were designed to increase vigilance and to provide warnings that patients might be aware. BIS values and ETACs were recorded at minimum intervals of 1 minute by means of an electronic recording of anesthesia data with the use of MetaVision software (iMDsoft), by direct electronic transfer of data to Microsoft Excel (Microsoft), or by direct electronic transfer of data with the use of TrendFace Solo software (ixellence). Manual records of anesthesia and digital photographs of monitor trends were used as alternative sources of data in the rare instances that the computer data or the electronic anesthesia records were incomplete.

Intraoperative awareness was assessed with the use of a modified Brice questionnaire.^{23,30} Patients were evaluated within 72 hours after surgery and at 30 days after extubation. Patients who, at either interview, reported memories of the period between "going to sleep" and "waking up" were contacted by a different evaluator, who asked additional structured questions.²³ Every patient who reported such memories was offered referral to a psychologist. After all the patients completed the study, three experts independently reviewed the responses to the questionnaire from patients who had reported memories and determined whether the reported event involved definite awareness, possible awareness, or no awareness. The experts assigned each event of definite or possible awareness to one of the categories of the Michigan Awareness Classification Instrument.³¹ The same committee also reviewed awareness events for the Michigan Awareness Control Study (NCT00689091).32 In the event of a divergence of opinion, a fourth expert reviewer, who reviews cases for the Anesthesia Awareness Registry of the American Society of Anesthesiologists, made the final determination.

STATISTICAL ANALYSIS

The primary outcome measure was the incidence of intraoperative awareness in the BIS and ETAC groups.²³ We based our projected estimates of the incidence of awareness among high-risk patients on the results of the B-Aware and the B-Unaware trials.^{5,11} The null hypothesis was that the BIS protocol is not superior to the ETAC protocol in preventing intraoperative awareness. The alternative hypothesis was that the BIS protocol is superior

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

Table 1. Baseline Characteristics of the Patients and Inclusion Criteria Met by Them.*						
Variable	BIS Group (N=2861)	ETAC Group (N=2852)	P Value†			
Age — yr	60±14.2	61±14.4	0.20			
Male sex — no. (%)	1621 (56.7)	1679 (58.9)	0.09			
Body-mass index‡	30±8.4	30±8.3	0.72			
Race — no. (%)∬			0.85			
White	2405 (84.1)	2388 (83.7)				
Black	357 (12.5)	369 (12.9)				
Other	99 (3.5)	95 (3.3)				
ASA physical status — no. (%)¶			0.08			
1	23 (0.8)	19 (0.7)				
2	468 (16.4)	407 (14.3)				
3	1416 (49.5)	1407 (49.3)				
4	954 (33.3)	1019 (35.7)				
Inclusion criteria met — no. of patients (%)						
Planned open-heart surgery	1004 (35.1)	1037 (36.4)	0.32			
Aortic <u>stenosis</u>	260 (9.1)	261 (9.2)	0.94			
Pulmonary hypertension	132 (4.6)	135 (4.7)	0.83			
Use of opiates	725 (25.3)	741 (26.0)	0.58			
Use of benzodiazepines	458 (16.0)	441 (15.5)	0.57			
Use of anticonvulsant drugs	231 (8.1)	192 (6.7)	0.05			
Daily alcohol consumption	464 (16.2)	423 (14.8)	0.15			
ASA status 4	954 (33.3)	1019 (35.7)	0.06			
End-stage lung disease	59 (2.1)	57 (2.0)	0.87			
History of intraoperative awareness	50 (1.7)	41 (1.4)	0.35			
History of or anticipated difficult intubation	234 (8.2)	235 (8.2)	0.93			
Cardiac ejection fraction <40%	235 (8.2)	276 (9.7)	0.05			
Marginal exercise tolerance	1297 (45.3)	1353 (47.4)	0.11			
Composite no. of inclusion criteria met			0.30			
Median	2	2				
Interquartile range	1–3	1–3				
Composite no. of preexisting medical conditions			0.04			
Median	2	2				
Interquartile range	1–3	1–3				

* Plus-minus values are means ±SD. BIS denotes bispectral index, and ETAC end-tidal anesthetic-agent concentration.

† P values were calculated with the use of Fisher's exact test or the chi-square test with Yates' correction for categorical variables. For differences between means, P values were calculated with the unpaired Student's t-test. For differences between medians, P values were calculated with the Mann-Whitney U test.

The body-mass index is the weight in kilograms divided by the square of the height in meters.

Race was determined on the basis of information in the medical records.

¶The American Society of Anesthesiologists (ASA) physical status classification system is used to assess a patient's physical status before surgery. A score of 1 indicates a normal, healthy patient; 2, a patient with mild systemic disease; 3, a patient with severe systemic disease; and 4, a patient with severe systemic disease that is a constant threat to life. A patient could have more than one inclusion criterion.

in preventing intraoperative awareness. We esti- reduction of 0.4 percentage points in the incidence mated that with 6000 patients, the study would of definite awareness with the BIS protocol, as have 87% power to detect a clinically significant compared with the ETAC protocol (from 0.5% in

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

the ETAC group to 0.1% in the BIS group), at a one-tailed alpha level of 0.05, with the use of Fisher's exact test. As a prespecified secondary analysis, we planned to use a one-sided Fisher's exact test to determine whether there was a lower incidence of definite or possible awareness in the BIS group than in the ETAC group.²³ As a post hoc secondary analysis, we used Fisher's exact test to determine whether there was a lower incidence in the BIS group of awareness that caused distress. A chi-square test, Fisher's exact test, unpaired Mann-Whitney U test, or unpaired Student's t-test was used, as appropriate, for other comparisons. A modified intention-to-treat analysis was performed, which included all patients who underwent randomization and who were assessed for intraoperative awareness. Other than the assessments of awareness events, all significance testing was two-sided, and P values of less than 0.05 were considered to indicate statistical significance. The statistical analysis was performed with the use of the R statistical software package, version 2.13.0 (R Foundation for Statistical Computing) and Analyze-it software, version 2.11 (Analyze-it Software).

RESULTS

PATIENTS

Of an estimated 49,000 patients who were screened, 6041 were enrolled over the course of a 25-month period, from May 2008 through May 2010. A total of 5809 patients were included in the trial, of whom 5713 (98.3%) completed at least one postoperative interview and were included in the primary outcome analysis (Fig. 1). A total of 5413 patients (93.2%) completed the postoperative interviews at both times — within 72 hours after surgery and at 30 days after extubation. All the patients were treated with the protocol to which they had been randomly assigned. In the case of 6 patients in the ETAC group, practitioners were aware of BIS values for part of the time during the surgery. Table 1 shows the baseline characteristics of the patients and their risk factors for awareness.

INTRAOPERATIVE AWARENESS

In total, 49 patients, including patients from all three enrollment sites, reported having memories of the period between "going to sleep" and "waking up" at the end of surgery. Experts determined that 9 patients had definite intraoperative awareness (incidence, 0.16%; 95% CI, 0.08 to 0.30), and 27 patients had definite or possible awareness (incidence, 0.47%; 95% CI, 0.32 to 0.68). There were fewer cases of awareness in the ETAC group than in the BIS group, which was contrary to the expected result (i.e., contrary to the alternative hypothesis of the trial). The grading of awareness events according to the Michigan Awareness Classification Instrument³¹ is shown in Table 2. The comparisons of the frequency of awareness events between the BIS group and the ETAC group are shown in Table 3. Additional information about awareness experiences is provided in the Supplementary Appendix, available with the full text of this article at NEJM.org.

Table 4 shows the characteristics of patients who had definite or possible awareness, as compared with the rest of the patients in the study. Patients who experienced awareness, as compared with patients who did not, met a median of one additional inclusion criterion and had a median of one additional preexisting medical condition. A total of 5 of the 9 patients who experienced definite awareness and 6 of the 18 patients who experienced possible awareness did not have either BIS values of greater than 60 or ETAC values of less than 0.7 age-adjusted MAC. Overall, during the maintenance of anesthesia, the BIS was less than 60 a median of 94.0% of the time (interguartile range, 93.6 to 100), and the ETAC was greater than 0.7 age-adjusted MAC a median of 84.8% of the time (interquartile range, 67.2 to 95.3).

There were no important differences between the groups in the doses of sedative, hypnotic, opioid analgesic, or neuromuscular-blocking drugs administered. Specifically, midazolam was administered to 80.8% of the patients in the BIS group and to 79.7% of the patients in the ETAC group. During maintenance of anesthesia, the median ETAC in the ETAC group and in the BIS group was 0.9 age-adjusted MAC (interquartile range, 0.8 to 1.0), and the median BIS in both groups was 41 (interquartile range, 38 to 45). The median ETAC values were similar at the three study sites (0.8 age-adjusted MAC in Chicago, 0.9 age-adjusted MAC in St. Louis, and 0.9 age-adjusted MAC in Winnipeg), as were the median BIS values (43 in Chicago, 41 in St. Louis, and 43 in Winnipeg). The 30-day mortality was 1.96% (57 of 2907 patients) in the BIS group and 2.21% (64 of 2902 patients) in the ETAC group (a difference of 0.24 percentage points; 95% CI, -0.50 to 0.99). In both groups, the median length of stay in the hospital was 7.0 days, and the median length of stay in the intensive care unit was 2.1 days.

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

595

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.



DISCUSSION

The results of the BAG-RECALL trial do not support the superiority of the BIS protocol over the ETAC protocol in preventing intraoperative awareness among patients at high risk for this complication. The overall incidence of awareness was lower than anticipated, suggesting that both protocols were likely to have had efficacy, but the finding of fewer cases in the ETAC group than in the BIS group was contrary to the expected result. The 95% confidence interval for the difference in definite awareness includes, at one extreme, a 0.03 percentagepoint benefit attributable to BIS monitoring, which represents a number needed to treat of 3333 highrisk patients to prevent one episode of awareness.

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

Table 2. Incidence of Definite and Possible Awareness Events.*										
Awareness	Michigan Awareness Classification									
	Class 1	Class 2	Class 3	Class 4	Class 5	Class 1D	Class 2D	Class 3D	Class 4D	Class 5D
	number of patients									
BIS group										
Definite	2	1	0	0	0	0	1	1	1	1
Possible	0	6	0	2	0	1	1	2	0	0
ETAC group										
Definite	1	0	0	0	0	0	0	1	0	0
Possible	5	0	0	1	0	0	0	0	0	0

* In the Michigan Awareness Classification Instrument, class 1 indicates isolated auditory perceptions; class 2, tactile perceptions (e.g., perception of surgical manipulation or endotracheal tube); class 3, pain; class 4, paralysis (e.g., a feeling that one cannot move, speak, or breathe); and class 5, paralysis and pain. D indicates that there is associated distress (e.g., reports of fear, anxiety, suffocation, sense of doom, or sense of impending death). BIS denotes bispectral index, and ETAC end-tidal anesthetic-agent concentration.

The BAG-RECALL trial addressed the major limitations of the B-Unaware trial.¹¹ In particular, the current study was substantially larger than the B-Unaware trial and was conducted at three international sites, and minor risk factors for awareness were not used as enrollment criteria. The findings of the B-Unaware trial¹¹ and the current trial regarding definite and possible awareness are congruent. Furthermore, similar to the findings in the B-Unaware trial, the results of the BAG-RECALL trial did not show that the ETAC protocol was associated with an increase in postoperative mortality^{21,22} or in the amount of anesthetic agent adminstered.¹¹

Several other key implications of the BAG-RECALL trial warrant discussion. ETAC alarms at less than 0.7 age-adjusted MAC should be considered to be an effective active comparator in future studies of the prevention of intraoperative awareness, and the ETAC protocol used in this trial should be evaluated in the general surgical population. With some awareness events apparently occurring with BIS values below 60, decreasing anesthetic concentration solely on the basis of a BIS value of less than 60 is not recommended. Also, the risk of awareness might be incrementally increased with additional risk factors and coexisting conditions. Finally, notwithstanding major advances in our understanding of consciousness and anesthesia,^{33,34} until we clarify fully the mechanisms and measurement of anesthetic-induced unconsciousness and amnesia, some patients are still likely to have this complication.

Despite its wide scope, the BAG-RECALL trial has a number of limitations. Most important, it was designed to study the BIS and ETAC protocols in high-risk patients undergoing general anesthesia that is based predominantly on a potent inhaled agent; its results should not be extrapolated beyond this population. Second, the ETAC protocol

Table 3. Between-Group Comparison of Awareness Experiences.*						
Outcome	BIS Group (N=2861)	ETAC Group (N=2852)	P Value†	Difference, BIS-ETAC		
	no. (%)			percentage points (95% CI)		
Definite awareness: primary outcome	7 <u>(0.24)</u>	2 <u>(0.07)</u>	0.98	0.17 (-0.03 to 0.38)		
Definite or possible awareness: pre- specified secondary outcome	19 (0.66)	8 (0.28)	0.99	0.38 (0.03 to 0.74)		
Distressing experience of awareness: post hoc secondary outcome	8 (0.28)	1 (0.04)	0.99	0.24 (0.04 to 0.45)		

* BIS denotes bispectral index, and ETAC end-tidal anesthetic-agent concentration.

† One-tailed P values were calculated to test the null hypothesis that BIS is <u>not</u> superior to ETAC and the <u>alternative</u> hypothesis that BIS is <u>superior</u> to ETAC. A <u>one-tailed</u> P value of less than 0.05 with the use of Fisher's exact test would suggest that the null hypothesis should be rejected and the alternative hypothesis accepted.

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

Table 4. Characteristics of Patients with and Patients without Intraoperative Awareness.*							
Characteristic	Definite or Possible Intraoperative Awareness (N = 27)	No Intraoperative Awareness (N = 5686)	P Value†				
Age — yr	61±13.2	60±14.3	0.89				
Male sex — no. (%)	15 (55.6)	3285 (57.8)	0.82				
Body-mass index <u>:</u>	30±10.1	30±8.4	0.73				
Race — no. (%)§			0.26				
White	20 (74.1)	4773 (83.9)					
Other	7 (25.9)	913 (16.1)					
Inclusion criteria met — no. of patients (%)¶							
Planned open-heart surgery	14 (51.9)	2027 (35.6)	0.08				
Aortic stenosis	4 (14.8)	517 (9.1)	0.30				
Pulmonary hypertension	3 (11.1)	264 (4.6)	0.13				
Use of opiates	8 (29.6)	1458 (25.6)	0.66				
Benzodiazepine use	5 (18.5)	894 (15.7)	0.60				
Anticonvulsant use	3 (11.1)	420 (7.4)	0.45				
Daily alcohol consumption	6 (22.2)	881 (15.5)	0.30				
ASA status 4	13 (48.1)	1960 (34.5)	0.14				
End-stage lung disease	1 (3.7)	115 (2.0)	0.43				
History of intraoperative awareness	3 (11.1)	88 (1.5)	0.02				
History of or anticipated difficult intubation	0	469 (8.2)	0.20				
Cardiac ejection fraction <40%	5 (18.5)	506 (8.9)	0.18				
Marginal exercise tolerance	14 (51.9)	2636 (46.4)	0.57				
Composite no. of inclusion criteria			0.002				
Median	3	2					
Interquartile range	2–4	1-3					
Composite no. of preexisting medical conditions			0.003				
Median	3	2					
Interquartile range	1-4	1-3					
Received midazolam — no. (%)	18 (66.7)	4566 (80.3)	0.08				
Bispectral index			0.70				
Median	40	41					
Interquartile range	37–45	38–45					
ETAC in MAC equivalents			0.88				
Median	0.87	0.90					
Interquartile range	0.84–1.02	0.80–0.99					

* Plus-minus values are means ±SD. ETAC denotes end-tidal anesthetic-gas concentration, and MAC minimum alveolar concentration.

† P values were calculated with the use of Fisher's exact test or the chi-square test with Yates' correction for categorical variables. For differences between means, P values were calculated with the unpaired Student's t-test; for differences between medians, P values were calculated with the Mann–Whitney U test.

the body-mass index is the weight in kilograms divided by the square of the height in meters.

 \S Race was determined on the basis of information in the medical records.

A patient could have more than one inclusion criterion.

The American Society of Anesthesiologists (ASA) physical status classification system is used to assess a patient's physical status before surgery. Scores range from 1 (indicating a normal, healthy patient) to 6 (indicating a brain-dead patient whose organs are being removed for donor purposes); a score of 4 indicates a patient with severe systemic disease that is a constant threat to life.

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

was evaluated against only one of many available electroencephalogram-derived monitors. Third, it is conceivable that a protocol that is based on both BIS and ETAC alerts would perform better than one based on either approach alone. Fourth, it is difficult to know whether practitioners might become desensitized to either protocol; many people find audible alerts distracting, especially with a high rate of false alarms.35 Fifth, when a rare outcome is assessed, missing data can have an important effect. Although a small number of patients were not interviewed postoperatively, as shown in Figure 1, in most cases, this was because they did not awaken and died before the initial interview. Finally, in the case of a rare event such as awareness, unidentified risk factors such as genetic resistance to anesthetic agents could have been unequally distributed between the BIS and ETAC groups despite randomization and could have confounded the results.

The BAG-RECALL trial did not demonstrate the superiority of a BIS protocol over a protocol incorporating standard ETAC monitoring for the prevention of intraoperative awareness. Implementation of an ETAC-based protocol would require a brief, structured educational program, measurement of the exhaled concentration of volatile anesthetic agent, routine setting of audible alarms for threshold ETAC values, and checklists to maintain vigilance by the practitioner. Similar to other structured protocols that have enjoyed widespread adoption,¹³⁻¹⁵ an ETAC protocol like the one used in this trial could be implemented for patients at high risk for awareness who are undergoing general anesthesia with a potent inhaled agent.

Supported by a grant (CFM-08/15/2007, to Dr. Avidan) awarded by the Foundation for Anesthesia Education and Research and the American Society of Anesthesiologists and by a grant from the Winnipeg Regional Health Authority and University of Manitoba Department of Anesthesia (to Dr. Jacobsohn). Departmental support was also provided by the Department of Anestthesiology at Washington University in St. Louis and the Department of Anesthesiology at University of Chicago.

No potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

We thank the many anesthesiologists, certified registered nurse anesthetists, anesthesia technicians, and surgeons at the participating institutions who helped with the conduct of this trial — in particular, Dr. Jeffrey Apfelbaum, Mr. Randy Branson, Ms. Penny Candalario, Ms. Laura Veremakis, and Dr. Troy Wildes for their contributions; and Dr. Christopher Kent of the University of Washington (Seattle), Dr. Kevin Tremper of the University of Michigan (Ann Arbor), and Dr. Charles Hantler of Washington University in St. Louis for serving as expert reviewers of awareness cases.

REFERENCES

1. The Joint Commission. Sentinel event alert: preventing, and managing the impact of, anesthesia awareness. Issue 32. October 6, 2004. (http://www.pmhd.org/ Medical-Staff/documents/SentinelAlert Issue32_Preventingandmanagingthe impactofanesthesiaawareness_Joi.pdf.)

 Leslie K, Chan MT, Myles PS, Forbes A, McCulloch TJ. Posttraumatic stress disorder in aware patients from the B-Aware trial. Anesth Analg 2010;110:823-8.

3. Phillips AA, McLean RF, Devitt JH, Harrington EM. Recall of intraoperative events after general anaesthesia and cardiopulmonary bypass. Can J Anaesth 1993; 40:922-6.

4. Ranta S, Jussila J, Hynynen M. Recall of awareness during cardiac anaesthesia: influence of feedback information to the anaesthesiologist. Acta Anaesthesiol Scand 1996;40:554-60.

5. Myles PS, Leslie K, McNeil J, Forbes A, Chan MT. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. Lancet 2004;363:1757-63.

6. Ghoneim MM, Block RI, Haffarnan M, Mathews MJ. Awareness during anesthesia: risk factors, causes and sequelae: a review of reported cases in the literature. Anesth Analg 2009;108:527-35.

7. Landrigan CP, Parry GJ, Bones CB, Hackbarth AD, Goldmann DA, Sharek PJ. Temporal trends in rates of patient harm resulting from medical care. N Engl J Med 2010;363:2124-34. [Erratum, N Engl J Med 2010;363:2573.]

8. Walder B, Lauber R, Zbinden AM. Accuracy and cross-sensitivity of 10 different anesthetic gas monitors. J Clin Monit 1993;9:364-73.

9. Eger EI II. Age, minimum alveolar anesthetic concentration, and minimum alveolar anesthetic concentration-awake. Anesth Analg 2001;93:947-53.

10. Gonsowski CT, Chortkoff BS, Eger EI II, Bennett HL, Weiskopf RB. Subanesthetic concentrations of desflurane and isoflurane suppress explicit and implicit learning. Anesth Analg 1995;80:568-72.

11. Avidan MS, Zhang L, Burnside BA, et al. Anesthesia awareness and the bispectral index. N Engl J Med 2008;358:1097-108.

12. Punjasawadwong Y, Boonjeungmonkol N, Phongchiewboon A. Bispectral index for improving anaesthetic delivery and postoperative recovery. Cochrane Database Syst Rev 2007:CD003843.

13. Pronovost P, Needham D, Berenholtz S, et al. An intervention to decrease catheter-related bloodstream infections in

the ICU. N Engl J Med 2006;355:2725-32. [Erratum, N Engl J Med 2007;356:2660.] **14.** Haynes AB, Weiser TG, Berry WR, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. N Engl J Med 2009;360:491-9.

15. de Vries EN, Prins HA, Crolla RM, et al. Effect of a comprehensive surgical safety system on patient outcomes. N Engl J Med 2010;363:1928-37.

16. Bates DW, Cohen M, Leape LL, Overhage JM, Shabot MM, Sheridan T. Reducing the frequency of errors in medicine using information technology. J Am Med Inform Assoc 2001;8:299-308.

17. American College of Physicians. Information on cost-effectiveness: an essential product of a national comparative effectiveness program. Ann Intern Med 2008;148:956-61.

 Weinstein MC, Skinner JA. Comparative effectiveness and health care spending — implications for reform. N Engl J Med 2010;362:460-5.

19. Emanuel EJ, Fuchs VR, Garber AM. Essential elements of a technology and outcomes assessment initiative. JAMA 2007;298:1323-5.

20. Orszag PR, Emanuel EJ. Health care reform and cost control. N Engl J Med 2010;363:601-3.

N ENGLJ MED 365;7 NEJM.ORG AUGUST 18, 2011

599

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.

21. Kertai MD, Pal N, Palanca BJ, et al. Association of perioperative risk factors and cumulative duration of low bispectral index with intermediate-term mortality after cardiac surgery in the B-Unaware trial. Anesthesiology 2010;112:1116-27.

22. Kertai MD, Palanca BJ, Pal N, et al. Bispectral index monitoring, duration of bispectral index below 45, patient risk factors, and intermediate-term mortality after noncardiac surgery in the B-Unaware trial. Anesthesiology 2011;114:545-56.

23. Avidan MS, Palanca BJ, Glick D, et al. Protocol for the BAG-RECALL clinical trial: a prospective, multi-center, randomized, controlled trial to determine whether a bispectral index-guided protocol is superior to an anesthesia gas-guided protocol in reducing intraoperative awareness with explicit recall in high risk surgical patients. BMC Anesthesiol 2009;9:8.

24. Boutron I, Moher D, Altman DG, Schulz KF, Ravaud P. Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: explanation and elaboration. Ann Intern Med 2008;148:295-309.

25. Schulz KF, Altman DG, Moher D. CON-SORT 2010 statement: updated guidelines for reporting parallel group randomized trials. Ann Intern Med 2010;152:726-32.

26. Moher D, Hopewell S, Schulz KF, et al. CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. BMJ 2010;340:c869.

27. American Society of Anesthesiologists Task Force on Intraoperative Awareness. Practice advisory for intraoperative awareness and brain function monitoring. Anesthesiology 2006;104:847-64.

28. Nickalls RW, Mapleson WW. Agerelated iso-MAC charts for isoflurane, sevoflurane and desflurane in man. Br J Anaesth 2003;91:170-4.

29. Nussmeier NA, Moskowitz GJ, Weiskopf RB, Cohen NH, Fisher DM, Eger EI II. In vitro anesthetic washin and washout via bubble oxygenators: influence of anesthetic solubility and rates of carrier gas inflow and pump blood flow. Anesth Analg 1988;67:982-7.

30. Brice DD, Hetherington RR, Utting JE. A simple study of awareness and dream-

ing during anaesthesia. Br J Anaesth 1970; 42:535-42.

31. Mashour GA, Esaki RK, Tremper KK, Glick DB, O'Connor M, Avidan MS. A novel classification instrument for intraoperative awareness events. Anesth Analg 2009;110:813-5.

32. Mashour GA, Tremper KK, Avidan MS. Protocol for the "Michigan Awareness Control Study": a prospective, randomized, controlled trial comparing electronic alerts based on bispectral index monitoring or minimum alveolar concentration for the prevention of intraoperative awareness. BMC Anesthesiol 2009;9:7.
33. Alkire MT, Hudetz AG, Tononi G. Consciousness and anesthesia. Science 2008;32:2:876-80.

34. Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. N Engl J Med 2010;363:2638-50.

35. Block FE Jr, Nuutinen L, Ballast B. Optimization of alarms: a study on alarm limits, alarm sounds, and false alarms, intended to reduce annoyance. J Clin Monit Comput 1999;15:75-83.

Copyright © 2011 Massachusetts Medical Society.

SPECIALTIES AND TOPICS AT NEJM.ORG

Specialty pages at the *Journal*'s Web site (**NEJM.org**) feature articles in cardiology, endocrinology, genetics, infectious disease, nephrology, pediatrics, and many other medical specialties. These pages, along with collections of articles on clinical and nonclinical topics, offer links to interactive and multimedia content and feature recently published articles as well as material from the NEJM archive (1812–1989).

The New England Journal of Medicine

Downloaded from nejm.org by JOHN VOGEL on August 18, 2011. For personal use only. No other uses without permission.