Do You Believe What You See or What You Hear?

Ultrasound versus Stethoscope for Perioperative Clinicians

Shiroh Isono, M.D., Ph.D., Warren S. Sandberg, M.D., Ph.D., Yandong Jiang, M.D., Ph.D.

ANY perioperative clinicians wear a stethoscope and frequently use it for auscultation to, among other things, confirm endotracheal intubation or to rule out esophageal or unintended bronchial intubation. As a matter of routine, auscultation has been used for decades immediately after intubation or whenever correct placement of the endotracheal tube is in doubt in the maintenance of endotracheal intubation. However, accumulating evidence indicates the superiority of other novel techniques to identify esophageal intubation^{1,2} when compared with the venerable practice of auscultation. Now, other forms of malposition, specifically endobronchial intubation, appear



"Perhaps the stethoscope is closer to a costume piece than ever before."

to be more readily and accurately detected *via* techniques besides auscultation. Perhaps the stethoscope is closer to a costume piece than ever before.

In this issue of ANESTHESIOLOGY, Ramsingh *et al.*³ demonstrate the superiority of point-of-care ultrasound over auscultation of breath sounds to differentiate tracheal and bronchial intubation. The study is a prospective, randomized, double-blinded, crossover trial, and was well designed and conducted. The authors found that point-of-care <u>ultra-</u> sound is a reliable technique to directly reveal tracheal intubation by detecting tracheal dilation and <u>detects bronchial</u> <u>intubation</u> by demonstrating <u>absent</u> contralateral pleural <u>lung sliding</u> on the <u>unintubated</u> side.

Ramsingh *et al.*³ reproduced the low sensitivity (66%) and specificity (59%) of auscultation to diagnose endobronchial intubation, similar to previous reports of auscultation to differentiate tracheal *versus* bronchial intubation.^{4–6} Even in a nonstressful environment, for instance in this study,³ experienced clinicians are, at best, able to detect no more than two thirds of definite, deliberate bronchial intubations. The detection rate for unplanned endobronchial intubation, especially under stressful conditions (*e.g.*, trauma) and/or by less experienced clinicians, is probably even lower.

Every technology and technique has its time. Some last longer than others, but many are replaced. Auscultation has been practiced for endotracheal tube positioning for decades, mainly due to lack of better techniques. We tabulated the advantages and disadvantages of various techniques of endotracheal tube position detection to put point-of-care ultrasound in context (table1). Point-of-care ultrasound has all the cost and speed advantages of auscultation, but vastly superior sensitivity and specificity. The cost of a point-of-care ultrasound device is reasonable and dropping fast. Having already displaced ste-

thoscopy for cardiac examination, ultrasound seems poised to take out another province of auscultation. The remaining two techniques, fiberoscopy and radiography, cannot compete due to cost, requirements for skilled (additional) operators or interpretation, and potential risk exposure.

Point-of-care ultrasound is superior to auscultation to detect bronchial intubation. However, the sensitivity (93%) and specificity (96%) of differentiating tracheal versus bronchial intubation with this technique are only close to (but do not reach) 100%. Perhaps the gold standard method, chest x-ray, is still needed if misplacement of endotracheal tube cannot be ruled out. However, Ramsingh et al.3 did not present the sensitivity and specificity that could be obtained by combining the two assessments: visualizing tracheal dilation and nonsymmetric lung pleural sliding. If tracheal dilation and nonsymmetric lung pleural sliding are mutually exclusive in adults, then observing tracheal dilation indicates that nonsymmetric lung pleural sliding should not be observed, and vice versa. Therefore, combining these two assessments would further increase the accuracy of differentiating tracheal versus bronchial intubation. Point-of-care ultrasound also demonstrates the distance between cricoid cartilage

Copyright © 2016, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2016; 124:00-00

Downloaded From: http://anesthesiology.pubs.asahq.org/pdfaccess.ashx?url=/data/Journals/JASA/0/ by John Vogel on 03/11/2016

Image: A. Johnson.

Corresponding article on page XXX.

Accepted for publication November 4, 2015. From the Department of Anesthesiology, Graduate School of Medicine, Chiba University, Chiba, Japan (S.I.); Departments of Anesthesiology, Surgery, and Biomedical Informatics (W.S.S.); and Department of Anesthesiology (Y.J.), Vanderbilt University Medical Center, Nashville, Tennessee.

Techniques	Sensitivity (%)	Specificity (%)	Harmful Exposure	Immediacy	Cost per Use	Comments
Auscultation	60–66 ³⁻⁶	59–93 ³⁻⁶	No	Present	No	Low sensitivity, low specificity
Fiberoscopy	91.77	98.6^{7}	Yes	Present	Over \$50 ⁸	Harmful and not cost-effective
Ultrasound	9 3 ³	963	No	Present	No	Highly accurate and cost-effective, no delay in diagnosis
Chest x-ray*	100	100	Yes	Absent	\$100 (the national median costs) ⁹	It is considered as the gold standard. ^{4,10} Highly accurate; x-ray is harmful; not cost-effective; delayed diagnosis

Editorial Views

(glottis) and the endotracheal tube cuff, so clinicians can both confirm the cuff in the trachea and also estimate the distance between the cuff and the carina. The tracheal length of adult humans, defined from glottis to carina, is 9.2 ± 1.5 cm (mean ± SD),¹⁰ and knowing the distance between glottis and the cuff helps clinicians assess the likelihood of bronchial intubation. Therefore, combining tracheal dilation, assessment of pleural sliding on both sides, and assessment of cuff location relative to the glottis (and hence, carina) should achieve sensitivity and specificity comparable to those obtained with chest x-ray recommended as a standard.^{4,11} This assertion is readily testable and should be done since the authors have the primary data.³

Like any new technique, using point-of-care ultrasound to detect the tube misplacement is not without challenge. First, the learning curve is unknown, as mentioned by Ramsingh et al., although their operators all had reasonably brief minimum practice periods. From other domains, it is encouraging to know that only 5 min of training allows trainees to detect esophageal intubation in cadavers using ultrasound, with a sensitivity of 97%.¹² Second, whether the degree of cuff inflation affects the detection of tracheal dilation observed by ultrasound has not been tested. Some clinicians inflate the cuff only until a relatively low-pressure leak around it has been occluded and no more. In theory, such a cuff conforms relatively closely to the trachea and may produce no dilation. Since there is no standard pressure or volume for endotracheal tube cuff inflation, a low-inflated cuff may be difficult to visualize. This remains to be explored before the sensitivity and specificity of tracheal dilation can be assumed in all hands. Third, the study was conducted on adults.³ It is unknown if the findings are reproducible in pediatric populations. For example, tracheal dilation requires an inflated cuff. Although uncuffed endotracheal tubes are falling out of fashion, they still have adherents. Certainly, an uncuffed tube does not create tracheal dilation. Fourth, ultrasound only indirectly detects bronchial intubation or tracheal intubation. It does not directly show the location of the cuff or tip of the tube. Lack of observable tracheal dilation could occur if the cuff is below the sternal notch or in the esophagus but not necessarily in a main stem bronchus. Therefore, it requires another method to detect bronchial or esophageal intubation by visualizing lung pleural sliding. To visualize the lung sliding, the sonographer has to change the mode of ultrasound. This requires additional effort.

In other published work, it has been proposed that ultrasound can be used dynamically in direct conjunction with the airway procedures for maximum benefit in airway management.¹³ This is because one can directly visualize the tube passing to the trachea or esophagus while it is being inserted. The previous investigators¹³ were concerned that the accuracy of detecting bronchial intubation with point-of-care ultrasound postintubation is not as high as that obtained by dynamically examining the airway during

Downloaded From: http://anesthesiology.pubs.asahq.org/pdfaccess.ashx?url=/data/Journals/JASA/0/ by John Vogel on 03/11/2016

intubation. In contrast, Ramsingh *et al.* found the accuracy of detecting bronchial intubation is high even when the assessment was conducted after the intubation was completed.

The studies of Ramsingh *et al.*³ and others^{12,14} open new areas to explore other potential methods with which the accuracy to detect misplacement of endotracheal tube could be further optimized as enumerated below. 1) To inflate the cuff with liquid instead of air. It was described previously that filling the cuff with fluid helps in revealing the cuff position by ultrasonagraphy.¹⁵ This method is proven valuable even in pediatric population.¹⁶ 2) To make the cuff with echogenic material. Such material enhances the image of the cuff and makes the cuff more visible relative to the tracheal region not interfaced with cuff. Such alterations could make the detection of cuff location much easier, and the sonographer can visualize the cuff and tracheal rings around the cuff, which confirms tracheal intubation, not esophageal or bronchial intubation. In other words, one-stop shopping.

The findings³ of Ramsingh *et al.* further undermine the perioperative role of the stethoscope (except perhaps as a fomite): accuracy using auscultation to detect intentional or unintentional bronchial intubation is low. Alternatively, point-of-care ultrasound produces the accuracy close to that obtained with chest x-ray, but using a device that is becoming prevalent and potentially ubiquitous in the perioperative environment. We expect that continuous improvement of image quality and clinical experience with point-of-care ultrasound will further prompt its application in detecting misplacement of endotracheal tubes and beyond and enable us to provide better patient care.

Competing Interests

The authors are not supported by, nor maintain any financial interest in, any commercial activity that may be associated with the topic of this article.

Correspondence

Address correspondence to Dr. Jiang: yandong.jiang@ vanderbilt.edu

References

1. Grmec S: Comparison of three different methods to confirm tracheal tube placement in emergency intubation. Intensive Care Med 2002; 28:701–4

- Knapp S, Kofler J, Stoiser B, Thalhammer F, Burgmann H, Posch M, Hofbauer R, Stanzel M, Frass M: The assessment of four different methods to verify tracheal tube placement in the critical care setting. Anesth Analg 1999; 88:766–70
- Ramsingh D, Frank E, Haughton R, Schilling J, Gimenez K, Banh E, Rinehart J, Cannesson M. Auscultation *versus* point of care ultrasound to determine tracheal versus bronchial intubation. A diagnostic accuracy study. ANESTHESIOLOGY 2015; xx:xx-x
- 4. Brunel W, Coleman DL, Schwartz DE, Peper E, Cohen NH: Assessment of routine chest roentgenograms and the physical examination to confirm endotracheal tube position. Chest 1989; 96:1043–5
- Sitzwohl C, Langheinrich A, Schober A, Krafft P, Sessler DI, Herkner H, Gonano C, Weinstabl C, Kettner SC: Endobronchial intubation detected by insertion depth of endotracheal tube, bilateral auscultation, or observation of chest movements: Randomised trial. BMJ 2010; 341:c5943
- Vezzani A, Manca T, Brusasco C, Santori G, Valentino M, Nicolini F, Molardi A, Gherli T, Corradi F: Diagnostic value of chest ultrasound after cardiac surgery: A comparison with chest X-ray and auscultation. J Cardiothorac Vasc Anesth 2014; 28:1527–32
- 7. Dietrich KA, Conrad SA, Romero MD, Lloyd JL, Fontenot EE: Accuracy of flexible fiberoptic endoscopy in identifying abnormal endotracheal tube positions. Pediatr Emerg Care 1990; 6:257–9
- Choure AJ, Manali ED, Krizmanich G, Gildea TR, Mehta AC. High price of bronchoscopy: Cost of maintenance and repair of flexible bronchoscopes. J Bronchol Intervent Pulmonol 2005; 12:147–50
- 9. Healthcare Price Transparency. Available at: https://www. nerdwallet.com/blog/health/2015/03/17/how-much-doesan-xray-cost/. Accessed March 17, 2015
- Pu J, Leader JK, Meng X, Whiting B, Wilson D, Sciurba FC, Reilly JJ, Bigbee WL, Siegfried J, Gur D: Three-dimensional airway tree architecture and pulmonary function. Acad Radiol 2012; 19:1395–401
- Rudraraju P, Eisen LA: Confirmation of endotracheal tube position: A narrative review. J Intensive Care Med 2009; 24:283–92
- 12. Ma G, Davis DP, Schmitt J, Vilke GM, Chan TC, Hayden SR: The sensitivity and specificity of transcricothyroid ultrasonography to confirm endotracheal tube placement in a cadaver model. J Emerg Med 2007; 32:405–7
- 13. Kristensen MS: Ultrasonography in the management of the airway. Acta Anaesthesiol Scand 2011; 55:1155–73
- Blaivas M, Tsung JW: Point-of-care sonographic detection of left endobronchial main stem intubation and obstruction *versus* endotracheal intubation. J Ultrasound Med 2008; 27:785–9
- 15. Hatfield A, Bodenham A: Ultrasound: An emerging role in anaesthesia and intensive care. Br J Anaesth 1999; 83:789–800
- Tessaro MO, Salant EP, Arroyo AC, Haines LE, Dickman E: Tracheal rapid ultrasound saline test (T.R.U.S.T.) for confirming correct endotracheal tube depth in children. Resuscitation 2015; 89:8–12