## Anesthesia, Microcirculation, and Wound Repair in Aging

Itay Bentov, M.D., Ph.D., May J. Reed, M.D.

#### ABSTRACT

Age-related changes in skin contribute to impaired wound healing after surgical procedures. Changes in skin with age include decline in thickness and composition, a decrease in the number of most cell types, and diminished microcirculation. The microcirculation provides tissue perfusion, fluid homeostasis, and delivery of oxygen and other nutrients. It also controls temperature and the inflammatory response. Surgical incisions cause further disruption of the microvasculature of aged skin. Perioperative management can be modified to minimize insults to aged tissues. Judicious use of fluids, maintenance of normal body temperature, pain control, and increased tissue oxygen tension are examples of adjustable variables that support the microcirculation. Anesthetic agents influence the microcirculation of a combination of effects on cardiac output, arterial pressure, and local microvascular changes. The authors examined the role of anesthetic management in optimizing the microcirculation and potentially improving postoperative wound repair in older persons. (ANESTHESIOLOGY 2014; 120:760-72)

URGICAL wound repair is a major problem in the older population,<sup>1</sup> who are at increased risk of wound dehiscence and infection.<sup>2</sup> As a specific example, surgical site infections (SSIs) are common (approximately 500,000 cases annually in the United States), lead to worse patient outcome (patients who develop SSI are twice as likely to die),<sup>3</sup> and are an enormous economic burden (1-10 billion dollars annually).<sup>4</sup> Many factors contribute to age-related changes in skin<sup>5</sup> and subsequent vulnerability to impaired wound healing and infection. Changes in skin with age (fig. 1) include a decline in epidermal and dermal thickness and composition, as well as a decrease in the number of most resident cell types. The dermal-epidermal junction is flattened and the microcirculation is diminished.<sup>6</sup> The latter is defined as blood flow through arterioles, capillaries, and venules and is the key system that affects the entire skin surface. In the aging patient, the microcirculation in the skin is <mark>reduced by 40%</mark> between the ages of 20 and 70 yr.<sup>7</sup> The microcirculation provides tissue perfusion, fluid hemostasis, and delivery of oxygen and other nutrients. It also controls temperature and the inflammatory response. Surgical incisions cause disruption of the microcirculation in the skin as manifested by local edema resulting from vasodilation and increased vascular permeability.

Perioperative management can be modified to optimize the microcirculation. Measures that support the microcirculation include careful use of fluids, normothermia, pain control, and smoking cessation.<sup>8</sup> Factors that can be influenced by intraoperative management (judicious use of fluids, maintenance of normal body temperature, pain control, and increased tissue oxygen tension) have been suggested to be beneficial as well. Most anesthetic agents also influence the microcirculation: a reduction in cardiac output and arterial pressure decreases flow in the microcirculation, whereas anesthetic-induced local microvascular changes and vasodilatation can increase perfusion.<sup>9</sup> Optimization of these variables plays an important role in enhancing the microcirculation in all patients, but is especially relevant if modifications could improve postoperative wound healing in the older population.

In this review, we will use skin as a representative organ to describe age-related changes that negatively affect the microcirculation and have subsequent impacts on wound healing and the incidence of postoperative infection. We will then examine the role of anesthesia management in minimizing detrimental effects on the microcirculation. A greater understanding of these variables could promote improvements that lead to better outcomes with respect to wound repair in older patients.

This article is featured in "This Month in Anesthesiology," page 1A.

Submitted for publication January 18, 2013. Accepted for publication September 24, 2013. From the Department of Anesthesiology and Pain Medicine (I.B.) and Division of Gerontology and Geriatric Medicine, Department of Medicine (M.J.R.), Harborview Medical Center, University of Washington, Seattle, Washington.

Copyright © 2014, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins. Anesthesiology 2014; 120:760-72



Fig. 1. Numerous changes in skin with age contribute to impaired wound healing.

В

#### Summary of Wound Repair and Aging

It has been nearly a century since it was noted that the rate of cutaneous scar formation after a wound is inversely related to the age of the patient.<sup>10</sup> Four decades ago, it was observed that older age was associated with an increased risk of postoperative disruption of the surgical wound, leading to higher mortality.<sup>11</sup> Recent data suggest that in patients older than 65 yr, development of SSI is associated with a two-fold increase in cost and a staggering four-fold increase in mortality.<sup>12</sup>

Wound healing ensues *via* a sequential chain of events (with variable overlap) that includes inflammation, tissue

#### Α Inflammation



Degranulation Phagocytosis Infiltration Cytokines (TGF-ß1, PDGF, TNFα, VEGF)

# tissue formation Fibroblast $\hat{\mathbf{G}}$

Proliferation and

Extracellular matrix formation Angiogenesis

Angiogenesis



#### Inflammation

Immediately after injury to the skin, local blood vessels constrict and circulating platelets attach to the endothelial

С



Extracellular matrix remodeling Wound contraction

Fig. 2. The stages of wound healing are a sequential chain of events that include: (A) inflammation, (B) proliferation and granulation tissue formation, and (C) extracellular matrix (ECM) deposition and tissue remodeling. PDGF = platelet-derived growth factor; TGF- $\beta$ 1 = transforming growth factor- $\beta$ 1; TNF- $\alpha$  = tumor necrosis factor- $\alpha$ ; VEGF = vascular endothelial growth factor.

wall to prevent further bleeding. The platelets aggregate and release their granules to form a fibrin clot. During this process, several mediators and cytokines that are also regulators of cell proliferation, extracellular matrix synthesis, and angiogenesis are released. As examples, transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) and platelet-derived growth factor elicit rapid chemotaxis of neutrophils, monocytes, and fibroblasts to the injured area, which stimulates generation of additional cytokines. The latter include the angiogenic factor vascular endothelial growth factor (VEGF) and the proinflammatory molecules tumor necrosis factor- $\alpha$  and interleukin-1 $\beta$ .<sup>16</sup>

Age-related changes in the inflammatory response (fig. 3A) result in alterations in cell adhesion, cell migration, and cytokine production. In an aged mouse model of impaired wound healing, reduced phagocytosis by macrophages and delayed T-cell infiltration into wounds were found in the aged mice relative to young mice. As expected, the production of most chemokines (as measured by messenger RNA levels) declined by 20–70% with age,<sup>17</sup> but levels of some proinflammatory cytokines increased. Subsequent studies highlighted the critical role of macrophages: when young mice were injected at a biopsy wound site with antimacrophage serum, they exhibited delayed closure of the wounds, similar to that of aged mice.<sup>18</sup> In contrast, aged mice wounds that were given peritoneal macrophages from young mice showed enhanced wound healing.<sup>19</sup> Interestingly, in old female mice, healing of full-thickness dermal wounds on the upper dorsum was accelerated by treadmill



Fig. 3. Age-related changes in every stage of the wound-healing process may delay and impair surgical wound repair in older patients. ECM = extracellular matrix; HA = hyaluronan; MMP = matrix metalloproteinases; SPARC = secreted protein acidic and rich in cysteine; TGF- $\beta$  = transforming growth factor- $\beta$ ; TIMP = tissue inhibitor of matrix metalloproteinases; VEGF = vascular endothelial growth factor.

exercise, potentially reflecting an exercise-induced antiinflammatory response in the wound.<sup>20</sup> Similarly, adult men (mean age, 61 yr) who exercised before an experimental wound showed a reduction in stress-related neuroendocrine responses that was accompanied by an accelerated wound healing.<sup>21</sup> Another therapeutic intervention is suggested by the finding that castration of male mice is associated with a reduced inflammatory response and results in acceleration of cutaneous wound healing.<sup>22</sup> This might reflect suppression of the androgen receptor, which enhances the inflammatory response through an increase in tumor necrosis factor- $\alpha$  expression.<sup>23</sup> The quantity and distribution of several growth factors are also different between aged and young mice. Expression of platelet-derived growth factor, epidermal growth factor, and their cognate receptors is delayed with increasing animal age.<sup>24</sup> Alterations of the inflammatory response are also found in aged humans. Although total leukocyte and neutrophil counts are slightly lower in samples from older individuals,<sup>25</sup> granulocyte adherence is greater in aged subjects especially in women.<sup>26</sup> Phagocytosis is decreased in neutrophils from old, compared with young, healthy donors, potentially secondary to reduced neutrophil CD16 expression in the aged.<sup>27</sup>

In summary, inflammation is an important part of the initial host response to injury and pathogens. Aging is sometimes associated with a persistent proinflammatory state; at the same time, there is a reduction in the ability to generate an acute inflammatory response during injury. This paradox can result in disrupted wound healing due to lack of synchronization between pro- and antiinflammatory responses.

#### **Proliferation and Tissue Formation**

Several hours after injury, reepithelization begins.<sup>28</sup> Wounded epidermal cells express integrin receptors, produce collagenase, and activate plasmin by plasminogen activator. These changes allow them to separate from neighboring cells, interact with and degrade extracellular matrix proteins, and enable movement from the dermis into the margins of the wound area. Epidermal cells in the wound margins begin to proliferate approximately 1 or 2 days after the injury, producing a scaffold of basement membrane proteins from the margins inward. During this process, mediators and cytokines (interleukins and  $\alpha$ - and  $\beta$ -chemokines) that regulate angiogenesis and influence the microcirculation are released.<sup>29</sup> Several days after the injury, macrophages, fibroblasts, and blood vessels simultaneously invade the wound.<sup>30</sup> Macrophages produce growth factors, such as TGF-B1 and platelet-derived growth factor. Fibroblasts synthesize a new matrix (first a provisional matrix of fibrin, collagen III, fibronectin, and hyaluronan; later a structural matrix of primarily collagen I replaces the provisional matrix). Blood vessels supply oxygen and nutrients, which is essential to sustain the newly formed granulation tissue. As an example, the deposition of collagen relies on proline hydroxlyase, an oxygendependent enzyme.<sup>31</sup>

Studies based on animal models demonstrate that proliferation of the cell types responsible for tissue formation is reduced with aging<sup>32</sup> (fig. 3B). As an example, punch biopsy samples obtained repeatedly over the life span of hamsters found that in vitro proliferative capacity of dermal fibroblasts mimicked in vivo dermal wound repair.33 In healthy human volunteers, superficial, split-thickness wound epithelization is delayed in older persons (older than 65 yr) when compared with the control group (18-55 yr old).<sup>34</sup> Most studies suggest that wound angiogenesis is also decreased by approximately 70% 1 week after injury in aged animals.<sup>35,36</sup> Others propose an altered, dysregulated response with some extracellular matrix components increased, some decreased, and many showing disrupted ultrastructure.<sup>37</sup> Impaired endothelial cell function and reduced VEGF expression are possible mechanisms of age-related deficits in angiogenesis, which has an adverse effect on the development of an effective microcirculation.<sup>38</sup> In an explant model, age-related deficiencies in angiogenesis were reversed, in part, by stimulation with angiogenic growth factors.<sup>39</sup>

#### Extracellular Matrix and Tissue Remodeling

During the last phase of wound healing, the extracellular matrix begins to remodel and the wound undergoes further contraction. Fibroblasts assume a myofibroblast phenotype characterized by bundles of a-smooth muscle actin-containing microfilaments. Synchronized collagen reorganization occurs by synthesis and catabolism (although at a much slower rate than in previous stages), which allows the granulation tissue to turn into a scar. Deposition and remodeling of collagen are slower in aged animals resulting in less scar formation.<sup>40</sup> Moreover, the collagen deposited has a looser, more disorganized matrix that has decreased tensile strength. The changes in aged collagen matrix reflect decreases in circulating factors, in particular reduced levels of TGF-B1-a potent stimulator of collagen synthesis.<sup>41</sup> Of note, dermal fibroblasts from aged and young donors exposed to TGF-B1 exhibit similar biosynthetic and contractile properties.<sup>42</sup>

Other matrix components<sup>43</sup> that are altered with age (fig. 3C) include: decreases in osteonectin (also known as secreted protein acidic and rich in cysteine), increases in thrombospondin,44 and alterations in fibronectin and laminin.45,46 Nonprotein components of the extracellular matrix include glycosaminoglycans, such as hyaluronan, which interact with other matrix components to maintain hydration in the dermis. Hyaluronan is a linear disaccharide polymer that can range from 2 to 25,000 disaccharides with molecular masses up to  $2 \times 10^4$  kDa. Hyaluronan size determines its biologic properties: high molecular-weight forms can inhibit proliferation and migration of many cell types, whereas middle and lower molecular-weight forms usually promote tissue formation.<sup>46</sup> Hyaluronan content is maintained in aged wound dermis, but its degradation is reduced.<sup>47</sup> Wound healing also requires matrix metalloproteinases (MMPs), which promote cell proliferation

and vessel ingrowth by degrading the existing extracellular matrix. MMP activity is balanced, in part, by endogenous tissue inhibitors of metalloproteinases. Aged tissues are associated with dysregulation of MMP activity,<sup>48</sup> with a tendency toward overexpression of MMPs<sup>49</sup> and reduced levels of tissue inhibitors of metalloproteinases.<sup>50</sup>

As highlighted above, alterations occur during each stage of wound repair in aging, and many of these changes negatively impact the microcirculation. Nonetheless, given sufficient time, aged animals eventually (age-related delay is approximately 30–40%) catch up to their young counterparts with respect to most aspects of tissue repair.<sup>51</sup>

### The Surgical Context of Wound Repair and Aging

Measures that support the microcirculation improve wound repair, thereby reducing the risk of postoperative dehiscence and infection.<sup>52</sup> General preoperative measures such as smoking cessation and optimal management of comorbid medical conditions have been reviewed in other contexts.<sup>53,54</sup> For the purpose of this review, we will focus on interventions in the perioperative setting.

#### **Oxygen Administration**

Wound healing is dependent upon adequate levels of oxygen.<sup>55</sup> Oxygen interacts with growth factor signaling and regulates numerous transduction pathways necessary for cell proliferation and migration.<sup>56</sup> It is also an indispensable factor for oxidative killing of microbes.<sup>57</sup> Consequently, the effects of oxygen tension on the outcome of surgical wounds have been best studied in the context of postoperative infection. Resistance to surgical wound infection is presumed to be oxygen dependent—with low oxygen tension viewed as a predictor of the development of infection,<sup>56</sup> particularly when subcutaneous tissue oxygenation (measured by a polarographic electrode) decreases to less than 40 mmHg.<sup>58</sup>

In two recent meta-analyses, one found that perioperative supplemental oxygen therapy exerts a significant beneficial effect on the prevention of SSIs,<sup>59</sup> whereas the other suggested a benefit only for specific subpopulations.<sup>60</sup> Although most authors suggest that supplemental oxygen during surgery is associated with a reduction in infection risk,<sup>61,62</sup> others propose it may be associated with an increased incidence of postoperative wound infection.<sup>63</sup> Notably, in the latter report, the sample size was small and there was a difference in the baseline characteristics of the groups. A prospective trial randomizing patients to either 30 or 80% supplemental oxygen during and 2 h after surgery did not find any difference in several outcome measures including death, pulmonary complications, and wound healing.<sup>64</sup> Of note, the administration of oxygen to aged subjects may be limited by the finding that although arterial oxygen tension did not decrease with age, there was reduced steady-state transfer of carbon monoxide in the lungs.<sup>65</sup> This indicates that oxygen transport could be diffusion-limited in older subjects, especially when oxygen consumption is increased. Furthermore, longitudinal studies of five healthy men over 3 decades showed impaired efficiency of maximal peripheral oxygen extraction,<sup>66</sup> suggesting that tissue oxygen uptake is reduced in the aged subjects.<sup>67</sup> This likely reflects a decrease in the number of capillaries as well as a reduction in mitochondrial enzyme activity.<sup>68</sup> Animal models (rabbit<sup>69</sup> and mouse<sup>69,70</sup>) have suggested that aging and ischemia have an additive effect on disruption of wound healing. Consequently, the potential benefit of increasing tissue oxygen tension during surgical wound repair in older patients should be further evaluated.

#### Fluid Management

Clinical signs of intravascular volume status are often difficult to evaluate in older persons.<sup>71</sup> Moreover, the repercussions of extremes of intravascular volume have harmful sequelae. As an example, hypovolemia decreases tissue oxygen concentrations,<sup>72</sup> whereas excessive fluid administration increases tissue edema, which can adversely affect healing.<sup>73</sup> Numerous types of fluids<sup>74</sup> and devices<sup>75</sup> have been evaluated as optimizers of volume status in the general surgical population, but lack of definition of liberal versus restrictive regimens precludes evidence-based guidelines.<sup>76</sup> When fluid administration was guided by subcutaneous oxygen tension rather than clinical criteria, patients received more fluids and accumulated more collagen in their surgical incisions.<sup>76</sup> However, in residents of nursing homes who are at a higher risk of impaired hydration (and subsequently reduced tissue oxygenation),<sup>77</sup> supplemental oral fluid intake neither reverses these deficits nor improves wound healing.<sup>78</sup> The need for more accurate determination of volume status is underscored by studies which showed that judicious use of fluids improves outcomes in the older population more than in the young population.<sup>79</sup> Goal-directed fluid therapy reduced the length of stay by an average of 2 days in general surgery cases where the mean age was 56-59 yr.<sup>80</sup> In an older group of patients (mean age, 75 yr) undergoing repair of femoral fractures, use of goal-directed therapy shortened the length of stay by 8 days.<sup>81</sup> Consequently, a strategy of administering fluids in a manner that maintains optimal hemodynamics and end-organ perfusion is recommended.

Anemia is common in the older population. More than 8% of men and 6% of women greater than 65 yr of age, and without severe comorbidities, have anemia as defined by hemoglobin levels less than 10 g/dl.<sup>82</sup> Perioperative anemia in the aged population is associated with worse outcome.<sup>83</sup> However, perioperative anemia results in an increase in erythrocyte transfusions, which are also correlated with adverse outcomes including SSI.<sup>84</sup> Low level of hemoglobin in young healthy subjects does not reduce subcutaneous tissue oxygenation,<sup>85</sup> suggesting that erythrocyte transfusions are not indicated to enhance wound healing. The optimal hemoglobin level to maximize wound healing in older patients remains to be elucidated.

#### **Temperature**

Mild perioperative hypothermia is common not only during general anesthesia but also during regional anesthesia.<sup>86</sup> Age is an independent risk factor for the development of hypothermia during anesthesia.<sup>87</sup> Mild hypothermia during the intraoperative period is associated with vasoconstriction as measured by skin temperature and subcutaneous tissue oxygen.<sup>88</sup> This markedly increases the risk of surgical wound infection, even after clean procedures such as hernia, breast, and varicose vein surgeries.<sup>89</sup> Thermoregulatory responses are decreased in the aged subjects,<sup>90</sup> mostly due to altered regulation of skin blood flow in the setting of a reduced microcirculation.<sup>91</sup> During general anesthesia with isoflurane<sup>92</sup> and sevoflurane,<sup>93</sup> the threshold for thermoregulatory vasoconstriction is reduced in the aged more than the young. The aged subjects are at additional risk of perioperative hypothermia because clinical signs (such as shivering) are absent at the same time thermoregulation is impaired.<sup>94</sup> Rewarming of the older patient takes significantly longer than younger adults, reflecting the same physiology that predisposes older adults to hypothermia.95 Consequently, it is prudent to maintain euthermia for every aged patients during the intraoperative and postoperative periods, regardless of the type of anesthesia.

Strategies to maintain normothermia during anesthesia must take into account specific changes in the microcirculation. The <u>initial decrease</u> in core temperature results from the <u>redistribution</u> of heat to the peripheral microcirculation. Prewarming in the preoperative area might prevent redistribution of core heat.<sup>96</sup> Combined strategies that use multiple modalities (prewarming with use of warmed fluids and forced-air warming devices), are more effective in prolonged surgeries and in the older population.<sup>97</sup>

#### Anesthetic Management Impacts the Microcirculation and Subsequent Wound Healing

The role of anesthesia technique in postoperative wound repair is not well studied. However, several lines of retrospective data analyses of outcomes after cancer surgery suggest that anesthesia technique can influence mechanisms that are relevant to tissue repair processes.<sup>98</sup> Wound healing and cancer progression share several pathways: cellular proliferation and migration are accelerated; the extracellular matrix undergoes greater turnover; and neovascularization is enhanced. These effects are mediated by a surge of inflammatory mediators, cytokines, and growth factors that are common to both wound repair and cancer. Although the endpoints in these studies were cancer recurrence or metastases, the subsequent changes in growth factor levels or activity of matrix-degrading MMPs<sup>99,100</sup> are likely relevant to tissue healing. It is notable, however, that the groups studied differed not only in the use of inhalational versus IV agents

but also in the use of other variables such as the amount of opiates used to control pain. Each of these components has collateral effects including changes in regional blood flow and the effective microcirculation. Nonetheless, the underlying mechanisms that influence cancer surgery outcomes should be noted when examining determinants of postoperative wound healing.

The Effect of Anesthesia Technique: General versus Regional. Although clinical and theoretical perceptions often advocate for regional anesthesia rather than general anesthesia in older patients, there is no difference in various outcome measures.<sup>101,102</sup> Comparisons are difficult: for example, in a retrospective analysis, neuroaxial anesthesia (epidural or spinal anesthesia) for total hip or knee replacement was associated with a lower risk of SSI than general anesthesia, but the general anesthesia group was older with more comorbidities making it difficult to form definitive recommendations.<sup>103</sup>

Effects of anesthesia on physiologic variables relevant to the microcirculation and subsequent wound repair have been examined and will be discussed here. For example, general anesthesia causes vasodilation by direct effects on the peripheral microcirculation<sup>104</sup> and indirectly by central inhibition of vasoconstriction.<sup>105</sup> Thoracic epidural anesthesia increases peripheral tissue oxygen tension, even outside the dermatome affected by the block.<sup>106</sup> Core hypothermia develops equally under general and epidural anesthesia due to vasodilation in the microcirculation in the skin and loss of thermoregulation. During epidural anesthesia, skin thermoregulation in the region affected by the epidural (lower body) is reduced independent of patient age. Not surprisingly, young patients are better able than older patients to maintain skin thermoregulation in the regions not affected by epidural anesthesia.<sup>107</sup>

Administration of typical doses of volatile or IV agents does not suppress the contribution of the endocrine response to the microcirculation.<sup>108</sup> In contrast, regional anesthesia (most notably neuroaxial blockade) blunts the endocrine stress response to surgery.<sup>109</sup> Continuous lumbar plexus and sciatic nerve blocks did not affect cortisol levels but attenuated the postoperative inflammatory response (lower C-reactive protein).<sup>110</sup> In a study of regional block after knee arthroplasty, clinical signs of inflammation were reduced although there were no detectable changes in levels of measured cytokines.<sup>111</sup> Epidural anesthesia reduces the plasma levels of norepinephrine during surgery when compared with general anesthesia.<sup>112</sup> This holds true when combined general and epidural anesthesia is compared with general anesthesia alone.<sup>113</sup> Norepinephrine and epinephrine induce vasoconstriction in the microcirculation<sup>114</sup> and also have effects on subsequent biological processes that influence wound repair, such as angiogenesis and inflammation. For example, norepinephrine significantly increased secretion of the angiogenic factor VEGF by ovarian cancer and melanoma cell lines.<sup>115,116</sup> Furthermore, epinephrine suppresses phagocytosis of soluble immune complexes (aggregated  $\gamma$ -globulin) by macrophages in a dose-dependent manner<sup>117</sup> (fig. 4).

**Volatile Anesthetics Agents.** Volatile anesthetic agents have several contradictory effects on microcirculatory flow. They decrease flow in the microcirculation by reducing arterial perfusion pressure and depressing myocardial contractility. At the same time, these agents can increase flow in the microcirculation by inducing a vasodilatory response.<sup>118</sup> As a result, clinically useful concentrations of volatile agents will often produce systemic hypotension and decrease regional tissue perfusion in a tissue- and agent-specific manner.<sup>119</sup> Muscle perfusion under anesthesia with volatile agents is better maintained in young subjects than in aged subjects.<sup>120</sup>

Not all effects of inhalation anesthesia are detrimental to tissue repair (fig. 5). Volatile anesthetics protect against ischemia-reperfusion injury of several organs (heart,<sup>121</sup> liver,<sup>122</sup> kidney,<sup>123</sup> and potentially others) by reducing necrosis and inflammation.<sup>124</sup> Several lines of evidence suggest that inhalation anesthetics exert their effects by affecting the microcirculation and influencing subsequent angiogenesis. For example, exposure to volatile anesthetics stimulates growth and proliferation of endothelial progenitor cells.<sup>125</sup> Conversely, volatile anesthetics, at clinically relevant doses, block transcription factor hypoxia-inducible factor-1 activity and expression of its downstream target genes. Hypoxia-inducible factor-1 is a transcriptional regulator of VEGF expression<sup>126</sup> and mediates angiogenic responses to reduced oxygen availability.<sup>127</sup> When renal proximal tubule cells were exposed to volatile anesthetics for 16h, there was an increased production and release of TGF-\$1, a potent stimulator of extracellular matrix synthesis, into the cell culture media. The role of TGF- $\beta$ 1 in the protective effect of volatile agents on ischemic injury to the kidney was specifically demonstrated in vivomice heterozygous for TGF- $\beta$ 1 (TGF- $\beta$ 1+/-) were not protected from ischemia-reperfusion injury by sevoflurane, and a neutralizing TGF-B1 antibody blocked renal protection with sevoflurane in TGF- $\beta$ 1+/+ mice.<sup>128</sup> Despite the data above, it is worth noting that volatile anesthetic agents differ



Fig. 4. The effects of local anesthetics and regional anesthesia on wound healing.



**Fig. 5.** The effects of volatile anesthetics on wound healing. ECM = extracellular matrix; HIF-1 = hypoxia-inducible factor-1; TGF- $\beta$ 1 = transforming growth factor- $\beta$ 1; VEGF = vascular endothelial growth factor.

in their effects on the microcirculation with respect to specific agents (desflurane *vs.* isoflurane *vs.* sevoflurane), target organs (heart *vs.* kidney *vs.* splanchnic), and clinical states (healthy *vs.* sepsis *vs.* hemorrhagic states).<sup>129</sup> Consequently, we cannot make meaningful recommendations regarding volatile anesthetics and wound-healing outcomes.

**Opioids.** Large doses of opioids are routinely used to achieve hemodynamic stability during the intraoperative period. Compared with other commonly used IV agents, opioids have the advantage of not depressing cardiac contractility.<sup>130</sup> Opioids have multiple effects on the microcirculation, which include inducing the release of nitric oxide<sup>131</sup> and directly relaxing smooth muscle cells.<sup>132</sup> Opioids can both increase and decrease body temperature. The hyperthermic response to opioids is mediated by the µ-receptor and the hypothermic response is mediated by the K-receptor.<sup>133</sup> Morphine has several downstream effects that can influence wound repair, such as activation of G-protein-coupled receptors to induce cellular proliferation<sup>134</sup> and activation of the VEGF receptor.<sup>135</sup> Activation of a receptor in a ligand-independent manner can occur through different mechanisms and includes cross-talk between signaling pathways. For example interleukin-8, an angiogenic factor, can transactivate VEGF receptor-2.136 Human endothelial cells express the opioid responsive µ-3 receptor,<sup>137</sup> but the effect of opioids on angiogenesis itself is still under investigation. Some suggest that opioids inhibit blood vessel growth<sup>138</sup> and that opioid antagonists promote angiogenesis,<sup>139</sup> probably via suppression of hypoxia-inducible factor-1 α-mediated VEGF transcription.<sup>140</sup> Others found that opioids stimulate angiogenesis and tumor growth by inhibition of apoptosis and promotion of cell cycle progression in breast cancers.<sup>141,142</sup>

There are conflicting findings regarding the direct effect of opioids on wound repair. The expression of  $\mu$ -receptors is decreased in the margins of chronic wounds.<sup>143</sup> Some describe opioid-induced, delayed wound healing that is dose dependent.<sup>144</sup> The latter is proposed to be mediated, in part, by the inhibition of peripheral neuropeptide release into the healing wound and reduced neurokinin receptor expression in inflammatory and parenchymal cells.<sup>145</sup> Others suggest that topical opioids accelerate wound healing<sup>146</sup> by up-regulating nitric oxide synthase and the VEGF receptor-2.<sup>147</sup>

Pain<sup>148</sup> (as well as psychological stress)<sup>149</sup> delays wound healing. Conversely, inappropriately high doses of opioids can also impair organ function, delay mobilization, and subsequent healing after surgery.<sup>150</sup> A recent meta-analysis that compared analgesia after surgery with opioids alone as compared with a multimodal approach (in which an epidural was used in addition to general anesthesia) did not find a difference in mortality between the groups but found a lower complication rate in the multimodal group.<sup>151</sup> Wound-healing rates were not reported in the meta-analysis. Further studies are needed to elucidate the effects of opioids and other painalleviating modalities on subsequent wound repair.

**Other IV Anesthetic/Sedative Agents.** It is common for combinations of different agents to be used during induction and maintenance of anesthesia.<sup>152</sup> To our knowledge, the effects of the IV agents on the microcirculation have been best studied in different models of shock states, such as profound hemorrhage. Consequently, the observed effects are usually transient and are of uncertain relevance to wound healing.

In general, most agents either produce no change in hemodynamic parameters or cause vasodilation and cardiac depression. In normovolemic rats, regional blood flow was similar for all anesthetic agents<sup>153</sup> although some authors have described skeletal muscle arteriolar vasodilation with propofol. In a rat model of abdominal surgery, dexmedetomidine attenuated the reduction of the microcirculatory blood flow to the intestinal mucosa.154 Ketamine is an exception as it usually induces vasoconstriction and produces an increase in blood pressure and cardiac output.<sup>155</sup> In certain models of hemorrhagic shock, at similar systolic blood pressures, propofol produced regional vasodilation, whereas ketamine resulted in vasoconstriction.<sup>156</sup> Of note, the potential benefit of propofol on end-organ perfusion may be offset by the finding that macromolecular leak occurred from venules during propofol/fentanyl administration, but not during ketamine anesthesia.<sup>157</sup> Midazolam (but not ketamine or propofol) stimulated the release of VEGF by smooth muscle cell cultures, which might induce further downstream changes in the microcirculation.<sup>158</sup> VEGF induces angiogenic responses by inducing proliferation and migration and inhibiting apoptosis of endothelial cells.<sup>159</sup>

Intravenous agents also have numerous effects on immune responses that might be relevant to subsequent wound healing. *In vitro*, clinically relevant concentrations of ketamine and midazolam (but not propofol) inhibited monocyte chemotaxis.<sup>160</sup> Thiopental and etomidate (but not ketamine, propofol, or methohexital) decreased chemotaxis by eosinophils.<sup>161</sup> Ketamine inhibits the inflammatory response of macrophages<sup>162</sup> and antigen-presenting cells.<sup>163</sup> Thiopental and midazolam, but not ketamine, suppressed neutrophil function.<sup>164</sup> In summary, the choice of IV anesthetic agents could potentially have beneficial effects on the microcirculation and subsequent wound healing. It is premature to make any specific recommendations, as cell and animal studies are insufficient and well-designed clinical studies do not exist.

**Local Anesthetics.** The effect of local anesthetic infiltration on wound healing has been studied in numerous models with conflicting results. Some suggest that exposure to local anesthetics enhances wound repair, others propose no effect or a negative impact.<sup>165</sup> Local anesthetics may positively influence wound healing by reducing the stress response and alleviating pain.<sup>166</sup> Intraarticular lidocaine, used to achieve pain management after knee surgery, increased oxygen tension in the subcutaneous tissue.<sup>167</sup> Local anesthetics can be detrimental by delaying the synthesis of collagen<sup>168</sup> and by an antiproliferative effect on mesenchymal cells.<sup>169</sup>

The response of the microcirculation to local anesthetics is not consistent. The first-known local anesthetic, cocaine,<sup>170</sup> induces vasoconstriction, even at small doses, by inhibition of norepinephrine uptake.<sup>171</sup> Modern local anesthetics (lidocaine<sup>172</sup> and bupivacaine<sup>173</sup>) have a dose-dependent effect: low concentrations cause vasoconstriction of arterioles, whereas high concentrations cause vasodilation. Human studies suggest that dose-dependent properties of lidocaine may be pronounced in aged tissues. This likely reflects a longer drug half-life in older individuals, as a result of agerelated decreases in hepatic blood flow and clearance.<sup>174</sup>

#### Summary

Nearly every anesthesiologist who provides care to adults will participate in the care of geriatric patients.<sup>175</sup> A growing older population is undergoing surgical procedures that are increasing in number and complexity.<sup>176</sup> Poor healing of surgical wounds is a major cause of morbidity, mortality, and substantial economic burden. Wound healing is dependent on the microcirculation that supplies the incision area. Measures that support the microcirculation during the perioperative period have a profound effect on wound healing. Some measures such as maintenance of normal body temperature and control of postoperative pain are supported by ample evidence and have been implemented in routine clinical care. Other measures, for example, the choice of anesthesia technique and use of opioids are supported by basic research but need further clinical studies. A better understanding of the effect of aging and anesthesia on the microcirculation can potentially assist in improving postoperative wound repair, thereby benefiting a growing older population.

#### Acknowledgments

The authors thank Mamatha Damodarasamy M.S., Division of Gerontology and Geriatric Medicine, Department of Medicine, Harborview Medical Center, University of Washington, Seattle, Washington, for her assistance and Yotam Bentov, Tufts University, for his skillful graphical support in designing and creating the figures for this article. This work was supported by the National Institute of Aging, Bethesda, Maryland (grant No. R03 AG042353 to Dr. Bentov), and by the Institute of Translational Health Sciences and the Department of Anesthesiology and Pain, University of Washington, Seattle, Washington (pilot grant to Dr. Bentov).

#### **Competing Interests**

The authors declare no competing interests.

#### Correspondence

Address correspondence to Dr. Bentov: Department of Anesthesia and Pain Medicine, University of Washington, Harborview Medical Center, 325 9th Avenue, Box 359724, Seattle, Washington 98104. itayb@uw.edu. Information on purchasing reprints may be found at www.anesthesiology. org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY'S articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

#### References

- 1. McGarry SA, Engemann JJ, Schmader K, Sexton DJ, Kaye KS: Surgical-site infection due to *Staphylococcus aureus* among elderly patients: Mortality, duration of hospitalization, and cost. Infect Control Hosp Epidemiol 2004; 25:461–7
- Lober CW, Fenske NA: Cutaneous aging: Effect of intrinsic changes on surgical considerations. South Med J 1991; 84:1444-6
- 3. Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ: The impact of surgical-site infections in the 1990s: Attributable mortality, excess length of hospitalization, and extra costs. Infect Control Hosp Epidemiol 1999; 20:725–30
- Perencevich EN, Sands KE, Cosgrove SE, Guadagnoli E, Meara E, Platt R: Health and economic impact of surgical site infections diagnosed after hospital discharge. Emerg Infect Dis 2003; 9:196–3
- 5. Zouboulis CC, Makrantonaki E: Clinical aspects and molecular diagnostics of skin aging. Clin Dermatol 2011; 29:3–14
- Montagna W, Carlisle K: Structural changes in ageing skin. Br J Dermatol 1990; 122(suppl 35):61–70
- 7. Tsuchida Y: The effect of aging and arteriosclerosis on human skin blood flow. J Dermatol Sci 1993; 5:175-81
- 8. Sørensen LT: Wound healing and infection in surgery. The clinical impact of smoking and smoking cessation: A systematic review and meta-analysis. Arch Surg 2012; 147:373–83
- Longnecker DE: Effects of general anesthetics on the microcirculation. Microcirc Endothelium Lymphatics 1984; 1:129–50
- 10. Du Noüy PL: Cicatrization of wounds: III. The relation between the age of the patient, the area of the wound, and the index of cicatrization. J Exp Med 1916; 24:461–70
- 11. Mendoza CB Jr, Postlethwait RW, Johnson WD: Veterans Administration cooperative study of surgery for duodenal ulcer. II. Incidence of wound disruption following operation. Arch Surg 1970; 101:396–8
- Kaye KS, Anderson DJ, Sloane R, Chen LF, Choi Y, Link K, Sexton DJ, Schmader KE: The effect of surgical site infection on older operative patients. J Am Geriatr Soc 2009; 57:46–54
- Singer AJ, Clark RA: Cutaneous wound healing. N Engl J Med 1999; 341:738–46
- 14. Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, Richards DR, McDonald-Smith GP, Gao H, Hennessy L, Finnerty CC, López CM, Honari S, Moore EE, Minei JP, Cuschieri J, Bankey PE, Johnson JL, Sperry J, Nathens AB, Billiar TR, West MA, Jeschke MG, Klein MB, Gamelli RL,

Gibran NS, Brownstein BH, Miller-Graziano C, Calvano SE, Mason PH, Cobb JP, Rahme LG, Lowry SF, Maier RV, Moldawer LL, Herndon DN, Davis RW, Xiao W, Tompkins RG; Inflammation and Host Response to Injury, Large Scale Collaborative Research Program: Genomic responses in mouse models poorly mimic human inflammatory diseases. Proc Natl Acad Sci U S A 2013; 110:3507–12

- Gottrup F, Agren MS, Karlsmark T: Models for use in wound healing research: A survey focusing on *in vitro* and *in vivo* adult soft tissue. Wound Repair Regen 2000; 8:83–96
- Tarnuzzer RW, Schultz GS: Biochemical analysis of acute and chronic wound environments. Wound Repair Regen 1996; 4:321–5
- Swift ME, Burns AL, Gray KL, DiPietro LA: Age-related alterations in the inflammatory response to dermal injury. J Invest Dermatol 2001; 117:1027–35
- Cohen BJ, Danon D, Roth GS: Wound repair in mice as influenced by age and antimacrophage serum. J Gerontol 1987; 42:295–301
- Danon D, Kowatch MA, Roth GS: Promotion of wound repair in old mice by local injection of macrophages. Proc Natl Acad Sci U S A 1989; 86:2018–20
- 20. Keylock KT, Vieira VJ, Wallig MA, DiPietro LA, Schrementi M, Woods JA: Exercise accelerates cutaneous wound healing and decreases wound inflammation in aged mice. Am J Physiol Regul Integr Comp Physiol 2008; 294:R179–84
- Emery CF, Kiecolt-Glaser JK, Glaser R, Malarkey WB, Frid DJ: Exercise accelerates wound healing among healthy older adults: A preliminary investigation. J Gerontol A Biol Sci Med Sci 2005; 60:1432–6
- 22. Ashcroft GS, Mills SJ: Androgen receptor-mediated inhibition of cutaneous wound healing. J Clin Invest 2002; 110:615–24
- 23. Lai JJ, Lai KP, Chuang KH, Chang P, Yu IC, Lin WJ, Chang C: Monocyte/macrophage androgen receptor suppresses cutaneous wound healing in mice by enhancing local TNF-alpha expression. J Clin Invest 2009; 119:3739–51
- Ashcroft GS, Horan MA, Ferguson MW: The effects of ageing on wound healing: Immunolocalisation of growth factors and their receptors in a murine incisional model. J Anat 1997; 190 (Pt 3):351–65
- 25. Hsieh MM, Everhart JE, Byrd-Holt DD, Tisdale JF, Rodgers GP: Prevalence of neutropenia in the U.S. population: Age, sex, smoking status, and ethnic differences. Ann Intern Med 2007; 146:486–92
- 26. Silverman EM, Silverman AG: Granulocyte adherence in the elderly. Am J Clin Pathol 1977; 67:49–52
- Butcher SK, Chahal H, Nayak L, Sinclair A, Henriquez NV, Sapey E, O'Mahony D, Lord JM: Senescence in innate immune responses: Reduced neutrophil phagocytic capacity and CD16 expression in elderly humans. J Leukoc Biol 2001; 70:881–6
- Clark RA: Basics of cutaneous wound repair. J Dermatol Surg Oncol 1993; 19:693–706
- 29. Gillitzer R, Goebeler M: Chemokines in cutaneous wound healing. J Leukoc Biol 2001; 69:513–21
- McClain SA, Simon M, Jones E, Nandi A, Gailit JO, Tonnesen MG, Newman D, Clark RA: Mesenchymal cell activation is the rate-limiting step of granulation tissue induction. Am J Pathol 1996; 149:1257–70
- 31. Mussini E, Hutton JJ Jr, Udenfriend S: Collagen proline hydroxylase in wound healing, granuloma formation, scurvy, and growth. Science 1967; 157:927–9
- 32. Reed MJ, Koike T, Puolakkainen P: Wound repair in aging. A review. Methods Mol Med 2003; 78:217–37
- Bruce SA, Deamond SF: Longitudinal study of *in vivo* wound repair and *in vitro* cellular senescence of dermal fibroblasts. Exp Gerontol 1991; 26:17–27
- Holt DR, Kirk SJ, Regan MC, Hurson M, Lindblad WJ, Barbul A: Effect of age on wound healing in healthy human beings. Surgery 1992; 112:293–7; discussion 297–8

- Swift ME, Kleinman HK, DiPietro LA: Impaired wound repair and delayed angiogenesis in aged mice. Lab Invest 1999; 79:1479–87
- 36. Reed MJ, Corsa A, Pendergrass W, Penn P, Sage EH, Abrass IB: Neovascularization in aged mice: Delayed angiogenesis is coincident with decreased levels of transforming growth factor beta1 and type I collagen. Am J Pathol 1998; 152:113–23
- 37. Ashcroft GS, Horan MA, Ferguson MW: Aging is associated with reduced deposition of specific extracellular matrix components, an upregulation of angiogenesis, and an altered inflammatory response in a murine incisional wound healing model. J Invest Dermatol 1997; 108:430–7
- Rivard A, Fabre JE, Silver M, Chen D, Murohara T, Kearney M, Magner M, Asahara T, Isner JM: Age-dependent impairment of angiogenesis. Circulation 1999; 99:111–20
- 39. Arthur WT, Vernon RB, Sage EH, Reed MJ: Growth factors reverse the impaired sprouting of microvessels from aged mice. Microvasc Res 1998; 55:260–70
- Reiser K, McGee C, Rucker R, McDonald R: Effects of aging and caloric restriction on extracellular matrix biosynthesis in a model of injury repair in rats. J Gerontol A Biol Sci Med Sci 1995; 50A:B40–7
- 41. Ashcroft GS, Dodsworth J, van Boxtel E, Tarnuzzer RW, Horan MA, Schultz GS, Ferguson MW: Estrogen accelerates cutaneous wound healing associated with an increase in TGF-beta1 levels. Nat Med 1997; 3:1209–15
- 42. Reed MJ, Vernon RB, Abrass IB, Sage EH: TGF-beta 1 induces the expression of type I collagen and SPARC, and enhances contraction of collagen gels, by fibroblasts from young and aged donors. J Cell Physiol 1994; 158:169–79
- 43. Trombetta-Esilva J, Bradshaw AD: The function of SPARC as a mediator of fibrosis. Open Rheumatol J 2012; 6:146–55
- 44. Agah A, Kyriakides TR, Letrondo N, Björkblom B, Bornstein P: Thrombospondin 2 levels are increased in aged mice: Consequences for cutaneous wound healing and angiogenesis. Matrix Biol 2004; 22:539–47
- 45. Kyriakides TR, Bornstein P: Matricellular proteins as modulators of wound healing and the foreign body response. Thromb Haemost 2003; 90:986–92
- 46. Sprenger CC, Plymate SR, Reed MJ: Aging-related alterations in the extracellular matrix modulate the microenvironment and influence tumor progression. Int J Cancer 2010; 127:2739–48
- 47. Reed MJ, Damodarasamy M, Chan CK, Johnson MN, Wight TN, Vernon RB: Cleavage of hyaluronan is impaired in aged dermal wounds. Matrix Biol 2013; 32:45–51
- Parrinello S, Coppe JP, Krtolica A, Campisi J: Stromalepithelial interactions in aging and cancer: Senescent fibroblasts alter epithelial cell differentiation. J Cell Sci 2005; 118(Pt 3):485–96
- 49. Ashcroft GS, Horan MA, Herrick SE, Tarnuzzer RW, Schultz GS, Ferguson MW: Age-related differences in the temporal and spatial regulation of matrix metalloproteinases (MMPs) in normal skin and acute cutaneous wounds of healthy humans. Cell Tissue Res 1997; 290:581–91
- Ashcroft GS, Herrick SE, Tarnuzzer RW, Horan MA, Schultz GS, Ferguson MW: Human ageing impairs injury-induced *in vivo* expression of tissue inhibitor of matrix metalloproteinases (TIMP)-1 and -2 proteins and mRNA. J Pathol 1997; 183:169–76
- 51. Reed MJ, Karres N, Eyman D, Vernon RB, Edelberg JM: Agerelated differences in repair of dermal wounds and myocardial infarcts attenuate during the later stages of healing. In Vivo 2006; 20(6B):801–6
- Hunt TK, Hopf HW: Wound healing and wound infection. What surgeons and anesthesiologists can do. Surg Clin North Am 1997; 77:587–6
- 53. Leaper DJ: Risk factors for and epidemiology of surgical site infections. Surg Infect (Larchmt) 2010; 11:283–7

- 54. Davis GB, Peric M, Chan LS, Wong AK, Sener SF: Identifying risk factors for surgical site infections in mastectomy patients using the National Surgical Quality Improvement Program database. Am J Surg 2013; 205:194–9
- 55. Tandara AA, Mustoe TA: Oxygen in wound healing—More than a nutrient. World J Surg 2004; 28:294–300
- Sundaresan M, Yu ZX, Ferrans VJ, Sulciner DJ, Gutkind JS, Irani K, Goldschmidt-Clermont PJ, Finkel T: Regulation of reactive-oxygen-species generation in fibroblasts by Rac1. Biochem J 1996; 318 (Pt 2):379–82
- 57. Babior BM: Oxygen-dependent microbial killing by phagocytes (first of two parts). N Engl J Med 1978; 298:659–68
- 58. Hopf HW, Hunt TK, West JM, Blomquist P, Goodson WH III, Jensen JA, Jonsson K, Paty PB, Rabkin JM, Upton RA, von Smitten K, Whitney JD: Wound tissue oxygen tension predicts the risk of wound infection in surgical patients. Arch Surg 1997; 132:997–4; discussion 1005
- 59. Qadan M, Akça O, Mahid SS, Hornung CA, Polk HC Jr: Perioperative supplemental oxygen therapy and surgical site infection: A meta-analysis of randomized controlled trials. Arch Surg 2009; 144:359–66; discussion 366–7
- 60. Togioka B, Galvagno S, Sumida S, Murphy J, Ouanes JP, Wu C: The role of perioperative high inspired oxygen therapy in reducing surgical site infection: A meta-analysis. Anesth Analg 2012; 114:334–42
- Greif R, Akça O, Horn EP, Kurz A, Sessler DI; Outcomes Research Group: Supplemental perioperative oxygen to reduce the incidence of surgical-wound infection. N Engl J Med 2000; 342:161–7
- 62. Belda FJ, Aguilera L, García de la Asunción J, Alberti J, Vicente R, Ferrándiz L, Rodríguez R, Company R, Sessler DI, Aguilar G, Botello SG, Ortí R; Spanish Reduccion de la Tasa de Infeccion Quirurgica Group: Supplemental perioperative oxygen and the risk of surgical wound infection: A randomized controlled trial. JAMA 2005; 294:2035–42
- 63. Pryor KO, Fahey TJ III, Lien CA, Goldstein PA: Surgical site infection and the routine use of perioperative hyperoxia in a general surgical population: A randomized controlled trial. JAMA 2004; 291:79–7
- 64. Meyhoff CS, Wetterslev J, Jorgensen LN, Henneberg SW, Høgdall C, Lundvall L, Svendsen PE, Mollerup H, Lunn TH, Simonsen I, Martinsen KR, Pulawska T, Bundgaard L, Bugge L, Hansen EG, Riber C, Gocht-Jensen P, Walker LR, Bendtsen A, Johansson G, Skovgaard N, Heltø K, Poukinski A, Korshin A, Walli A, Bulut M, Carlsson PS, Rodt SA, Lundbech LB, Rask H, Buch N, Perdawid SK, Reza J, Jensen KV, Carlsen CG, Jensen FS, Rasmussen LS; PROXI Trial Group: Effect of high perioperative oxygen fraction on surgical site infection and pulmonary complications after abdominal surgery: The PROXI randomized clinical trial. JAMA 2009; 302:1543–50
- Guénard H, Marthan R: Pulmonary gas exchange in elderly subjects. Eur Respir J 1996; 9:2573–7
- 66. McGuire DK, Levine BD, Williamson JW, Snell PG, Blomqvist CG, Saltin B, Mitchell JH: A 30-year follow-up of the Dallas Bedrest and Training Study: I. Effect of age on the cardiovascular response to exercise. Circulation 2001; 104:1350–7
- Poole D, Behnke B, Musch T: Capillary hemodynamics and oxygen pressures in the aging microcirculation. Microcirculation 2006; 13:289–99
- 68. Coggan AR, Spina RJ, King DS, Rogers MA, Brown M, Nemeth PM, Holloszy JO: Histochemical and enzymatic comparison of the gastrocnemius muscle of young and elderly men and women. J Gerontol 1992; 47:B71–6
- 69. Wu L, Xia YP, Roth SI, Gruskin E, Mustoe TA: Transforming growth factor-beta1 fails to stimulate wound healing and impairs its signal transduction in an aged ischemic ulcer model: Importance of oxygen and age. Am J Pathol 1999; 154:301–9
- Quirinia A, Viidik A: The impact of ischemia on wound healing is increased in old age but can be countered by hyperbaric oxygen therapy. Mech Ageing Dev 1996; 91:131–44

- McGee S, Abernethy WB III, Simel DL: The rational clinical examination. Is this patient hypovolemic? JAMA 1999; 281:1022–9
- Arkiliç CF, Taguchi A, Sharma N, Ratnaraj J, Sessler DI, Read TE, Fleshman JW, Kurz A: Supplemental perioperative fluid administration increases tissue oxygen pressure. Surgery 2003; 133:49–55
- 73. Brandstrup B, Tønnesen H, Beier-Holgersen R, Hjortsø E, Ørding H, Lindorff-Larsen K, Rasmussen MS, Lanng C, Wallin L, Iversen LH, Gramkow CS, Okholm M, Blemmer T, Svendsen PE, Rottensten HH, Thage B, Riis J, Jeppesen IS, Teilum D, Christensen AM, Graungaard B, Pott F; Danish Study Group on Perioperative Fluid Therapy: Effects of intravenous fluid restriction on postoperative complications: Comparison of two perioperative fluid regimens: A randomized assessorblinded multicenter trial. Ann Surg 2003; 238:641–8
- Grocott MP, Mythen MG, Gan TJ: Perioperative fluid management and clinical outcomes in adults. Anesth Analg 2005; 100:1093–106
- Green D, Paklet L: Latest developments in peri-operative monitoring of the high-risk major surgery patient. Int J Surg 2010; 8:90–9
- 76. Bundgaard-Nielsen M, Secher NH, Kehlet H: 'Liberal' vs. 'restrictive' perioperative fluid therapy—A critical assessment of the evidence. Acta Anaesthesiol Scand 2009; 53:843–51
- 77. Stotts NA, Hopf HW: The link between tissue oxygen and hydration in nursing home residents with pressure ulcers: Preliminary data. J Wound Ostomy Continence Nurs 2003; 30:184–90
- 78. Stotts NA, Hopf HW, Kayser-Jones J, Chertow GM, Cooper BA, Wu HS: Increased fluid intake does not augment capacity to lay down new collagen in nursing home residents at risk for pressure ulcers: A randomized, controlled clinical trial. Wound Repair Regen 2009; 17:780–8
- Spahn DR, Chassot PG: CON: Fluid restriction for cardiac patients during major noncardiac surgery should be replaced by goal-directed intravascular fluid administration. Anesth Analg 2006; 102:344–6
- Gan TJ, Soppitt A, Maroof M, el-Moalem H, Robertson KM, Moretti E, Dwane P, Glass PS: Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. ANESTHESIOLOGY 2002; 97:820–6
- Sinclair S, James S, Singer M: Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: Randomised controlled trial. BMJ 1997; 315:909–12
- 82. Endres HG, Wedding U, Pittrow D, Thiem U, Trampisch HJ, Diehm C: Prevalence of anemia in elderly patients in primary care: Impact on 5-year mortality risk and differences between men and women. Curr Med Res Opin 2009; 25:1143–58
- 83. Wu WC, Schifftner TL, Henderson WG, Eaton CB, Poses RM, Uttley G, Sharma SC, Vezeridis M, Khuri SF, Friedmann PD: Preoperative hematocrit levels and postoperative outcomes in older patients undergoing noncardiac surgery. JAMA 2007; 297:2481–8
- Dunne JR, Malone D, Tracy JK, Gannon C, Napolitano LM: Perioperative anemia: An independent risk factor for infection, mortality, and resource utilization in surgery. J Surg Res 2002; 102:237–44
- 85. Hopf HW, Viele M, Watson JJ, Feiner J, Weiskopf R, Hunt TK, Noorani M, Yeap H, Ho R, Toy P: Subcutaneous perfusion and oxygen during acute severe isovolemic hemodilution in healthy volunteers. Arch Surg 2000; 135:1443–9
- 86. Sessler DI: Mild perioperative hypothermia. N Engl J Med 1997; 336:1730–7
- 87. Frank SM, Beattie C, Christopherson R, Norris EJ, Rock P, Parker S, Kimball AW Jr: Epidural *versus* general anesthesia, ambient operating room temperature, and patient age as predictors of inadvertent hypothermia. ANESTHESIOLOGY 1992; 77:252–7

- Sheffield CW, Sessler DI, Hopf HW, Schroeder M, Moayeri A, Hunt TK, West JM: Centrally and locally mediated thermoregulatory responses alter subcutaneous oxygen tension. Wound Repair Regen 1996; 4:339–45
- Melling AC, Ali B, Scott EM, Leaper DJ: Effects of preoperative warming on the incidence of wound infection after clean surgery: A randomised controlled trial. Lancet 2001; 358:876–80
- Hajat S, Kovats RS, Lachowycz K: Heat-related and coldrelated deaths in England and Wales: Who is at risk? Occup Environ Med 2007; 64:93–100
- Holowatz LA, Thompson-Torgerson C, Kenney WL: Aging and the control of human skin blood flow. Front Biosci (Landmark Ed) 2010; 15:718–39
- 92. Kurz A, Plattner O, Sessler DI, Huemer G, Redl G, Lackner F: The threshold for thermoregulatory vasoconstriction during nitrous oxide/isoflurane anesthesia is lower in elderly than in young patients. ANESTHESIOLOGY 1993; 79:465–9
- 93. Ozaki M, Sessler DI, Matsukawa T, Ozaki K, Atarashi K, Negishi C, Suzuki H: The threshold for thermoregulatory vasoconstriction during nitrous oxide/sevoflurane anesthesia is reduced in the elderly. Anesth Analg 1997; 84:1029–33
- 94. Vassilieff N, Rosencher N, Sessler DI, Conseiller C: Shivering threshold during spinal anesthesia is reduced in elderly patients. ANESTHESIOLOGY 1995; 83:1162–6
- 95. Carli F, Gabrielczyk M, Clark MM, Aber VR: An investigation of factors affecting postoperative rewarming of adult patients. Anaesthesia 1986; 41:363–9
- 96. de Brito Poveda V, Clark AM, Galvão CM: A systematic review on the effectiveness of prewarming to prevent perioperative hypothermia. J Clin Nurs 2013; 22:906–18
- 97. Moola S, Lockwood C: Effectiveness of strategies for the management and/or prevention of hypothermia within the adult perioperative environment. Int J Evid Based Healthc 2011; 9:337–45
- 98. Snyder GL, Greenberg S: Effect of anaesthetic technique and other perioperative factors on cancer recurrence. Br J Anaesth 2010; 105:106–15
- 99. Looney M, Doran P, Buggy DJ: Effect of anesthetic technique on serum vascular endothelial growth factor C and transforming growth factor  $\beta$  in women undergoing anesthesia and surgery for breast cancer. ANESTHESIOLOGY 2010; 113:1118–25
- 100. Deegan CA, Murray D, Doran P, Moriarty DC, Sessler DI, Mascha E, Kavanagh BP, Buggy DJ: Anesthetic technique and the cytokine and matrix metalloproteinase response to primary breast cancer surgery. Reg Anesth Pain Med 2010; 35:490–5
- 101. O'Hara DA, Duff A, Berlin JA, Poses RM, Lawrence VA, Huber EC, Noveck H, Strom BL, Carson JL: The effect of anesthetic technique on postoperative outcomes in hip fracture repair. ANESTHESIOLOGY 2000; 92:947–57
- 102. Roy RC: Choosing general *versus* regional anesthesia for the elderly. Anesthesiol Clin North America 2000; 18:91–104, vii
- 103. Chang CC, Lin HC, Lin HW, Lin HC: Anesthetic management and surgical site infections in total hip or knee replacement: A population-based study. ANESTHESIOLOGY 2010; 113:279–84
- 104. Kobayashi Y, Yoshida K, Noguchi M, Wakasugi Y, Ito H, Okabe E: Effect of enflurane on contractile reactivity in isolated canine mesenteric arteries and veins. Anesth Analg 1990; 70:530–6
- 105. Xiong J, Kurz A, Sessler DI, Plattner O, Christensen R, Dechert M, Ikeda T: Isoflurane produces marked and nonlinear decreases in the vasoconstriction and shivering thresholds. ANESTHESIOLOGY 1996; 85:240–5
- 106. Treschan TA, Taguchi A, Ali SZ, Sharma N, Kabon B, Sessler DI, Kurz A: The effects of epidural and general anesthesia on tissue oxygenation. Anesth Analg 2003; 96:1553–7
- 107. Frank SM, Shir Y, Raja SN, Fleisher LA, Beattie C: Core hypothermia and skin-surface temperature gradients.

Epidural *versus* general anesthesia and the effects of age. ANESTHESIOLOGY 1994; 80:502–8

- Kehlet H: Manipulation of the metabolic response in clinical practice. World J Surg 2000; 24:690–5
- 109. Moraca RJ, Sheldon DG, Thirlby RC: The role of epidural anesthesia and analgesia in surgical practice. Ann Surg 2003; 238:663–73
- 110. Bagry H, de la Cuadra Fontaine JC, Asenjo JF, Bracco D, Carli F: Effect of a continuous peripheral nerve block on the inflammatory response in knee arthroplasty. Reg Anesth Pain Med 2008; 33:17–3
- 111. Martin F, Martinez V, Mazoit JX, Bouhassira D, Cherif K, Gentili ME, Piriou P, Chauvin M, Fletcher D: Antiinflammatory effect of peripheral nerve blocks after knee surgery: Clinical and biologic evaluation. ANESTHESIOLOGY 2008; 109:484–90
- 112. Breslow MJ, Parker SD, Frank SM, Norris EJ, Yates H, Raff H, Rock P, Christopherson R, Rosenfeld BA, Beattie C: Determinants of catecholamine and cortisol responses to lower extremity revascularization. The PIRAT Study Group. ANESTHESIOLOGY 1993; 79:1202–9
- 113. Yokoyama M, Itano Y, Katayama H, Morimatsu H, Takeda Y, Takahashi T, Nagano O, Morita K: The effects of continuous epidural anesthesia and analgesia on stress response and immune function in patients undergoing radical esophagectomy. Anesth Analg 2005; 101:1521–7
- 114. Jensen JA, Jonsson K, Goodson WH III, Hunt TK, Roizen MF: Epinephrine lowers subcutaneous wound oxygen tension. Curr Surg 1985; 42:472–4
- 115. Lutgendorf SK, Cole S, Costanzo E, Bradley S, Coffin J, Jabbari S, Rainwater K, Ritchie JM, Yang M, Sood AK: Stressrelated mediators stimulate vascular endothelial growth factor secretion by two ovarian cancer cell lines. Clin Cancer Res 2003; 9:4514–21
- 116. Yang EV, Kim SJ, Donovan EL, Chen M, Gross AC, Webster Marketon JI, Barsky SH, Glaser R: Norepinephrine upregulates VEGF, IL-8, and IL-6 expression in human melanoma tumor cell lines: Implications for stress-related enhancement of tumor progression. Brain Behav Immun 2009; 23:267–75
- 117. Abrass CK, O'Connor SW, Scarpace PJ, Abrass IB: Characterization of the beta-adrenergic receptor of the rat peritoneal macrophage. J Immunol 1985; 135:1338–41
- 118. Kawano T, Oshita S, Takahashi A, Tsutsumi Y, Tanaka K, Tomiyama Y, Kitahata H, Nakaya Y: Molecular mechanisms underlying ketamine-mediated inhibition of sarcolemmal adenosine triphosphate-sensitive potassium channels. ANESTHESIOLOGY 2005; 102:93–101
- 119. Hartman JC, Pagel PS, Proctor LT, Kampine JP, Schmeling WT, Warltier DC: Influence of desflurane, isoflurane and halothane on regional tissue perfusion in dogs. Can J Anaesth 1992; 39:877–87
- Dwyer R, Howe J: Peripheral blood flow in the elderly during inhalational anaesthesia. Acta Anaesthesiol Scand 1995; 39:939–44
- 121. De Hert SG, Turani F, Mathur S, Stowe DF: Cardioprotection with volatile anesthetics: Mechanisms and clinical implications. Anesth Analg 2005; 100:1584–93
- 122. Beck-Schimmer B, Breitenstein S, Urech S, De Conno E, Wittlinger M, Puhan M, Jochum W, Spahn DR, Graf R, Clavien PA: A randomized controlled trial on pharmacological preconditioning in liver surgery using a volatile anesthetic. Ann Surg 2008; 248:909–18
- 123. Lee HT, Ota-Setlik A, Fu Y, Nasr SH, Emala CW: Differential protective effects of volatile anesthetics against renal ischemia-reperfusion injury *in vivo*. ANESTHESIOLOGY 2004; 101:1313–24
- 124. Raphael J: Physiology and pharmacology of myocardial preconditioning. Semin Cardiothorac Vasc Anesth 2010; 14:54–9

- 125. Lucchinetti E, Zeisberger SM, Baruscotti I, Wacker J, Feng J, Zaugg K, Dubey R, Zisch AH, Zaugg M: Stem cell-like human endothelial progenitors show enhanced colony-forming capacity after brief sevoflurane exposure: Preconditioning of angiogenic cells by volatile anesthetics. Anesth Analg 2009; 109:1117–26
- 126. Puppo M, Battaglia F, Ottaviano C, Delfino S, Ribatti D, Varesio L, Bosco MC: Topotecan inhibits vascular endothelial growth factor production and angiogenic activity induced by hypoxia in human neuroblastoma by targeting hypoxia-inducible factor-1alpha and -2alpha. Mol Cancer Ther 2008; 7:1974–84
- 127. Itoh T, Namba T, Fukuda K, Semenza GL, Hirota K: Reversible inhibition of hypoxia-inducible factor 1 activation by exposure of hypoxic cells to the volatile anesthetic halothane. FEBS Lett 2001; 509:225–9
- 128. Lee HT, Kim M, Kim J, Kim N, Emala CW: TGF-beta1 release by volatile anesthetics mediates protection against renal proximal tubule cell necrosis. Am J Nephrol 2007; 27:416–24
- 129. Turek Z, Sykora R, Matejovic M, Cerny V: Anesthesia and the microcirculation. Semin Cardiothorac Vasc Anesth 2009; 13:249–58
- 130. Kawakubo A, Fujigaki T, Uresino H, Zang S, Sumikawa K: Comparative effects of etomidate, ketamine, propofol, and fentanyl on myocardial contractility in dogs. J Anesth 1999; 13:77–2
- 131. Stefano GB, Salzet M, Magazine HI, Bilfinger TV: Antagonism of LPS and IFN-gamma induction of iNOS in human saphenous vein endothelium by morphine and anandamide by nitric oxide inhibition of adenylate cyclase. J Cardiovasc Pharmacol 1998; 31:813–20
- 132. White DA, Reitan JA, Kien ND, Thorup SJ: Decrease in vascular resistance in the isolated canine hindlimb after graded doses of alfentanil, fentanyl, and sufentanil. Anesth Analg 1990; 71:29–4
- 133. Xin L, Geller EB, Adler MW: Body temperature and analgesic effects of selective mu and kappa opioid receptor agonists microdialyzed into rat brain. J Pharmacol Exp Ther 1997; 281:499–7
- Law PY, Loh HH: Regulation of opioid receptor activities. J Pharmacol Exp Ther 1999; 289:607–24
- 135. Singleton PA, Lingen MW, Fekete MJ, Garcia JG, Moss J: Methylnaltrexone inhibits opiate and VEGF-induced angiogenesis: Role of receptor transactivation. Microvasc Res 2006; 72:3–11
- 136. Petreaca ML, Yao M, Liu Y, Defea K, Martins-Green M: Transactivation of vascular endothelial growth factor receptor-2 by interleukin-8 (IL-8/CXCL8) is required for IL-8/ CXCL8-induced endothelial permeability. Mol Biol Cell 2007; 18:5014–23
- 137. Cadet P, Mantione KJ, Bilfinger TV, Stefano GB: Differential expression of the human mu opiate receptor from different primary vascular endothelial cells. Med Sci Monit 2004; 10:BR351–5
- 138. Pasi A, Qu BX, Steiner R, Senn HJ, Bär W, Messiha FS: Angiogenesis: Modulation with opioids. Gen Pharmacol 1991; 22:1077–9
- 139. Faramarzi N, Abbasi A, Tavangar SM, Mazouchi M, Dehpour AR: Opioid receptor antagonist promotes angiogenesis in bile duct ligated rats. J Gastroenterol Hepatol 2009; 24:1226–9
- 140. Roy S, Balasubramanian S, Wang J, Chandrashekhar Y, Charboneau R, Barke R: Morphine inhibits VEGF expression in myocardial ischemia. Surgery 2003; 134:336–44
- 141. Gupta K, Kshirsagar S, Chang L, Schwartz R, Law PY, Yee D, Hebbel RP: Morphine stimulates angiogenesis by activating proangiogenic and survival-promoting signaling and promotes breast tumor growth. Cancer Res 2002; 62:4491–8
- 142. Farooqui M, Li Y, Rogers T, Poonawala T, Griffin RJ, Song CW, Gupta K: COX-2 inhibitor celecoxib prevents chronic

morphine-induced promotion of angiogenesis, tumour growth, metastasis and mortality, without compromising analgesia. Br J Cancer 2007; 97:1523–31

- 143. Bigliardi PL, Sumanovski LT, Büchner S, Rufli T, Bigliardi-Qi M: Different expression of mu-opiate receptor in chronic and acute wounds and the effect of beta-endorphin on transforming growth factor beta type II receptor and cytokeratin 16 expression. J Invest Dermatol 2003; 120:145–52
- 144. Rook JM, McCarson KE: Delay of cutaneous wound closure by morphine *via* local blockade of peripheral tachykinin release. Biochem Pharmacol 2007; 74:752–7
- 145. Rook JM, Hasan W, McCarson KE: Morphine-induced early delays in wound closure: Involvement of sensory neuropeptides and modification of neurokinin receptor expression. Biochem Pharmacol 2009; 77:1747–55
- 146. Küchler S, Wolf NB, Heilmann S, Weindl G, Helfmann J, Yahya MM, Stein C, Schäfer-Korting M: 3D-wound healing model: Influence of morphine and solid lipid nanoparticles. J Biotechnol 2010; 148:24–30
- 147. Poonawala T, Levay-Young BK, Hebbel RP, Gupta K: Opioids heal ischemic wounds in the rat. Wound Repair Regen 2005; 13:165–74
- 148. McGuire L, Heffner K, Glaser R, Needleman B, Malarkey W, Dickinson S, Lemeshow S, Cook C, Muscarella P, Melvin WS, Ellison EC, Kiecolt-Glaser JK: Pain and wound healing in surgical patients. Ann Behav Med 2006; 31:165–72
- 149. Kiecolt-Glaser JK, Marucha PT, Malarkey WB, Mercado AM, Glaser R: Slowing of wound healing by psychological stress. Lancet 1995; 346:1194–6
- 150. Kehlet H: Multimodal approach to control postoperative pathophysiology and rehabilitation. Br J Anaesth 1997; 78:606–17
- 151. Nishimori M, Low JH, Zheng H, Ballantyne JC: Epidural pain relief *versus* systemic opioid-based pain relief for abdominal aortic surgery. Cochrane Database Syst Rev 2012; 7:CD005059
- 152. Eger EI II: Characteristics of anesthetic agents used for induction and maintenance of general anesthesia. Am J Health Syst Pharm 2004; 61(suppl 4):S3–10
- 153. Seyde WC, McGowan L, Lund N, Duling B, Longnecker DE: Effects of anesthetics on regional hemodynamics in normovolemic and hemorrhaged rats. Am J Physiol 1985; 249(1 Pt 2):H164–73
- 154. Yeh YC, Sun WZ, Ko WJ, Chan WS, Fan SZ, Tsai JC, Lin TY: Dexmedetomidine prevents alterations of intestinal microcirculation that are induced by surgical stress and pain in a novel rat model. Anesth Analg 2012; 115:46–3
- 155. Brookes ZL, Brown NJ, Reilly CS: Intravenous anaesthesia and the rat microcirculation: The dorsal microcirculatory chamber. Br J Anaesth 2000; 85:901–3
- 156. Brookes ZL, Brown NJ, Reilly CS: Response of the rat cremaster microcirculation to hemorrhage *in vivo*: Differential effects of intravenous anesthetic agents. Shock 2002; 18:542–8
- 157. Brookes ZL, Brown NJ, Reilly CS: Differential effects of intravenous anaesthetic agents on the response of rat mesenteric microcirculation *in vivo* after haemorrhage. Br J Anaesth 2002; 88:255–63
- 158. Tanabe K, Dohi S, Matsuno H, Hirade K, Kozawa O: Midazolam stimulates vascular endothelial growth factor release in aortic smooth muscle cells: Role of the mitogenactivated protein kinase superfamily. ANESTHESIOLOGY 2003; 98:1147–54
- 159. Ferrara N, Gerber HP, LeCouter J: The biology of VEGF and its receptors. Nat Med 2003; 9:669–76
- 160. Krumholz W, Reussner D, Hempelmann G: The influence of several intravenous anaesthetics on the chemotaxis of human monocytes *in vitro*. Eur J Anaesthesiol 1999; 16:547–9

- 161. Krumholz W, Abdulle O, Knecht J, Hempelmann G: Effects of i.v. anaesthetic agents on the chemotaxis of eosinophils *in vitro*. Br J Anaesth 1999; 83:333–5
- 162. Li CY, Chou TC, Wong CS, Ho ST, Wu CC, Yen MH, Ding YA: Ketamine inhibits nitric oxide synthase in lipopolysaccharide-treated rat alveolar macrophages. Can J Anaesth 1997; 44:989–95
- 163. Ohta N, Ohashi Y, Fujino Y: Ketamine inhibits maturation of bone marrow-derived dendritic cells and priming of the Th1type immune response. Anesth Analg 2009; 109:793–800
- 164. Nishina K, Akamatsu H, Mikawa K, Shiga M, Maekawa N, Obara H, Niwa Y: The inhibitory effects of thiopental, midazolam, and ketamine on human neutrophil functions. Anesth Analg 1998; 86:159–65
- 165. Brower MC, Johnson ME: Adverse effects of local anesthetic infiltration on wound healing. Reg Anesth Pain Med 2003; 28:233–40
- 166. Bamigboye AA, Hofmeyr GJ: Local anaesthetic wound infiltration and abdominal nerves block during caesarean section for postoperative pain relief. Cochrane Database Syst Rev 2009: CD006954
- 167. Akça O, Melischek M, Scheck T, Hellwagner K, Arkiliç CF, Kurz A, Kapral S, Heinz T, Lackner FX, Sessler DI: Postoperative pain and subcutaneous oxygen tension. Lancet 1999; 354:41–2
- 168. Chvapil M, Hameroff SR, O'Dea K, Peacock EE Jr: Local anesthetics and wound healing. J Surg Res 1979; 27:367–71

- 169. Lucchinetti E, Awad AE, Rahman M, Feng J, Lou PH, Zhang L, Ionescu L, Lemieux H, Thébaud B, Zaugg M: Antiproliferative effects of local anesthetics on mesenchymal stem cells: Potential implications for tumor spreading and wound healing. ANESTHESIOLOGY 2012; 116: 841–56
- 170. Calatayud J, González A: History of the development and evolution of local anesthesia since the coca leaf. ANESTHESIOLOGY 2003; 98:1503–8
- 171. Lange RA, Cigarroa RG, Yancy CW Jr, Willard JE, Popma JJ, Sills MN, McBride W, Kim AS, Hillis LD: Cocaine-induced coronary-artery vasoconstriction. N Engl J Med 1989; 321:1557–62
- 172. Johns RA, DiFazio CA, Longnecker DE: Lidocaine constricts or dilates rat arterioles in a dose-dependent manner. ANESTHESIOLOGY 1985; 62:141–4
- 173. Johns RA, Seyde WC, DiFazio CA, Longnecker DE: Dosedependent effects of bupivacaine on rat muscle arterioles. ANESTHESIOLOGY 1986; 65:186–91
- 174. Nation RL, Triggs EJ, Selig M: Lignocaine kinetics in cardiac patients and aged subjects. Br J Clin Pharmacol 1977; 4:439–48
- 175. Prough DS: Anesthetic pitfalls in the elderly patient. J Am Coll Surg 2005; 200:784–94
- 176. Etzioni DA, Liu JH, O'Connell JB, Maggard MA, Ko CY: Elderly patients in surgical workloads: A population-based analysis. Am Surg 2003; 69:961–5