David C. Warltier, M.D., Pb.D., Editor

Anesthesiology 2003; 99:1415-32

© 2003 American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Pulmonary Arterial Hypertension

Pathophysiology and Anesthetic Approach Gilbert Blaise, M.D.,* David Langleben, M.D.,† Bernard Hubert, M.D.‡

IN 1998, the World Health Organization sponsored a symposium on pulmonary hypertension at which a new, more clinically useful classification system was adopted.§ Traditionally, pulmonary hypertension had been classified as being primary or secondary. The classification proposed at the World Health Organization symposium divides pulmonary hypertension into five distinct categories (table 1): (1) pulmonary arterial hypertension (PAH) associated primary pulmonary hypertension (PPH) (familial and sporadic) and PAH related to collagen vascular disease, congenital systemic to pulmonary shunts, HIV infection, portopulmonary hypertension, drugs such as anorexigens, and toxins; (2) PAH linked with disorders of the respiratory system and/or hypoxemia; (3) pulmonary venous hypertension, including mitral valve disease, chronic left ventricular dysfunction, and pulmonary venoocclusive disease; (4) PAH due to chronic thrombotic and/or embolic disease; and (5) PAH attributed to disorders directly affecting the pulmonary vasculature (inflammatory pulmonary capillary hemangiomatosis).

Pathophysiology

Pulmonary arterial hypertension is characterized by a progressive increase in pulmonary arterial pressure

(PAP; mean pressure > 25 mmHg at rest or 30 mmHg during exercise) in association with variable degrees of pulmonary vascular remodelling, vasoconstriction, and *in situ* thrombosis.¹ No underlying cause can be found for some PAH, and secondary forms of PAH are related to collagen vascular disease, congenital systemic to pulmonary shunts, HIV infection, portopulmonary hypertension, and drugs.²

Primary pulmonary hypertension is a rare disease with an annual incidence of 1-2 per million. Six to 12% of cases are inherited in an autosomal dominant manner with reduced penetrance. PPH occurs three times more frequently in women than in men.³ Recently, mutations of the bone morphogenetic protein receptor type 2 gene (BMPR2) have been identified as causing many cases of familial PPH. BMPR2 encodes a type II receptor member of the transforming growth factor B superfamily of cellsignaling molecules⁴; after ligand binding, type II receptors form heteromeric complexes with membranebound type I receptors initiating phosphorylation of the type I receptor.⁵ This pathway seems to be critical in both cell differentiation and cell growth, with specificity mediated through transcriptional factors. The sporadic form of PPH is also associated with mutations of the gene encoding the protein receptor BMPR2 in at least 26% of cases.⁶⁻⁸ The 2000 International PPH Consortium pointed to the possibility that additional factors, either environmental or genetic, are required for disease pathogenesis. Recent publications have confirmed the important role of heredity in the disease.^{9,10} The sex bias for disease presentation suggests a role for either hormonal factors or an X-linked locus.³

In the normal pulmonary circulation, pressure and resistance are 80–90% lower than in the systemic circulation. Pulmonary arteries larger than 1 mm in ID are elastic in nature and have well-developed internal and external laminae with a less distinct medial layer than systemic arteries. Most pulmonary arteries run adjacent to the airways. Distal to the respiratory bronchioles, the smooth muscle layer is reduced, and the arteries are only partially muscularized or nonmuscularized.¹¹ Vascular tone is normally very low, even if the pulmonary vessels are highly reactive to hypoxia and endogenous constric-

^{*} Professor, Department of Anesthesia, Hôpital Notre-Dame du Centre hospitalier de l'Université de Montréal. † Associate Professor, Director of Department of Cardiology, Centre for Pulmonary Vascular Disease, Sir Mortimer B. Davis Jewish General Hospital, McGill University, Montreal, Quebec, Canada. ‡ Anesthesia-Intensive Care Intern, Department of Anesthesia and Intensive Care Medicine, CHU de Liège, Liege, Belgium.

Received from the Department of Anesthesiology and Medicine, Université de Montréal, Montreal, Quebec, Canada.

Submitted for publication March 27, 2001. Accepted for publication February 20, 2003. Dr. Blaise is supported by the Canadian Institutes of Health Research, Ottawa, Ontario, Canada. Dr. Langleben is a Chercheur-Boursier Clinicien of the Fonds de la recherche en Santé du Québec, Montreal, Quebec, Canada, and is supported in part by the Bank of Montreal Centre for the Study of Heart Disease in Women, Jewish General Hospital, McGill University, Montreal, Quebec, Canada.

Address reprint requests to Dr. Blaise: Department of Anesthesia, CHUM-Hôpital Notre-Dame, Pavillon Deschamps, local FS-1136, 1560 Sherbrooke East Street, Montreal, Quebec, Canada H2L 4M1. Address electronic mail to: blaisegi@sympatico.ca. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

[§]World Symposium on Primary Pulmonary Hypertension, 1998. Available at: http://www.who.int/ncd/cvd/pph.homl. Accessed December 5, 2000.

Table 1. Diagnosis Classification of Pulmonary Hypertension Proposed at the World Symposium on Primary Pulmonary Hypertension, 1998

Pulmonary arterial hypertension Primary pulmonary hypertension (sporadic, familial) Pulmonary arterial hypertension related to collagen vascular disease (scleroderma, lupus, rheumatoid arthritis) Congenital systemic-to-pulmonary shunts (Eisenmenger syndrome), portopulmonary hypertension HIV infection, drugs, and toxins
Pulmonary hypertension associated with disorders of the
respiratory system and/or hypoxemia
Parenchymal lung disease (chronic obstructive pulmonary disease, interstitial pulmonary fibrosis, and cystic fibrosis)
Chronic alveolar hypoxemia (exposure to long-term low oxygen tension such as in high altitudes)
Pulmonary venous hypertension
Mitral valve disease
Chronic left ventricular dysfunction
Pulmonary venoocclusion disease
Pulmonary hypertension due to chronic thrombotic and/or embolic disease
Thromboembolic obstruction of proximal pulmonary arteries
Obstruction of distal pulmonary arteries
Pulmonary hypertension due to disorders directly affecting the
Inflammatory
Pulmonary capillary hemangiomatosis

World Symposium on Primary Pulmonary Hypertension, 1998. Available at: http://www.who.int/ncd/cvd/pph.html. Accessed May 12, 2000.

tors.¹²⁻¹⁵ Sympathetic innervation of the pulmonary circulation does exist, and its activation increases pulmonary tone, although there are still controversies regarding the presence of α_1 receptors.^{16,17} There are α_2 -adrenergic receptors in human pulmonary tissue. However, their specific localization (nerves, vessels, bronchi) is not known.¹⁸ We are not aware of studies of the effects of α_2 agonists on isolated pulmonary vessels. Both PPH and other forms of PAH may share a common pathophysiology. It has been proposed that muscularization of the terminal portion of the pulmonary arterial vascular tree, caused by smooth muscle cell (SMC) hyperplasia, is the earliest change.¹⁹ There is great heterogeneity in the phenotype of pulmonary vascular SMCs,²⁰ depending on the size and location of the pulmonary arteries. Different phenotypes respond differently to vasoactive factors and exhibit differential pharmacology. For example, K⁺ channels are differentially distributed,²¹ and cell-specific differences are in the endothelin-1 (ET-1) system.²² In established PAH, the pulmonary arteries are characterized by intimal fibrosis, medial hypertrophy, adventitial proliferation, and obliteration of small arteries. Plexiform lesions are found in many PAH cases. These lesions resemble renal glomeruli with multiple channels lined by endothelial cells. There is a suggestion from the work of Tuder et al.^{23,24} that this is a form of neoplastic lesion reflecting dysregulation of endothelial cell growth, rather than a smooth muscle abnormality, the principal event in PPH. The plexiform lesions are monoclonal in PPH, and polyclonal in other forms of PAH.^{25,26} Plexiform lesions might also represent an angiogenic response to local ischemia or hypoxia.²⁷

Acute pulmonary hypertension induces a severe and sudden increase of right ventricular afterload, with an increased end-diastolic volume, a reduced ejection fraction, and a decreased stroke volume of the right ventricle.²⁸ However, chronic pulmonary hypertension leads to progressive systolic pressure overload of the right ventricle that dilates and hypertrophies, resulting in gradual right ventricular dysfunction.²⁹ In acute pulmonary hypertension, when pulmonary vascular resistance (PVR) increases abruptly, it is unusual for the right ventricle to be able to generate a mean pressure greater than 40 mmHg.³⁰⁻³² In the presence of more severe vascular obstruction, the heightened PVR induces a decrease in the cardiac index. In chronic pulmonary hypertension, increases in PVR gradually worsen right ventricular failure.³³ The ejection fraction of the right ventricle is also gradually reduced.

Decreased venous return compromises right ventricular preload and pulmonary blood flow. It may result from positive intrathoracic pressure during mechanical ventilation, such as positive end-expiratory pressure. High positive end-expiratory pressure also evokes alveolar overdistension and compression of the capillary network in the alveolar wall and interstitium. This condition increases PVR and reduces pulmonary blood flow.³⁴

In pulmonary hypertension, the increased PVR limits right ventricular stroke volume and the volume available for left ventricular filling. The left ventricle is also compressed as the intraventricular septum moves paradoxically to the left during systole. Left ventricular septal bowing reduces left ventricular volume in early diastole and can impair left ventricular filling in the most important phase of rapid filling.³⁵⁻³⁷ Both mechanisms lead to low cardiac output and reduced arterial pressure. Decreased systemic vascular resistance, which occurs after anesthetic administration, depresses systemic arterial pressure. Hypotension reduces coronary perfusion pressure, which can result in myocardial ischemia and cause right-sided heart failure. The right ventricle normally receives coronary blood flow in both systole and diastole. During pulmonary hypertension and augmented right ventricular wall stress, coronary blood flow to the right ventricle is dramatically decreased during systole if right ventricular systolic pressure is equal to or higher than systemic pressure, and could also become limited during diastole if right ventricular end-diastolic pressures are increased.

Hypoxemia can occur in conjunction with reduced cardiac output and pulmonary blood flow. It can also result from right-to-left intracardiac shunting. As many as 30% of adults have a patent foramen ovale.³⁸ Normally, there is no right-to-left shunting across the foramen ovale. If right atrial pressure exceeds left atrial pressure,

right-to-left shunting can occur, culminating in systemic oxygen desaturation. Finally, patients may have a restrictive pattern on pulmonary function tests or a low carbon monoxide diffusion capacity, leading to more severe hypoxemia and pulmonary hypertension.

Biology

In PAH, platelet activity is enhanced; serotonin, plasminogen activator inhibitor, and fibrinopeptide A concentrations are increased; and thrombomodulin is decreased.^{39,40} Thrombosis is often found in situ in pulmonary arterioles of PAH patients. The role of serotonin (5-hydroxytryptamine [5-HT]) in the development of PAH has been an enigma.⁴¹⁻⁴³ 5-HT promotes SMC proliferation, pulmonary arterial vasoconstriction, and local microthrombosis.⁴⁴ Alterations in 5-HT turnover, leading to an increased availability of free 5-HT in the vicinity of the pulmonary artery wall, have been proposed to be a pathophysiologic process.⁴⁵ The major source of stored 5-HT is the platelet-dense granule. There is evidence that alterations in platelet 5-HT storage and/or heightened platelet consumption by the lung may trigger the development of PAH. Decreased platelet 5-HT concentration with enhanced plasma concentrations of free 5-HT has been reported in many disorders associated with PAH, including anorexigen intake, portal hypertension,^{46,47} Raynaud phenomenon, and collagen vascular disease.⁴² In human pulmonary arteries, serotonin acts mainly via 5-HT $_{\rm 1B}$ and not via 5-HT $_{\rm 2A}$ receptors. $^{\rm 48,49}$ This explains the lack of effect of ketanserin, a 5-HT_{2A} receptor antagonist, on pulmonary hypertension in humans. Pulmonary vasoconstriction is mainly evoked through activation of the 5- HT_{1B} receptor, and smooth muscle proliferation is induced through activation of the 5-HT_{2A} receptor. 5-HT is taken up by SMC through the 5-HT transporter. The involvement of the 5-HT transporter in pulmonary hypertension and remodelling is based on several experimental observations⁵⁰⁻⁵³ and suggests that it plays a major role in hypoxia-induced vascular remodelling through its ability to mediate the mitogenic action of 5-HT.

Besides 5-HT and coagulation, several other mechanisms controlling pulmonary vascular tone and plasticity can be altered in pulmonary hypertension. The balance between endothelium-dependent vasodilators, such as prostacyclin and nitric oxide, and vasoconstrictors, such as ET-1 and thromboxane, is modified toward higher concentrations of vasoconstrictors and lower concentrations of vasodilators. Patients with either PPH or secondary pulmonary hypertension have increased excretion of thromboxane A_2 metabolites and reduced excretion of prostacyclin metabolites.⁵⁴ Endothelial nitric oxide synthase expression is reduced in the pulmonary circulation of patients with PPH compared to control subjects⁵⁵; however, this is still controversial.⁵⁶

Table 2. Possible Causes of Primary Pulmonary Hypertension

- 1. BMPR2 mutation and vascular smooth muscle cell proliferation
- 2. Monoclonal proliferation of endothelial cells in plexiform lesions
- Inhibition or down-regulation of the K_v channel in pulmonary artery SMC
- 4. Excess endothelial production of constrictors (ET-1, TXA₂) *versus* dilator mediators (nitric oxide, prostaglandin)
- 5. Serotonin excess
- 6. Thrombosis in situ

Endothelin-1 is a potent vasoconstrictor and mitogen. Its concentrations are increased in experimental pulmonary hypertension^{57,58} and in human PAH.⁵⁹⁻⁶⁴ ET-1 produced by pulmonary endothelial cells may contribute to increased PVR and to the pathogenesis of PPH.⁶⁵

K⁺ channels are transmembrane-spanning proteins that contain a pore with great selectivity for the K^+ ion. They are tonically active in vascular smooth muscle, allowing a slow efflux of K⁺ along their intracellular/ extracellular concentration gradient of 145/5 mm. There are several types of K⁺ channels; one of them, the voltage gated (Kv), has a voltage sensor and contributes to membrane potential in SMC. Inhibition of Kv channels results in accumulation of positively charged K⁺ ions within cells, raising the membrane potential to more positive levels (depolarization), which activates the voltage-gated L-type Ca^{2+} channel.⁶⁶ Ca^{2+} then enters the cells, activating their contractile apparatus, leading to vasoconstriction and possibly initiating cell proliferation. Inhibition of Kv could be one of the mechanisms of hypoxic pulmonary vasoconstriction.67,68 In humans with PPH, Kv1.5 mRNA concentrations are reduced in pulmonary artery SMC.⁶⁹ It is possible that decreased expression or function of K⁺ channels in pulmonary artery SMC of patients with PPH could initiate and/or maintain pulmonary hypertension and play a role in the pathogenesis of PPH.⁷⁰ It is intriguing that Kv2.1 is also inhibited by dexfenfluramine, a weight-loss drug that is associated with the development of PAH.

Vascular remodelling is a prominent feature of PPH. BMPR2, a member of the tumor growth factor (TGF)- β receptor family, regulates cell proliferation in response to ligand binding. The ligands for the TGF- β receptor family include TGF- β , bone morphogenetic protein, and activin. These growth factors have pleiotropic effects on endothelial cells and vascular SMCs depending on the context of the signal and the specific TGF receptor family members to which they bind.^{71,72} It has been postulated that mutations in BMPR2 in patients with PPH lead to loss of the inhibitory action of bone morphogenetic protein on the growth and the proliferative response of vascular endothelial and smooth cells in the pulmonary vasculature⁴ (table 2).

Primary pulmonary hypertension is also characterized

Anesthesiology, V 99, No 6, Dec 2003

Copyright © by the American Society of Anesthesiologists. Unauthorized reproduction of this article is prohibited.

by endothelial cell proliferation. Migration and proliferation of pulmonary endothelial cells and angiogenesis might be the initial phenomenon in the pathogenesis of PPH. This view is supported by the expression of BMPR2 on endothelial cells and plexiform lesions. Mutations in this receptor are likely to affect bone morphogenetic protein signaling in endothelial cells and myofibroblasts within obliteration lesions in PPH.⁷³

Moreover, the endothelial cell proliferation might be explained by mutations in activin receptor-like kinase (ALK)-1 found abundantly in the pulmonary vasculature. Normally, TGF- β binding ALK-1 attenuates the endothelial proliferative response induced by TGF-B binding to ALK-5.74 Trembath et al.75 have recently identified amino acid changes in ALK-1 in patients with pulmonary hypertension associated with hereditary hemorrhagic telangiectasia, which might result in unopposed TGF- β signaling through ALK-5 and endothelial cell proliferation. The role of vascular endothelial growth factor in the pathophysiology of PAH is controversial because the expression of vascular endothelial growth factor and its receptor are closely correlated with the formation of the plexiform lesion in human pulmonary hypertension,⁷⁶ and on the opposite, blockade of the vascular endothelial growth factor 2 receptor potentiates hypoxic pulmonary hypertension,77 and cell-based gene transfer of vascular endothelial growth factor attenuates experimental pulmonary hypertension.78

Pulmonary hypertension can be induced by different drugs and pathologic conditions. The first association between the anorexigen aminorex and pulmonary hypertension was observed in the 1960s.79,80 PAH was later associated with fenfluramine or dexfenfluramine intake, and it was subsequently shown that PAH occurs approximately 30 times more frequently in patients receiving these anorectic agents for more than 3 months compared to the general population.^{81,82} Fenfluramine releases 5-HT from platelets and inhibits its reuptake into platelets and pulmonary endothelial cells.^{83,84} Fenfluramine was widely coprescribed with phentermine, a combination that became known as "fen/phen." Phentermine can prolong the effect of 5-HT and increase the concentration of 5-HT by inhibiting monoamine oxidase B, which metabolizes 5-HT. The fen/phen association has been shown to induce valvular disease⁸⁵ similar to that observed after exposure to 5-HT-like drugs, such as ergotamine and methysergide, and to increased 5-HT concentrations found in carcinoid syndrome. Most of the anorexigens are also serotonin-transporter substrates⁸⁶ and thus get translocated into pulmonary vascular cells, where their effects may become amplified. Anorexigens are also Ky channel blockers,87 and one of their targets is Kv2.1.69,88 Anorexigen-induced Kv channel inhibition and membrane depolarization might contribute to pulmonary vasoconstriction.^{89,90} In addition to its effect on Ca^{2+} entry *via* the L-type Ca^{2+} channel, dexfenfluramine

Table 3. Signs of Disease Severity

- 1. Dyspnea at rest
- 2. Low cardiac output with metabolic acidosis

- 4. Signs of right heart failure (large V wave on jugularis vein,
- peripheral edema, hepatomegaly)
- 5. Syncope

also promotes vasoconstriction by enhancing Ca²⁺ release from the sarcoplasmic reticulum.^{88,91} Furthermore, fenfluramine reduces Kv1.5 mRNA concentrations by 50% in pulmonary artery SMC from normotensive patients,⁹² suggesting that inhibited gene transcription and expression of Kv channels may play an important role in anorexigen-induced PAH. In addition to a synergistic interaction between fenfluramine and phentermine favoring PAH and cardiac vascular disease,⁹³ fenfluramine may precipitate secondary forms of PAH. Fenfluramine and dexfenfluramine are no longer available for clinical use.

Symptoms and Evaluation

Careful diagnostic evaluation aimed at identifying various etiologies is essential for appropriate management of pulmonary hypertension. Symptoms are not specific, and the most frequent symptom is progressive dyspnea. Other common signs and symptoms include chest pain secondary to right ventricular ischemia, fatigue, peripheral edema, near syncope, and syncope. Syncope is a serious complication of pulmonary hypertension and portends a poor prognosis^{94,95} (table 3).

Clinical examination can help to detect pulmonary hypertension and right-sided heart failure. The signs of pulmonary hypertension depend on the severity of the disorder. A loud pulmonic component of the second heart sound is suggestive of increased PAP. Patients with right-sided heart overload may have a left parasternal heave.

A murmur of tricuspid regurgitation that may increase in intensity during inspiration can develop as the right ventricle dilates. Signs such as an increase in jugular venous pressure, neck veins pulsations (giant systolic V waves), peripheral edema, hepatomegaly, and ascites are indicative of right-sided heart failure. Dilatation of the pulmonary valve annulus produces the diastolic decrescendo murmur of pulmonary valve regurgitation, the Graham Steell murmur. Right ventricular S3 gallop is characteristic of advanced right ventricle failure and has a poor prognosis.⁹⁶

The diagnostic evaluation of patients with suspected pulmonary hypertension includes echocardiography, electrocardiography, chest radiographs, pulmonary function tests, ventilation/perfusion scanning, pulmonary angiography, spiral computed tomography, serologic testing, and liver function testing (table 4). Echocardiography is the screening method of choice.

^{3.} Hypoxemia

 Table 4. Recommended Complementary Examinations before

 Anesthesia in Patients with Pulmonary Hypertension

- 1. Electrocardiography
- 2. Chest radiology
- 3. Measurement of arterial blood gases
- Echocardiography: information obtained includes size of right heart (dilation or hypertrophy), tricuspid regurgitation, myocardial function, shift of intravenous septum, patency of foramen ovale, estimation of pulmonary pressure, left heart function
- Cardiac catheterization: information obtained includes pulmonary pressure, cardiac output, response to vasodilators, patency of foramen ovale, status of coronary circulation

Anatomic and functional data involving ventricular function, valvular abnormalities, and intracardiac shunts can be assessed. Echocardiography may show right ventricular hypertrophy, dilatation of the right heart chamber with impairment of left ventricular filling, and paradoxical motion of the interventricular septum. Doppler studies provide an estimate of pulmonary artery systolic pressure by measuring regurgitant flow across the tricuspid valve or by directly measuring systolic flow velocity across the pulmonary valve.96 Electrocardiographic abnormalities reflect right ventricular and right atrial enlargement as well as right ventricular hypertrophy. Chest radiography may show enlarged central and right and left pulmonary arteries.⁹⁴ Increased size of the cardiac silhouette may reflect an enlarged right ventricle and right atrium. Ventilation/perfusion scans, pulmonary angiograms, and spiral computed tomograms are useful to identify thromboembolic disease. Serologic tests may indicate collagen vascular diseases such as scleroderma, systemic lupus erythematosus, rheumatoid arthritis, HIV infection, liver diseases, and other rare conditions.¹

Cardiac catheterization remains the gold standard for the assessment of PAH. Right-sided heart catheterization confirms the presence of increased pressure (mean PAP > 25 mmHg), and the absence of pulmonary venous hypertension is suggested by normal mean capillary wedge pressure (< 15 mmHg). Central venous pressure is an important parameter to follow because it indicates the degree of right-sided heart failure. A low cardiac index is also of important prognostic significance. Hemodynamic abnormalities predict survival in patients with PPH.⁹⁷ Finally, right-sided heart catheterization is necessary for testing the efficacy of vasodilator drugs.

Vasodilator trials are performed using short-acting vasodilators such as nitric oxide, epoprostenol, or adenosine.⁹⁸⁻¹⁰⁰ The ideal response to these substances is pulmonary arterial vasodilatation with an increase in cardiac output, a decrease and a reduction of pulmonary resistance. Acute testing is only indicative of the acutely reversible component of pulmonary hypertension. As discussed above, some chronic therapies such as prostacyclin and nitric oxide may show benefit, even in the absence of an acute response.

Treatment Options

Until recently, the medical treatment of pulmonary hypertension was limited to anticoagulation, oxygen, and high-dose calcium channel blockers for responders, in association with diuretics and digoxin where indicated.

Thrombosis of small pulmonary arteries (in situ thrombosis) is seen in most patients who are dying of pulmonary hypertension. Evidence of thrombin activity and fibrinogen consumption has been found in patients with PPH.^{101,102} Anticoagulation therapy has been studied in uncontrolled case series, and, indeed, the long-term use of warfarin is associated with improved survival. Warfarin is administered in doses to maintain the international normalized ratio at 2-2.5 times the control level. The use of unfractionated or low-molecular-weight heparin has not been examined. Heparin might provide similar antithrombotic efficacy and potentially offer some benefit through inhibition of SMC proliferation. However, heparin has a synergistic effect on endothelial proliferation when combined with endothelial cell growth factor,¹⁰³ and, given the importance of endothelial proliferation in PAH, its application must be carefully examined. Oxygen therapy may be useful if arterial desaturation occurs at rest (hemoglobin saturation < 90%) or during physical activity.

High-dose calcium channel blockers were the first class of drugs that were shown to have dramatic, beneficial, long-term effects in selected patients with PPH.¹⁰⁴ The presumed mechanism is through vasodilatation and the subsequent decrease in mean PAP. Cardiac output may increase as a result of right ventricular afterload reduction. Patients tend to be responders or nonresponders to high doses of Ca²⁺ channel blockers, with only approximately 15-25% of patients responding. Only nifedipine and diltiazem have been tested rigorously. It is not known whether Ca²⁺ channel blockers that release nitric oxide, such as amlodipine, are more effective than other classes of medications. In patients who are responsive (defined as > 20% reduction of PVR and > 20%decrease in PAP), a dose-response relation seems to exist in terms of magnitude of action. High doses of Ca²⁺ channel blockers are necessary to achieve maximum benefit, and once achieved, the beneficial effect may be stable for many years. The indiscriminate use of calcium channel blockers in patients with PPH also has great potential for harm. There is no evidence that nonresponders benefit from them. Systemic hypotension producing reflex tachycardia, sympathetic stimulation, and right ventricular ischemia are detrimental effects of calcium channel blockers that ultimately may worsen survival. Acute vasodilation testing with nitric oxide and prostacyclin has been used as a technique to identify patients who may respond to high-dose calcium channel blockers. In a clinical trial, pulmonary and systemic effects of a nicardipine ($0.06 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) infusion and a changing fraction of inspired oxygen (Fio₂) was evaluated in patients with pulmonary hypertension secondary to chronic obstructive disease. Nicardipine reduces pulmonary resistance and increases cardiac index independent of Fio₂ without changing arterial and mixed venous content.¹⁰⁵ No controlled studies have examined the effect of calcium channel blockers in PAH secondary to causes other than PPH.

Nitric oxide used as an inhaled gas is a selective pulmonary vasodilator.¹⁰⁶ It acts by stimulating soluble guanylate cyclase and increasing cyclic guanosine monophosphate. The vasodilator response to inhaled nitric oxide is an index of the reversibility of pulmonary hypertension.¹⁰⁷ Long-term treatment of PAH with inhaled nitric oxide has been studied in some patients.¹⁰⁸ However, inhaled nitric oxide therapy is currently cumbersome and expensive and requires a fairly sophisticated system because it is administered only during the initial part of the inspiration phase.¹⁰⁹⁻¹¹¹ Patient mobility is limited by the need for a gas cylinder and an injector to provide nitric oxide. Because the effect of inhaled nitric oxide is mainly limited to the pulmonary circulation, there is no systemic vasodilatation and hypotension. The influence of inhaled nitric oxide on pulmonary remodeling has not definitively been established. If inhaled nitric oxide inhibits remodeling in newborn rats treated with monocrotaline,¹¹² inhaled nitric oxide has no effect on pulmonary remodeling in adults rats treated with monocrotaline.^{113,114} Nitric oxide could inhibit pulmonary vascular remodeling by two possibilities¹¹⁵: (1) a reduction in pressure/shear stress,² a direct antimitogenic effect on smooth muscles via inhibition of the expression/production of ET-1 and/or platelet-derived growth factor,¹¹⁶ or inhibition of serine elastase.¹¹⁷ The effect of inhaled nitric oxide may be potentiated by phosphodiesterase-5 inhibitors, which specifically suppress cyclic guanosine monophosphate catabolism and prolong its action. Two medications that inhibit phosphodiesterase-5 are available: dipyridamole and sildenafil.118-126 Sildenafil has already been described in case reports as a therapy for severe pulmonary hypertension.¹²⁷ Its effect might be explained by two mechanisms of action: increased cyclic guanosine monophosphate concentration and opening of K⁺ channels.¹²⁸

Prostaglandins such as epoprostenol sodium (the synthetic salt of prostacyclin), a U.S. Food and Drug Administration-approved treatment for PPH, have been studied extensively over the past decade in patients with PPH and secondary PAH. Prostacyclin increases cyclic adenosine monophosphate by stimulating adenylate cyclase, leading to vasodilatation. It increases cardiac output and heart rate and decreases mean PAP and right atrial pressure.^{129,130} Epoprostenol may have beneficial effects other than vasodilatation when given chronically, possibly because of its antiplatelet or antiproliferative properties.^{131,132} Lack of an acute vasodilator response does not preclude a positive effect of chronic treatment, and many patients with a poor acute response have greatly benefited from chronic therapy. In fact, epoprostenol is not intended for patients with an acute response to vasodilator therapy.¹³³ The epoprostenol dose must be increased gradually during the first year of therapy to prevent symptom recurrence. Its half-life in the blood is short (3-5 min), and it must be administered intravenously. Inhaled epoprostenol is experimental and not approved. Continuous epoprostenol infusion requires the placement of an indwelling venous catheter and is associated with the risk of infection. Epoprostenol therapy leads to a marked improvement in functional capacity and in the survival of patients with PPH,^{134,135} with improvement in functional capacity and hemodynamics in patients with PAH related to scleroderma-type diseases.¹³⁶ Despite the huge increase in price of inhaled nitric oxide treatment in some countries, epoprostenol treatment is more expensive than inhaled nitric oxide.¹⁰⁸ Intravenous use of prostaglandin (PG) I₂ is somewhat limited by its many adverse effects, including systemic hypotension, flushing, chest pain, headache, and diarrhea.

Treprostinil is a more stable analog of prostacyclin at room temperature with a longer half-life and can be administered subcutaneously. The Food and Drug Administration Advisory Committee has recommended its approval, and a recent trial has demonstrated that its long-term use improves exercise tolerance in patients with PPH.¹³⁷ Continuous infusion of trepostinil, another stable prostacyclin analog, has improved function in patients with PAH.¹³⁸ Iloprost is a prostacyclin analog that can be administered by inhalation. The major advantage of this inhalation strategy is that lower doses of the drug, with minimal systemic effects, can be used.^{139,140} Unfortunately, its short half-life requires frequent inhalation, and it is unclear whether the magnitude of longterm effects is sustained. Inhaled prostacyclin treatment could be combined with a phosphodiesterase-5 inhibitor such as sildenafil such that lower doses of drug can be used and the adverse effects can be minimized. 125,141-144

Wilkens *et al.*¹²⁰ conclude that sildenafil causes a longlasting reduction in mean PAP and PVR, with further additional improvement after inhalation of iloprost. These data suggest that small doses of a phosphodiesterase type V inhibitor may be a useful adjunct to inhaled iloprost in the management of pulmonary hypertension. Beraprost, an oral prostacyclin analog, is reasonably well absorbed and produces beneficial effects in patients with PPH.¹⁴⁵ Its efficacy is currently being examined in controlled clinical investigations.

The role of ET-1 as a mediator of PAH has been suggested.⁶⁵ Antagonists that are nonselective (block both endothelin-A [ET-A] and endothelin-B [ET-B] receptors) or are ET-A receptor selective have been developed and tested in animal models of pulmonary hypertension.¹⁴⁶ ET-A receptors on smooth muscle mediate vasoconstriction and promote SMC proliferation. ET-B receptors seem to be involved in ET-1 clearance, 147,148 and a subpopulation of ET-B receptors located in the endothelium mediates vasodilation through the release of nitric oxide and prostacyclin.¹⁴⁹ There may be an advantage to selectively block the ET-A receptor, leaving ET-B receptor function intact.¹³² Clinical trials with bosentan, a nonselective inhibitor, already suggest a beneficial effect in patients with PAH, 150-152 as do studies with sitaxsentan, an ET-A selective inhibitor. Bosentan has recently been approved by the Food and Drug Administration for the oral treatment of PPH.¹⁵³

It is likely that combination therapy using several medications acting through different and complementary pathways, for example, prostacyclin (intravenous or inhaled), inhaled nitric oxide, ET-A blockers, and phosphodiesterase inhibitors will be shown to be useful in the medical management of pulmonary hypertension.

Finally, the advances in our understanding of the pathogenesis of PAH will eventually lead to the development of novel approaches focusing directly on abnormal proliferation of endothelial cells¹⁵⁴ or regression of established pulmonary vascular remodelling.¹⁵⁵⁻¹⁵⁷ Genes for prostacyclin synthase and nitric oxide synthase have been transfected into the airways as well as small pulmonary arteries, and transiently overexpressed in mice.^{158,159} Gene therapy directed at the pulmonary bed, using nitric oxide synthase,¹⁶⁰ prostacyclin synthase,⁹ or perhaps BMPR2 is a possible future approach. Antiangiogenic therapy may also prove to be beneficial when given in the presence of endothelial proliferation.

Perioperative Care

The management of PPH or secondary PAH is a challenge for the anesthesiologist because the risk of rightsided heart failure is markedly increased. Knowledge of the pathophysiology of the disease and the therapeutic possibility of treating patients in the perioperative phase are essential.

The anesthetic management of patients with PAH is generally described in case reports of obstetric anesthesia and cardiac surgery.¹⁶¹⁻¹⁶³ In PAH, several mechanisms are implicated in right-sided heart failure, such as inadequate preload of the right ventricle, increased afterload of the right ventricle, hypotension, and hypoxemia. During anesthesia, all of these situations can occur. It is essential that the anesthesiologist avoid precipitating right-sided heart failure, resulting in low cardiac output.

Before anesthesia, chronic medical treatment already being administered for pulmonary hypertension and right-sided heart failure should be continued. If already being given, continuous intravenous epoprostenol therapy should be maintained at the same dose because it offers hemodynamic benefit and abrupt discontinuation of the drug can lead to syncope and death. Oxygen is useful in patients with hypoxemia (oxygen saturation < 90%) because hypoxemia causes pulmonary vasoconstriction and increases PVR.¹⁶⁴ Cardiac glycosides, such as digoxin, may improve cardiac output in patients with PPH and right ventricular failure and induce a significant reduction in circulating norepinephrine.¹⁶⁵ However, digitalis glycosides have a narrow therapeutic range. Digoxin blood concentration and blood electrolytes should be monitored as the risk of toxicity increases with hypokalemia. Diuretics should be used judiciously to control edema from right-sided heart dysfunction. Excessive diuresis may be extremely dangerous through a reduction of right ventricular preload. If overdiuresis occurs in the presence of acute right-sided heart failure secondary to acute myocardial infarction, it is less commonly an issue in the hypertrophied right side of the heart in chronic pulmonary hypertension. Fluid administration and induction of diuresis during anesthesia

should be based on careful hemodynamic monitoring, surveillance of filling pressure, cardiac chamber volume, intracardiac flow pattern (determined by echocardiography), and clinical response. Vasoactive medication should be added with caution. Sedatives should be administered with care, avoiding any

Sedatives should be administered with care, avoiding any drugs that potentially decrease systemic blood pressure.² The anesthesiologist must try to maintain optimal pulmonary blood flow. All factors that increase PVR should be avoided and corrected if they occur. Because hypoxia and hypercarbia augment PVR, ventilation and oxygenation should be controlled, and acidosis should be corrected. Changes in PVR in response to acidosis are small in the presence of normal alveolar oxygen tensions, but in the presence of alveolar hypoxia, they are greatly enhanced. Therefore, vasoconstriction may be augmented by increases in arterial hydrogen ion concentration, alveolar partial pressure of carbon dioxide, or both. Reducing arterial carbon dioxide tension ($Paco_2$) and increasing pH produce a consistent and reproducible decrease in PVR in infants with pulmonary hypertension.¹⁶⁶ Most of the sedative drugs and general anesthetic agents reduce systemic vascular resistance, which may lead, particularly in fixed cardiac output states, to decreased systemic arterial pressure and coronary perfusion.95

Calcium channel blockers given as treatment of chronic pulmonary hypertension should be continued despite any possible interaction with anesthetics on myocardium or vascular resistance.^{167,168} Particular attention should be paid to maintain sinus rhythm. Patients

BLAISE ET AL.

Table 5. Treatment of Pulmonary Hypertension during Surgery

Inhaled nitric oxide: 20-40 ppm. ^{226,227}
<i>Milrinone</i> (phosphodiesterase III inhibitor): $50 = \mu g/kg$ bolus
followed by a perfusion of 0.5–0.75 $\mu g \cdot kg^{-1} \cdot min^{-1}$
Dypiridamole: 0.2-0.6 mg/kg intravenously over 15 min; to be
repeated every 12 hours. ²²⁸⁻²³²
Inhaled prostacyclin: two modalities of application
1. Intermittent administration: 50 μ g is diluted in 50 ml saline
and nebulized in 15 min, which aerosolizes a dose between
14 and 17 μ g; this treatment must be repeated every
hour. ^{226,233–235}
2. Continuous administration at a concentration of
50 na \cdot ka ⁻¹ \cdot min ⁻¹ . ²³⁶
Prostacyclin, 1.5 mg, can be dissolved in 100 ml sterile
alveine buffer (final concentration, 15 μ g/ml); the drug is
administered by means of an inline nebulizer connected to
the inspiratory line. ²³⁷
If no nebulizing device is available, prostacyclin can be
infused intravenously at a dose between 4 and
$10 \text{ ng} \cdot \text{kg}^{-1} \cdot \text{min}^{-1.238}$
These medications should be weaned slowly after the
pulmonary hemodynamic response in the postoperative
period.
Regional anesthesia can be provided using nerve block,
brachial plexus block, and lumbar plexus block.
Epidural anesthesia should be induced slowly, and a mixture
of local anesthetics and opioids should be given to reduce
the dose of local anesthetics and then the hypotension.
Hypotension during the procedure should be treated
according to the etiology (bleeding etc.). Phenylephrine and
norepinephrine have been used to treat persistent systemic
hypertension; norepinephrine has the advantages of being
both a vasoconstrictor and a positive inotropic agent. This
medication should be titrated according to the clinical
response.
If regional anesthesia is used, arterial and CVP lines are
recommended, and specific treatment for pulmonary
hypertension is required.

CVP = central venous pressure; ppm = parts per million.

may deteriorate rapidly with the onset of atrial fibrillation or flutter because the atrial component to ventricular filling can be critical in maintaining cardiac output. High $Paco_2$ may induce arrhythmias during general anesthesia, especially in the presence of some anesthetics (*e.g.*, halothane). Increased $Paco_2$ with acidosis can increase PAP and right ventricular afterload.

Insertion of catheters or wires into the central circulation may provoke cardiac arrhythmia, so flow-directed thermodilution catheters should be introduced carefully. Difficulties may be encountered in passing the catheters into the pulmonary artery in patients with severe pulmonary hypertension.

If hypotension occurs despite euvolemia, inotropic agents are essential to improve cardiac output. Dobutamine is a β -adrenergic agent that increases cyclic adenosine monophosphate concentrations in both myocardium and vascular smooth muscle, resulting in inotropic, chronotropic, systemic, and pulmonary vasodilator effects. Dobutamine has been used extensively in patients with pulmonary hypertension after cardiopulmonary by-

pass (CPB) and for the evaluation of pulmonary vascular reactivity before cardiac transplantation. This agent decreases PVR in experimental pulmonary hypertension.¹⁶⁹ Phosphodiesterase-3 inhibitors, such as milrinone, reduce both systemic vascular resistance and PVR and augment contractility by increasing intracellular concentrations of cyclic adenosine monophosphate concentration.¹⁷⁰ They may be useful in supporting patients with pulmonary hypertension. Systemic hypotension is the limiting factor because of a lack of pulmonary specificity. Most of the medications used to treat pulmonary hypertension are vasodilators; their effects on systemic arterial pressure are difficult to predict because they are due to a balance among the decrease in pulmonary and systemic resistance, their impact on cardiac function, and oxygenation. As a consequence, therapy must be individualized to the patient's response.

In case of persistent hypotension, the use of vasoconstrictors such as phenylephrine and norepinephrine can augment coronary perfusion pressure and avoid right ventricular ischemia. Norepinephrine also provides inotropic support. Norepinephrine is metabolized by the pulmonary vascular endothelium. Therefore, its metabolism is reduced in PPH, and its concentrations increase.¹⁷¹⁻¹⁷³ Successful weaning for CPB in patients with acute pulmonary hypertension has been achieved with left atrial norepinephrine infusion combined with inhaled nitric oxide or PGE₁ infusion (table 6).^{174,175} Concomitant administration of inhaled nitric oxide or inhaled prostacyclin may be helpful.¹⁷⁶ Reductions in PVR and mean PAP with enhanced cardiac performance and minimal or no effects on systemic pressure have been reported. Nitric oxide and prostacyclin act via different signaling pathways, and their effects are additive without increased toxicity.177 Prostacyclin decreases PVR and augments cardiac output and systemic oxygen delivery when acutely administered to patients with PPH. The effect of inhaled nitric oxide and inhaled prostacyclin could be improved by phosphodiesterase inhibitors, such as dipyridamole and sildenafil.^{120,122}

Epidural anesthesia has been used successfully for cesarean delivery in patients with pulmonary hypertension.¹⁶³ The most significant risk in this situation is the reduction of venous return and arterial pressure consequent to sympathetic blockade. Also, the rapid changes in vascular volume and vascular tone peripartum pose a significant challenge. Few reports have described the effect of thoracic epidural anesthesia on PVR. Sympathetic innervation of the pulmonary circulation has been elaborated. No α_1 -adrenergic receptors are present in the pulmonary circulation.^{178,179} Therefore, sympathetic innervation does not contribute to basal vasomotor tone. Thoracic epidural has no impact on basal pulmonary artery tone. However, thoracic epidural anesthesia has cardiac effects that can be deleterious in patients with pulmonary hypertension. High thoracic epidural anes-

Table 6. Anesthetic Considerations

Preoperative medications	Agents
 Maintain all pulmonary vasodilators, such as intravenous or inhaled prostacyclin, Ca⁺⁺ antagonists, phosphodiesterase-5 inhibitors (sildenafil, dypiridamole), endothelin receptor antagonists (Bosentan), and oxygen. If pulmonary hypertension has been discovered in the immediate preoperative period and if the surgery cannot be delayed, a treatment with sildenafil (50–100 mg daily) and L-arginine (15 g daily) should be started as soon as possible. Heparin should replace indirect anticoagulant until the surgical procedure. Premedication: Slight sedation (midazolam) is allowed as long 	Few se action of pulmona been re pulmona ibe <i>et al</i> on the depende
as respiratory acidosis is not induced.	model.
 Opioids, such as fentanyl, alfentanil, sufentanil, and remifentanil, should be used at a dose to block the cardiorespiratory response of intubation. They have no direct vascular effect on pulmonary vessels. Lidocaine, 1 mg/kg, can also suppress the response to intubation. Propofol, 1–2 mg/kg; pentothal, 1–2 mg/kg; or etomidate, 0.2–0.4 mg/kg, may be used. Depolarizing or nondepolarizing muscle relaxants could be used. Maintenance 	attenuat tion. Pro vasodila sponse t ulate er but by d nary vas cyclin. A by nitric
Volatile anesthetics, such as isoflurane, desflurane, or sevoflurane, can be administered (isoflurane has been the most commonly used). Opioids should be maintained at a surgical analgesic level. Muscle relaxation should be maintained. Monitoring Arterial line: CVP or pulmonary artery catheter: TFF if available	could b Propofo induced ence gu smooth
Postoperative treatment Hospitalization in an intensive care unit Optimal analgesia with continuous epidural, regional block, or parenteral opioids	bradykii on the induced had no

CVP = central venous pressure; TEE = transesophageal echocardiography.

thesia, from the first to the fifth thoracic level, blocks cardiac afferent and efferent sympathetic fibers with loss of chronotropic and inotropic drive to the myocardium.¹⁸⁰ Heart rate may decrease as a result of blockade of the cardioaccelerator fibers arising from T1 to T4. The heart rate may decrease as a result of a decrease in right atrial filling, which reduces outflow from intrinsic chronotropic stretch receptors located in the right atrium.¹⁸¹

Postoperative Period

Patients with pulmonary hypertension who undergo surgery often die suddenly during the first postoperative days.¹⁸² Possible etiologies include a progressive increase in pulmonary vascular tone, acute pulmonary vasospasm, pulmonary thromboembolism, cardiac arrhythmia, heightened sympathetic tone, and fluid shifts. All precautions should be taken to avoid hypoxemia, hypotension, and hypovolemia. Postoperative control of pain should be effective. Any therapy to decrease PVR and improve pulmonary blood flow should be weaned with caution.

Pulmonary Vasoreactivity to Anesthetic Agents

studies have been performed on the intrinsic of anesthetic agents on the vasoreactivity of the ary vasculature. The effect of propofol on the ary circulation is controversial. Propofol has ported to cause either an increase in PVR or ary vasodilation in transplant patients.^{183,184} Hor-¹⁸⁵ have recently studied the effects of propofol pulmonary vascular response to endotheliument and -independent vasodilators in a canine They tested the hypothesis that propofol could e endothelium-dependent pulmonary vasodilataopofol reduces acetylcholine-induced pulmonary tation. However, it does not change the reto bradykinin. Acetylcholine and bradykinin stimndothelium-dependent pulmonary vasodilation, lifferent mechanisms. Bradykinin-induced pulmosodilation is mediated by nitric oxide and prosta-Acetylcholine-induced vasodilatation is mediated c oxide and a cytochrome P-450 metabolite that e endothelium-derived hyperpolarizing factor. ol has no effect on the pulmonary vasodilatation by nitric oxide, implying that it does not influanylyl cyclase activity in pulmonary vascular muscle. The normal response to nitric oxide and nin indicates that propofol has a selective effect endothelial signaling pathway for acetylcholinevasodilatation.¹⁸⁶ In a canine model, propofol effect on basal pulmonary vascular tone but increased the pulmonary response to vasoconstrictors.¹⁸⁷ It is not known whether the same response could occur in humans. Propofol has been used as an induction agent without problems in patients with PPH.188

Ketamine may produce sympathetic nervous system activation and increased concentrations of epinephrine and norepinephrine in plasma.¹⁸⁹ In vitro, ketamine increases PVR in rat lung.¹⁹⁰ However, a study performed using isolated rabbit pulmonary arteries showed a relaxant effect of ketamine, suggesting an intrinsic endothelium-independent vasodilator action on the pulmonary circulation.¹⁹¹ Relaxation caused by ketamine could be mediated by inhibition of calcium release from intracellular storage sites, or by a calcium channelblocking effect. In a rat thoracic aorta model, Miyawaki et al.192 found that ketamine inhibited acetylcholineinduced relaxation but not sodium nitroprusside-induced relaxation, implying that it suppressed nitric oxide formation in the endothelium. Ogawa et al.¹⁹³ have shown recently in canine pulmonary artery preparations that ketamine attenuates endothelium-dependent pulmonary vasorelaxation in response to acetylcholine and bradykinin by inhibiting both the nitric oxide and the endothelium-derived hyperpolarizing factor components

Cause of Pulmonary Hypertension and Surgical Intervention	Anesthesia Technique and Medication Used	Outcome
Eisenmenger syndrome ²³⁹	103 Anesthesia 68 General anesthesia 19 Neuroaxial anesthesia 16 Combined anesthesia	Mortality 14% Mortality 18% Mortality 5% Mortality 7%
Major surgery: cesarean delivery, hysterectomy, laparotomy, vascular surgery		Mortality 24%
Minor surgery: hernia repair, tubal ligation, dental work, dilatation and curettage, extremity surgery		Mortality 4%
PPH Cesarean delivery ²²⁵	EDA 18 mg ropivacaine + 5 mg morphine	Hypotension Bradycardia 0.25 mg atropine 10 mg ephedrine
	Postoperative analgesia	1.7 mg phenylephrine
	Epidural morphine Oral calcium channel blockers and low- molecular-weight heparin	Immediate outcome good Patient died 19 months later, waiting for heart-lung transplantation
PPH	General anesthesia	Acute pulmonary hypertension after intubation, low CO (1.2 l/min)
Lung transplantion ²⁴⁰	250 μg fentanyl 3.6 mg midazolam 70 mg succinylcholine 0.5–1% isoflurane	
	TEE Dilatation of RV Major T regurgitation Underfilling of LV Shift of IV septum	Treatment Epinephrine $2 \times 50 \ \mu g + perfusion$ $0.01 \ \mu g \cdot kg^{-1} \cdot min^{-1}$ Cardiac arrest after thoracotomy, open
		chest cardiac massage, catheterization of femoral vessels Beginning of CPB Good final outcome
PPH	FDA	Good outcome
Cesarean delivery ¹⁶³	300 mg lidocaine, 1% 70 mg bupivacaine, 0.5% 0.1 mg fentanyl 20 mg aerosolized iloprost Eurosemide for positive volume into	
	postoperative period	
PPH	General anesthesia	
Lung transplantation-	Etomidate Rocuronium	Carolac arrest Carolac massage Canulation of femoral vessels CPB
	Nitroghyporing	Good outcome
PPH Cesarean delivery ²¹⁹	Nitroglycerine 50–100 mg/h <i>via</i> PA catheter to reduce PAP 1 mg alfentanil	Good outcome
	16 mg etomidate 100 mg succinylcholine Isoflurane	
	$50\%/50\% N_2/O_2$ After delivery 5 U oxytocin Nitroglycerin replaced by 2–5 ng , kg ⁻¹ , min ⁻¹	
	intravenous prostacyclin Because of vaginal bleeding	
	60,000 ng/h Finally, 30 mg oral nefedipin and weaning from	
	prostacyclin	

Table 7. Summary of Recent Case Reports of Perioperative Management of Pulmonary Hypertension

(Table continues)

Anesthesiology, V 99, No 6, Dec 2003

Table 7. (continued)

Cause of Pulmonary Hypertension and Surgical Intervention	Anesthesia Technique and Medication Used	Outcome
Severe pulmonary hypertension due to valvular heart disease ¹⁸⁸	General anesthesia: TCI	Good outcome; valves repaired 1 month after surgery
(mitral stenosis and regurgitation,	8 ng/ml remifentanil	
Coorresp delivery	$2.5 \ \mu g/m \ proposol$	
Cesarean delivery	Postsurgical pain management	
	3-ml/h infusion of a mixture of 0.06%	
	bupivacaine	
1 μg/ml s	1 μ g/ml sulfentanil	-
Primary and secondary hypertension ²⁴¹	Epidural anesthesia	Good outcome
Vaginal delivery: seven patients	General anesthesia	
Cesarean delivery: one patient	200 mg thiopenthal	
	100 mg succinylcholine	
	100 μ g fentanyl 0.5% balothano	
	50%/50% N ₂ O/O ₂	Death 12 h after admission to ICU
PPH	Epidural anesthesia	
Cesarean delivery ¹⁶³	300 mg lidocaine, 2%	
	70 mg bupivacaine, 0.5%	
	100 μ g fentanyl	
	$20 \ \mu g$ aerosolized lioprost Perfusion of 2.400 ml Ringer's lactate	
	Postoperative treatment	
	lloprost inhalation 6 times/day	
	Furosemide to treat 1,600 positive balance	
	Analgesia	
	Intravenous paracetamol and morphine	Good outcome

CO = cardiac output; CPB = cardiopulmonary bypass; EDA = epidural analgesia; ICU = intensive care unit; IV = intravenous; LV = left ventricle; N_2O/O_2 = nitrous oxideoxygen ratio; PA = pulmonary artery; PAP = pulmonary artery pressure; PPH = primary pulmonary hypertension; RV = right ventricle; TEE = transesophageal echocardiography. TCI = target-controlled infusion.

of the response.¹⁹⁴ These results reveal the differential reactivity of the pulmonary vascular bed and that the clinical effect of an agent is always mediated *via* the integration of several systems. Like ketamine, etomidate selectively attenuates endothelium-dependent canine pulmonary vasorelaxation by inhibiting the nitric oxide and endothelium-derived hyperpolarizing factor components of the response. It seems likely that etomidate and ketamine suppress the pulmonary vasorelaxant response to acetylcholine and bradykinin by reducing the increase in endothelial Ca²⁺, an essential step in the production of nitric oxide and endothelium-derived hyperpolarizing factor in response to receptor activation.¹⁹³

Inhalational Anesthetic Agents

Isoflurane anesthesia exerts differential effects on the vasoreactivity of the pulmonary circulation. It attenuates the magnitude of hypoxic pulmonary vasoconstriction. Isoflurane potentiates the vasodilator response to β_1 adrenoceptor activation. However, it has no effect on the vasoconstriction response to α_1 adrenoreceptor activation.^{195,196}

Adenosine triphosphate-sensitive potassium channels play an important role in the regulation of vascular smooth muscle tone.¹⁹⁷ Adenosine triphosphate-sensitive potassium channel activation mediates the vasodilator effect of many endogenous mediators, such as adenosine, PGI₂, and nitric oxide. Isoflurane inhibits endothelium-dependent relaxation in aorta as well as in isolated pulmonary arteries. It selectively attenuates the pulmonary vasorelaxant response by inhibiting the activity of the adenosine triphosphate-sensitive potassium channels, which regulate the synergy between nitric oxide and PGI₂. It has no effect on baseline pulmonary circulation tone.¹⁹⁸ Conversely, isoflurane suppresses the pulmonary vasoconstrictor response to hypotension. In this situation, it does not modify the influence of endogenous potassium-channel activation on the pulmonary vascular response to hypotension. Possibly, isoflurane may attenuate the pulmonary vasoconstrictor response to hypotension through a differential effect on pulmonary sympathetic adrenoceptors in the vasculature.199

Neither halothane nor enflurane exerts any action on the baseline pulmonary circulation, but these agents reduce the pulmonary vasodilator effect mediated by

Anesthesiology, V 99, No 6, Dec 2003

adenosine triphosphate-sensitive potassium channel agonists.²⁰⁰ Desflurane shares this latter feature, unlike sevoflurane, which has no effect. Desflurane potentiates pulmonary vasoconstriction in response to adrenoceptor activation, in contrast to isoflurane.²⁰¹

Sykes et al.²⁰² demonstrated in the dog lung that nitrous oxide seemed to attenuate hypoxic pulmonary vasoconstriction. Other investigators have obtained little or no action when nitrous oxide is added to inhaled gas mixtures.²⁰³ Nitrous oxide has minimal effects on hemodynamics but can still depress myocardial contractility.²⁰⁴ In infants, nitrous oxide does not produce the increases in PAP and PVR observed in adults.²⁰⁵ Nitrous oxide should be used with caution in adult patients with increased PVR, particularly in the presence of right ventricular dysfunction.²⁰⁶ In a dog model, nitrous oxide had no effect on pulmonary vascular tone in an isolated lobe but increased PVR in intact animals.²⁰⁷ Despite a moderate impact on pulmonary pressure, nitrous oxide has been used several times without detrimental results in patients with pulmonary hypertension. 206, 208, 209

Anesthesia for Special Procedures

Immediate Preoperative Preparation

All medications specially prescribed for treating pulmonary hypertension, such as prostacyclin, Ca^{2+} antagonists, phosphodiesterase inhibitors, ET receptor antagonists, and oxygen therapy, should be continued until the procedure and after surgery. Coumadin should be changed to heparin before the procedure.

If the patient has had no specific treatment for pulmonary hypertension and if the surgery cannot be delayed to establish the best treatment, phosphodiesterase inhibitors (50–100 mg sildenafil) should be given daily, with L-arginine supplementation (15 g/day). This is especially indicated if the patient has clinical signs of pulmonary hypertension and demonstrates poor exercise tolerance. For premedication, a sedative, such as benzodiazepine, is administered to decrease anxiety.

Anesthesia for Bilateral Lung Transplantation

Induction of anesthesia can be accomplished with opioids (5-10 μ g/kg fentanyl or 0.5-1 μ g/kg sufentanil), 1 mg/kg lidocaine, 2-3 mg/kg pentothal, or 0.5-1 mg/kg propofol with muscle relaxants. Muscles relaxants releasing histamine such as atracurium or cisatracurium should be avoided. Mask ventilation with 100% O₂ and a volatile anesthetic such as isoflurane could be used to deepen the level of anesthesia before intubation. If necessary, norepinephrine should be infused to treat hypotension evoked by induction agents. If the patient is not fastened (this situation is not frequent because it takes some time to organize the transplantation), rapid-sequence induction can be used after hyperventilation to reduce $Paco_2$ and increase $Paco_2$. However, in this alternative, the induction is not as smooth, and it is more difficult to control the pulmonary and systemic hemodynamics.

Mechanical ventilation should allow enough time to let the lung deflate and avoid thoracic hyperinflation with reduction of venous return.²¹⁰ Anesthesia maintenance is based on opioids, volatile agents, and muscle relaxants. Total intravenous anesthesia could also be used. Specific pulmonary vasodilators, such as inhaled nitric oxide and inhaled prostacyclin, should be immediately available in the operating room.

The most difficult part of the anesthesia usually occurs during one-lung ventilation of the native lung. Acute exacerbation of pulmonary hypertension can occur because total cardiac output is going only to one lung. A pulmonary vasodilator, such as inhaled nitric oxide, may be useful. However, CPB is often needed to complete the procedure. Inhaled nitric oxide can be administered before reperfusion because it decreases reperfusion injury.^{211–213} After the procedure, there is usually no problem of pulmonary hypertension, and if patients were on PG infusion before surgery, it can be weaned rapidly after transplantation. The management then becomes similar to that for other major thoracic procedures. Epidural thoracic analgesia is indicated as soon as coagulation status is in the normal range.

Anesthesia for Cardiac Procedures

The use of CPB induces a pulmonary inflammatory response (acute lung injury) marked by pulmonary hypertension and hypoxemia. The inflammatory response can be prevented by inhaled nitric oxide and nitric oxide administered through the CPB machine; 20 parts per million (ppm) NO should be given throughout the procedure.^{214–216} Phosphodiesterase-3 inhibitors, such as milrinone, are helpful to wean from CPB, and a bolus can be added to the CPB reservoir before weaning. The dosage for milrinone is a 25- to 50- μ g/kg bolus, followed by infusion at 0.375–0.750 μ g · kg⁻¹ · min⁻¹. Hypotension and low cardiac output should be treated with norepinephrine.

If pulmonary pressure remains high despite treatment with inhaled nitric oxide, right ventricular failure may occur. The latter can be diagnosed with use of direct myocardial vision, measurement of pulmonary hemodynamics, and transesophageal echocardiography and can be treated with a phosphodiesterase-5 inhibitor (dipyridamole). Finally, inhaled prostacyclin should be given if the previous treatment has not sufficiently decreased pulmonary pressure. Weaning from inhaled nitric oxide should be performed progressively while following the PAP response. The process can be facilitated by phosphodiesterase-5 inhibitors or L-arginine (table 5).

Anesthesia for Peripheral and General Surgery

Peripheral surgery can be completed under regional block (e.g., brachial plexus, lumbar plexus nerve block). Epidural anesthesia can be used. Prostacyclin and nitric oxide have a synergistic inhibitory effect on platelet function,²¹⁷ and prostacyclin inhibits platelet aggregation in vitro.218 In patients with PPH, platelet aggregation was increased because of a higher concentration of thromboxane A₂ and 5-HT, and this abnormality was corrected in 80% of patients treated with continuous intravenous prostacyclin.131 In one patient, vaginal bleeding after delivery was enhanced after treatment by intravenous prostacyclin. The bleeding stopped after changing the route of administration of the drug to inhalation from intravenous.²¹⁹ Inhaled prostacyclin has no effect on platelet function and on bleeding after cardiac surgery.²²⁰ There are no data on the safety of epidural anesthesia in patients treated with intravenous prostacyclin. The reduction of vaginal bleeding after changing the route of prostacyclin administration (intravenous vs. inhaled) could be due to an effect on platelet function or a change in uterine tone. There are no reports that long-term intravenous infusion of prostacyclin produces epidural bleeding after regional anesthesia. If inhaled, prostacyclin treatment is not a contraindication to epidural anesthesia. Platelet function testing, if easily available, would give useful information before proceeding with epidural anesthesia if the patients are treated with intravenous prostacyclin. The induction of epidural anesthesia should be progressive to avoid sudden hypotension, and morphine or morphine derivatives should be added to local anesthetics to improve analgesia and reduce the amount of local anesthetics as well as the hemodynamic consequences of regional block. Spinal anesthesia with local anesthetics is not the technique of choice because the hemodynamic changes at induction and during recession of the block can be rapid and poorly tolerated. Systemic hypotension may be treated with either phenylephrine or norepinephrine. Norepinephrine acting primarily through B1-adrenergic receptors has a positive inotropic effect on the myocardium.^{171,172,221} A central intravenous line or pulmonary artery catheter and an arterial cannula are useful to monitor hemodynamics and to administer medications.

If general anesthesia is required, induction can be completed using a mixture of opioids, hypnotics, and volatile anesthetics, as previously explained, and maintenance comprising a mixture of volatile anesthetics, opioids, and muscle relaxants. A combination of epidural and general anesthesia is a good choice, and the epidural can also be used for postoperative analgesia. Inhaled nitric oxide should be available and given (concentration 20 - 40 ppm) to treat pulmonary hypertension crises. If this treatment is inefficient, dipyridamole should be administered to potentiate inhaled nitric oxide. Finally, aerosolized prostacyclin could add its effect to the previously mentioned medications to decrease pulmonary pressure.

Patients should be admitted to the intensive care unit in the postoperative period and monitored closely. Pain management should be optimal, and inhaled nitric oxide, if used during the procedure, should be weaned progressively.

Anesthesia in the Obstetrics Population

Epidural Anesthesia for Delivery

Several hemodynamic objectives should be reached: maintain the pulmonary pressure as low as possible and the systemic pressure within the 15% above and below the basal level (the systemic pressure should always be higher than pulmonary pressure), avoid dysrhythmias and tachycardia, and maintain sinusal rhythm. An arterial line and a central venous or pulmonary catheter should be used for monitoring or for drug administration. Patients should be admitted to the intensive care unit after delivery.

Epidural Anesthesia for Vaginal Delivery

Pain induced by labor should be treated in these patients with epidural analgesia *via* a mixture of local anesthetics and opioids in a low concentration because it produces good analgesia, keeps sufficient muscle tone, and has a minor effect on blood pressure. Forceps delivery, which decreases patient effort and has hemodynamic consequences, is the technique of choice.²²²⁻²²⁴

Cesarean Delivery

Both general and epidural anesthesia have been used for cesarean delivery in patients with pulmonary hypertension.^{163,188,219,225} Special care should be taken to avoid perioperative hemodynamic instability. Patient positioning is important.

The surgical procedure can lead to excessive bleeding and hypovolemia. Uterine contraction after delivery may return a large bolus of blood to the circulation. This can be poorly tolerated in patients with severe pulmonary hypertension and mitral stenosis. The sudden hypervolemia can be treated with vasodilators, such as nitroglycerine, and diuretics.

A blood pressure cuff inflated between the arterial and venous pressures around the thighs, can suddenly and reversibly decrease right ventricular filling by reducing venous return. Air or amniotic fluid embolism could acutely increase pulmonary pressure.

Induction of general anesthesia is based on opioids: fentanyl, sufentanil, or remifentanil. Lidocaine (1 mg/kg) reduces pulmonary and hemodynamic reactions during intubation. Induction can be further achieved by pentothal, propofol, or etomidate. Succinylcholine can be used for intubation. Anesthesia is maintained with use of sufentanil or remifentanil infusion (if these opioids have been used for induction), volatile anesthetics such as isoflurane, or propofol infusion.

If long-acting opioids have been used (fentanyl or sufentanil), the resuscitation team must be prepared to intubate and ventilate the newborn, who will have depressed respiration; naloxone has been administered in some cases (a special team should be ready to take care of the newborn). Postoperative analgesia may be achieved by intravenous, epidural, or intrathecal opioids. If the epidural approach is taken, a mixture of opioids and low-dose bupivacaine can be infused continuously (table 7).

Summary

Pulmonary arterial hypertension is a serious clinical problem associated with significant morbidity and mortality. A better understanding of disease pathophysiology will contribute to the development of new therapies and improve the long-term prognosis of patients. Selective pulmonary vasodilators are available and can be given separately or in combination. Hemodynamic monitoring and transesophageal echocardiography are diagnostic tools that are essential for the diagnosis but also give direct information on the efficacy of the therapeutic used.

References

Rubin LJ: Primary pulmonary hypertension. N Engl J Med 1997; 336:111-7
 Gaine S: Pulmonary hypertension. JAMA 2000; 284:3160-8

3. Loyd JE, Butler MG, Foroud TM, Conneally PM, Phillips JA III, Newman JH: Genetic anticipation and abnormal gender ratio at birth in familial primary pulmonary hypertension. Am J Respir Crit Care Med 1995; 152:93-7

4. Liu F, Ventura F, Doody J, Massague J: Human type II receptor for bone morphogenic proteins (BMPs): Extension of the two-kinase receptor model to the BMPs. Mol Cell Biol 1995; 15:3479-86

5. Massague J, Chen YG: Controlling TGF-beta signaling. Genes Dev 2000; 14:627-44

6. Thomson JR, Machado RD, Pauciulo MW, Morgan NV, Humbert M, Elliott GC, Ward K, Yacoub M, Mikhail G, Rogers P, Newman J, Wheeler L, Higenbottam T, Gibbs JS, Egan J, Crozier A, Peacock A, Allcock R, Corris P, Loyd JE, Trembath RC, Nichols WC: Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-beta family. J Med Genet 2000; 37:741-5

7. Deng Z, Morse JH, Slager SL, Cuervo N, Moore KJ, Venetos G, Kalachikov S, Cayanis E, Fischer SG, Barst RJ, Hodge SE, Knowles JA: Familial primary pulmonary hypertension (gene PPH1) is caused by mutations in the bone morphogenetic protein receptor-II gene. Am J Hum Genet 2000; 67:737-44

8. Machado RD, Pauciulo MW, Thomson JR, Lane KB, Morgan NV, Wheeler L, Phillips JA III, Newman J, Williams D, Galie N, Manes A, McNeil K, Yacoub M, Mikhail G, Rogers P, Corris P, Humbert M, Donnai D, Martensson G, Tranebjaerg L, Loyd JE, Trembath RC, Nichols WC: BMPR2 haploinsufficiency as the inherited molecular mechanism for primary pulmonary hypertension. Am J Hum Genet 2001; 68:92-102

9. Geraci MW, Hoshikawa Y, Yeager M, Golpon H, Gesell T, Tuder RM, Voelkel NF: Gene expression profiles in pulmonary hypertension. Chest 2002; 121:1048-58

10. Grunig E, Mereles D, Arnold K, Benz A, Olschewski H, Miltenberger-Miltenyi G, Borst MM, Abushi A, Seeger W, Winkler J, Hoper MM, Bartram CR, Kubler W, Janssen B: Primary pulmonary hypertension is predominantly a hereditary disease. Chest 2002; 121:81S-28

11. Meyrick B, Reid L: Pulmonary hypertension: Anatomic and physiologic correlates. Clin Chest Med 1983; 4:199-217

12. McMurtry IF, Hookway BW, Roos S: Red blood cells play a crucial role in maintaining vascular reactivity to hypoxia in isolated rat lungs. Chest 1977; 71:253-6

13. Barer G, Emery C, Stewart A, Bee D, Howard P: Endothelial control of the

pulmonary circulation in normal and chronically hypoxic rats. J Physiol 1993; 463:1-16

14. MacLean MR, McCulloch KM, Baird M: Endothelin ETA- and ETB-receptormediated vasoconstriction in rat pulmonary arteries and arterioles. J Cardiovasc Pharmacol 1994; 23:838-45

15. McCulloch KM, Docherty CC, Morecroft I, MacLean MR: EndothelinB receptor-mediated contraction in human pulmonary resistance arteries. Br J Pharmacol 1996; 119:1125-30

16. Troncy E, Francoeur M, Salazkin I, Yang F, Charbonneau M, Leclerc G, Vinay P, Blaise G: Extra-pulmonary effects of inhaled nitric oxide in swine with and without phenylephrine. Br J Anaesth 1997; 79:631-40

17. Hebb C: Motor innervation of the pulmonary blood vessels of mammals, The Pulmonary Circulation and Interstitial Space. Edited by Fishman AP, Hecht HH. Chicago, University of Chicago Press, 1969, pp 195-217

18. Berkowitz DE, Price DT, Bello EA, Page SO, Schwinn DA: Localization of messenger RNA for three distinct alpha 2-adrenergic receptor subtypes in human tissues: Evidence for species heterogeneity and implications for human pharma-cology ANESTHESIOLOGY 1994; 81:1235-44

19. Heath D, Smith P, Gosney J, Mulcahy D, Fox K, Yacoub M, Harris P: The pathology of the early and late stages of primary pulmonary hypertension. Br Heart J 1987; 58:204-13

20. Frid MG, Moiseeva EP, Stenmark KR: Multiple phenotypically distinct smooth muscle cell populations exist in the adult and developing bovine pulmonary arterial media in vivo. Circ Res 1994; 75:669-81

21. Michelakis ED, Reeve HL, Huang JM, Tolarova S, Nelson DP, Weir EK, Archer SL: Potassium channel diversity in vascular smooth muscle cells. Can J Physiol Pharmacol 1997; 75:889-97

22. Tchekneva E, Lawrence ML, Meyrick B: Cell-specific differences in ET-1 system in adjacent layers of main pulmonary artery: A new source of ET-1. Am J Physiol Lung Cell Mol Physiol 2000; 278:L813-L821

23. Tuder RM, Chacon M, Alger L, Wang J, Taraseviciene-Stewart L, Kasahara Y, Cool CD, Bishop AE, Geraci M, Semenza GL, Yacoub M, Polak JM, Voelkel NF: Expression of angiogenesis-related molecules in plexiform lesions in severe pulmonary hypertension: Evidence for a process of disordered angiogenesis. J Pathol 2001; 195:367-74

24. Tuder RM, Voelkel NF: Plexiform lesion in severe pulmonary hypertension: Association with glomeruloid lesion. Am J Pathol 2001; 159:382-3

25. Voelkel NF, Cool C, Lee SD, Wright L, Geraci MW, Tuder RM: Primary pulmonary hypertension between inflammation and cancer. Chest 1998; 114: 225S-30S

26. Lee SD, Shroyer KR, Markham NE, Cool CD, Voelkel NF, Tuder RM: Monoclonal endothelial cell proliferation is present in primary but not secondary pulmonary hypertension. J Clin Invest 1998; 101:927-34

27. Cool CD, Stewart JS, Werahera P, Miller GJ, Williams RL, Voelkel NF, Tuder RM: Three-dimensional reconstruction of pulmonary arteries in plexiform pulmonary hypertension using cell-specific markers: Evidence for a dynamic and heterogeneous process of pulmonary endothelial cell growth. Am J Pathol 1999; 155:411-9

28. Mercat A, Meyer G: Embolie pulmonaire grave, Insuffisance circulatoire aiguë. Edited by Richard C, Vincent JL. Paris, Arnette, 1994, pp 313-29

29. Morpurgo M: Le coeur dans l'hypertension arterielle pulmonaire, Hypertension arterielle pulmonaire. Edited by Weitzenblum E, Denolin H. Paris, Masson, 1995, pp 117-27

30. Benotti JR, Dalen JE: The natural history of pulmonary embolism. Clin Chest Med 1984; 5:403-10

31. Dalen JE, Banas JS Jr, Brooks HL, Evans GL, Paraskos JA, Dexter L: Resolution rate of acute pulmonary embolism in man. N Engl J Med 1969; 280:1194–9

32. McIntyre KM, Sasahara AA: The hemodynamic response to pulmonary embolism in patients without prior cardiopulmonary disease. Am J Cardiol 1971; 28:288-94

33. Wiedemann HP, Matthay RA: The management of acute and chronic cor pulmonale, Heart-Lung Interactions in Health and Disease. Edited by Scarf SM, Cassidy SS. New-York, Marcel Dekker, 1991, pp 915-82

34. Jenkins J, Lynn A, Edmonds J, Barker G: Effects of mechanical ventilation on cardiopulmonary function in children after open-heart surgery. Crit Care Med 1985; 13:77–80

35. Boxt LM, Katz J, Kolb T, Czegledy FP, Barst RJ: Direct quantitation of right and left ventricular volumes with nuclear magnetic resonance imaging in patients with primary pulmonary hypertension. J Am Coll Cardiol 1992; 19:1508-15

36. Louie EK, Rich S, Brundage BH: Doppler echocardiographic assessment of impaired left ventricular filling in patients with right ventricular pressure overload due to primary pulmonary hypertension. J Am Coll Cardiol 1986; 8:1298 -306

37. Marcus JT, Vonk NA, Roeleveld RJ, Postmus PE, Heethaar RM, Van Rossum AC, Boonstra A: Impaired left ventricular filling due to right ventricular pressure overload in primary pulmonary hypertension: Noninvasive monitoring using MRI. Chest 2001; 119:1761-5

38. Hagen PT, Scholz DG, Edwards WD: Incidence and size of patent foramen ovale during the first 10 decades of life: An autopsy study of 965 normal hearts. Mayo Clin Proc 1984; 59:17-20

39. Frank H, Mlczoch J, Huber K, Schuster E, Gurtner HP, Kneussl M: The effect of anticoagulant therapy in primary and anorectic drug-induced pulmonary hypertension. Chest 1997; 112:714-21

40. Welsh CH, Hassell KL, Badesch DB, Kressin DC, Marlar RA: Coagulation and fibrinolytic profiles in patients with severe pulmonary hypertension. Chest 1996; 110:710-7

41. Egermayer P, Town GI, Peacock AJ: Role of serotonin in the pathogenesis of acute and chronic pulmonary hypertension. Thorax 1999; 54:161-8

42. Herve P, Launay JM, Scrobohaci ML, Brenot F, Simonneau G, Petitpretz P, Poubeau P, Cerrina J, Duroux P, Drouet L: Increased plasma serotonin in primary pulmonary hypertension. Am J Med 1995; 99:249-54

43. MacLean MR: Endothelin-1 and serotonin: Mediators of primary and secondary pulmonary hypertension? J Lab Clin Med 1999; 134:105-14

44. Fanburg BL, Lee SL: A new role for an old molecule: Serotonin as a mitogen. Am J Physiol 1997; 272:L795-L806

45. MacLean MR, Herve P, Eddahibi S, Adnot S: 5-hydroxytryptamine and the pulmonary circulation: Receptors, transporters and relevance to pulmonary arterial hypertension. Br J Pharmacol 2000; 131:161-8

46. Laffi G, Marra F, Gresele P, Romagnoli P, Palermo A, Bartolini O, Simoni A, Orlandi L, Selli ML, Nenci GG: Evidence for a storage pool defect in platelets from cirrhotic patients with defective aggregation. Gastroenterology 1992; 103:641-6

47. Liu H, Lee SS: Cardiopulmonary dysfunction in cirrhosis. J Gastroenterol Hepatol 1999; 14:600-8

48. MacLean MR, Clayton RA, Templeton AG, Morecroft I: Evidence for 5-HT1like receptor-mediated vasoconstriction in human pulmonary artery. Br J Pharmacol 1996; 119:277-82

49. Morecroft I, Heeley RP, Prentice HM, Kirk A, MacLean MR: 5-hydroxytryp-tamine receptors mediating contraction in human small muscular pulmonary arteries: Importance of the 5-HT1B receptor. Br J Pharmacol 1999; 128:730-4

50. Eddahibi S, Fabre V, Boni C, Martres MP, Raffestin B, Hamon M, Adnot S: Induction of serotonin transporter by hypoxia in pulmonary vascular smooth muscle cells: Relationship with the mitogenic action of serotonin. Circ Res 1999; 84;329-36

51. Eddahibi S, Raffestin B, Pham I, Launay JM, Aegerter P, Sitbon M, Adnot S: Treatment with 5-HT potentiates development of pulmonary hypertension in chronically hypoxic rats. Am J Physiol 1997; 272:H1173-81

52. Eddahibi S, Hanoun N, Lanfumey L, Lesch KP, Raffestin B, Hamon M, Adnot S: Attenuated hypoxic pulmonary hypertension in mice lacking the 5-hydroxytryptamine transporter gene. J Clin Invest 2000; 105:1555-62

53. Eddahibi S, Humbert M, Fadel E, Raffestin B, Darmon M, Capron F, Simonneau G, Dartevelle P, Hamon M, Adnot S: Hyperplasia of pulmonary artery smooth muscle cells is causally related to overexpression of the serotonin transporter in primary pulmonary hypertension. Chest 2002; 121:978-88

54. Christman BW, McPherson CD, Newman JH, King GA, Bernard GR, Groves BM, Loyd JE: An imbalance between the excretion of thromboxane and prostacyclin metabolites in pulmonary hypertension. N Engl J Med 1992; 327:70-5

55. Giaid A, Saleh D: Reduced expression of endothelial nitric oxide synthase in the lungs of patients with pulmonary hypertension. N Engl J Med 1995; 333:214-21

56. Xue C, Johns RA: Endothelial nitric oxide synthase in the lungs of patients with pulmonary hypertension. N Engl J Med 1995; 333:1642-4

57. Frasch HF, Marshall C, Marshall BE: Endothelin-1 is elevated in monocrotaline pulmonary hypertension. Am J Physiol 1999; 276:L304-10

58. Stelzner TJ, O'Brien RF, Yanagisawa M, Sakurai T, Sato K, Webb S, Zamora M, McMurtry IF, Fisher JH: Increased lung endothelin-1 production in rats with idiopathic pulmonary hypertension. Am J Physiol 1992; 262:L614-L620

59. Cacoub P, Dorent R, Nataf P, Carayon A, Riquet M, Noe E, Piette JC, Godeau P, Gandjbakhch I: Endothelin-1 in the lungs of patients with pulmonary hypertension. Cardiovasc Res 1997; 33:196–200

60. Kumar P, Kazzi NJ, Shankaran S: Plasma immunoreactive endothelin-1 concentrations in infants with persistent pulmonary hypertension of the newborn. Am J Perinatol 1996; 13:335-41

61. Ishikawa S, Miyauchi T, Sakai S, Ushinohama H, Sagawa K, Fusazaki N, Kado H, Sunagawa H, Honda S, Ueno H: Elevated levels of plasma endothelin-1 in young patients with pulmonary hypertension caused by congenital heart disease are decreased after successful surgical repair. J Thorac Cardiovasc Surg 1995; 110:271-3

62. Allen SW, Chatfield BA, Koppenhafer SA, Schaffer MS, Wolfe RR, Abman SH: Circulating immunoreactive endothelin-1 in children with pulmonary hypertension: Association with acute hypoxic pulmonary vasoreactivity. Am Rev Respir Dis 1993; 148:519–22

63. Yoshibayashi M, Nishioka K, Nakao K, Saito Y, Matsumura M, Ueda T, Temma S, Shirakami G, Imura H, Mikawa H: Plasma endothelin concentrations in patients with pulmonary hypertension associated with congenital heart defects: Evidence for increased production of endothelin in pulmonary circulation. Circulation 1991; 84:2280-5

64. Stewart DJ, Levy RD, Cernacek P, Langleben D: Increased plasma endothelin-1 in pulmonary hypertension: Marker or mediator of disease? Ann Intern Med 1991; 114:464-9

65. Giaid A, Yanagisawa M, Langleben D, Michel RP, Levy R, Shennib H, Kimura S, Masaki T, Duguid WP, Stewart DJ: Expression of endothelin-1 in the lungs of patients with pulmonary hypertension. N Engl J Med 1993; 328:1732-9

66. Archer S, Rusch N: Potassium Channels in Cardiovascular Biology. Norwell, Massachusetts, Kluwer/Plenum, 2001

67. Weir EK, Archer SL: The mechanism of acute hypoxic pulmonary vasoconstriction: The tale of two channels. FASEB J 1995; 9:183-9 68. Post JM, Hume JR, Archer SL, Weir EK: Direct role for potassium channel inhibition in hypoxic pulmonary vasoconstriction. Am J Physiol 1992; 262:C882-C890

69. Michelakis ED, Weir EK: The pathobiology of pulmonary hypertension: Smooth muscle cells and ion channels. Clin Chest Med 2001; 22:419-32

70. Archer SL, Souil E, Dinh-Xuan AT, Schremmer B, Mercier JC, El Yaagoubi A, Nguyen-Huu L, Reeve HL, Hampl V: Molecular identification of the role of voltage-gated K+ channels, Kv1.5 and Kv2.1, in hypoxic pulmonary vasoconstriction and control of resting membrane potential in rat pulmonary artery myocytes. J Clin Invest 1998; 101:2319–30

71. Pepper MS, Vassalli JD, Orci L, Montesano R: Biphasic effect of transforming growth factor-beta 1 on in vitro angiogenesis. Exp Cell Res 1993; 204:356-63

72. Battegay EJ, Raines EW, Seifert RA, Bowen-Pope DF, Ross R: TGF-beta induces bimodal proliferation of connective tissue cells via complex control of an autocrine PDGF loop. Cell 1990; 63:515-24

73. Atkinson C, Stewart S, Imamura T, Trembath RC, Morrell NW: Immunolocalization of BMPR-II and TGF-B type I and II receptors in primary plexogenic pulmonary hypertension. J Heart Lung Transplant 2001; 20:149

74. Oh SP, Seki T, Goss KA, Imamura T, Yi Y, Donahoe PK, Li L, Miyazono K, ten Dijke P, Kim S, Li E: Activin receptor-like kinase 1 modulates transforming growth factor-beta 1 signaling in the regulation of angiogenesis. Proc Natl Acad Sci U S A 2000; 97:2626-31

75. Trembath RC, Thomson JR, Machado RD, Morgan NV, Atkinson C, Winship I, Simonneau G, Galie N, Loyd JE, Humbert M, Nichols WC, Morrell NW, Berg J, Manes A, McGaughran J, Pauciulo M, Wheeler L: Clinical and molecular genetic features of pulmonary hypertension in patients with hereditary hemorrhagic telangiectasia. N Engl J Med 2001; 345:325-34

76. Hirose S, Hosoda Y, Furuya S, Otsuki T, Ikeda E: Expression of vascular endothelial growth factor and its receptors correlates closely with formation of the plexiform lesion in human pulmonary hypertension. Pathol Int 2000; 50: 472-9

77. Taraseviciene-Stewart L, Kasahara Y, Alger L, Hirth P, Mc MG, Waltenberger J, Voelkel NF, Tuder RM: Inhibition of the VEGF receptor 2 combined with chronic hypoxia causes cell death-dependent pulmonary endothelial cell proliferation and severe pulmonary hypertension. FASEB J 2001; 15:427-38

78. Campbell AI, Zhao Y, Sandhu R, Stewart DJ: Cell-based gene transfer of vascular endothelial growth factor attenuates monocrotaline-induced pulmonary hypertension. Circulation 2001; 104:2242-8

79. Gurtner HP: Pulmonary hypertension, "plexogenic pulmonary arteriopathy" and the appetite depressant drug aminorex: Post or propter? Bull Physiopathol Respir (Nancy) 1979; 15:897–923

80. Kay JM, Smith P, Heath D: Aminorex and the pulmonary circulation. Thorax 1971; 26:262-70

81. Abenhaim L, Moride Y, Brenot F, Rich S, Benichou J, Kurz X, Higenbottam T, Oakley C, Wouters E, Aubier M, Simonneau G, Begaud B: Appetite-suppressant drugs and the risk of primary pulmonary hypertension. International Primary Pulmonary Hypertension Study Group. N Engl J Med 1996; 335:609-16

82. Fishman AP: Etiology and pathogenesis of primary pulmonary hypertension: A perspective. Chest 1998; 114:2428-78

83. Fristrom S, Airaksinen MM, Halmekoski J: Release of platelet 5-hydroxytryptamine by some anorexic and other sympathomimetics and their acetyl derivatives. Acta Pharmacol Toxicol (Copenh) 1977; 41:218-24

84. Buczko W, De Gaetano G, Garattini S: Effect of fenfluramine on 5-hydroxytryptamine uptake and release by rat blood platelets. Br J Pharmacol 1975; 53:563-8

85. Connolly HM, Crary JL, McGoon MD, Hensrud DD, Edwards BS, Edwards WD, Schaff HV: Valvular heart disease associated with fenfluramine-phentermine. N Engl J Med 1997; 337:581-8

86. Rothman RB, Ayestas MA, Dersch CM, Baumann MH: Aminorex, fenfluramine, and chlorphentermine are serotonin transporter substrates: Implications for primary pulmonary hypertension. Circulation 1999; 100:869-75

87. Weir EK, Reeve HL, Huang JM, Michelakis E, Nelson DP, Hampl V, Archer SL: Anorexic agents aminorex, fenfluramine, and dexfenfluramine inhibit potassium current in rat pulmonary vascular smooth muscle and cause pulmonary vasoconstriction. Circulation 1996; 94:2216-20

 Patel AJ, Lazdunski M, Honore E: Kv2.1/Kv9.3, an ATP-dependent delayedrectifier K+ channel in pulmonary artery myocytes. Ann N Y Acad Sci 1999; 868:438-41

89. Weir EK, Reeve HL, Johnson G, Michelakis ED, Nelson DP, Archer SL: A role for potassium channels in smooth muscle cells and platelets in the etiology of primary pulmonary hypertension. Chest 1998; 114:2008-48

90. Michelakis ED, Weir EK, Nelson DP, Reeve HL, Tolarova S, Archer SL: Dexfenfluramine elevates systemic blood pressure by inhibiting potassium currents in vascular smooth muscle cells. J Pharmacol Exp Ther 1999; 291:1143-9

91. Reeve HL, Archer SL, Soper M, Weir EK: Dexfenfluramine increases pulmonary artery smooth muscle intracellular Ca2+, independent of membrane potential. Am J Physiol 1999; 277:L662-6

92. Wang J, Juhaszova M, Conte JV Jr, Gaine SP, Rubin LJ, Yuan JX: Action of fenfluramine on voltage-gated K+ channels in human pulmonary-artery smoothmuscle cells (letter). Lancet 1998; 352:290

93. Wellman PJ, Maher TJ: Synergistic interactions between fenfluramine and phentermine. Int J Obes Relat Metab Disord 1999; 23:723-32

94. Rich S, Dantzker DR, Ayres SM, Bergofsky EH, Brundage BH, Detre KM,

Fishman AP, Goldring RM, Groves BM, Koerner SK: Primary pulmonary hypertension: A national prospective study. Ann Intern Med 1987; 107:216-23

 Mikhail GW, Gibbs JS, Yacoub MH: Pulmonary and systemic arterial pressure changes during syncope in primary pulmonary hypertension. Circulation 2001: 104:1326-7

96. McGoon MD, Fuster V, Freeman WK, Edwards WD, Scott JP: Pulmonary hypertension, Mayo Clinic Practice of Cardiology, 3rd edition. Edited by Guiliani ER. St. Louis, Mosby, 1996, pp 1815-36

97. D'Alonzo GE, Barst RJ, Ayres SM, Bergofsky EH, Brundage BH, Detre KM, Fishman AP, Goldring RM, Groves BM, Kernis JT: Survival in patients with primary pulmonary hypertension: Results from a national prospective registry. Ann Intern Med 1991; 115:343-9

98. Sitbon O, Brenot F, Denjean A, Bergeron A, Parent F, Azarian R, Herve P, Raffestin B, Simonneau G: Inhaled nitric oxide as a screening vasodilator agent in primary pulmonary hypertension: A dose-response study and comparison with prostacyclin. Am J Respir Crit Care Med 1995; 151:384-9

99. Sitbon O, Humbert M, Jagot JL, Taravella O, Fartoukh M, Parent F, Herve P, Simonneau G: Inhaled nitric oxide as a screening agent for safely identifying responders to oral calcium-channel blockers in primary pulmonary hypertension. Eur Respir J 1998; 12:265-70

100. Pepke-Zaba J, Higenbottam TW, Dinh-Xuan AT, Stone D, Wallwork J: Inhaled nitric oxide as a cause of selective pulmonary vasodilatation in pulmonary hypertension. Lancet 1991; 338:1173-4

101. Langleben D, Moroz LA, McGregor M, Lisbona R: Decreased half-life of fibrinogen in primary pulmonary hypertension. Thromb Res 1985; 40:577-80

102. Eisenberg PR, Lucore C, Kaufman L, Sobel BE, Jaffe AS, Rich S: Fibrinopeptide A levels indicative of pulmonary vascular thrombosis in patients with primary pulmonary hypertension. Circulation 1990; 82:841-7

103. Weatherford DA, Sackman JE, Reddick TT, Freeman MB, Stevens SL, Goldman MH: Vascular endothelial growth factor and heparin in a biologic glue promotes human aortic endothelial cell proliferation with aortic smooth muscle cell inhibition. Surgery 1996; 120:433-9

104. Rich S, Kaufmann E, Levy PS: The effect of high doses of calcium-channel blockers on survival in primary pulmonary hypertension. N Engl J Med 1992; 327:76-81

105. Saadjian A, Philip-Joet F, Hot B, Reynaud-Gaubert M, Durand A, Levy S, Arnaud A: Effects of nicardipine on pulmonary and systemic vascular reactivity to oxygen in patients with pulmonary hypertension secondary to chronic obstructive lung disease. J Cardiovasc Pharmacol 1991; 17:731-7

106. Clark RH, Kueser TJ, Walker MW, Southgate WM, Huckaby JL, Perez JA, Roy BJ, Keszler M, Kinsella JP: Low-dose nitric oxide therapy for persistent pulmonary hypertension of the newborn. Clinical Inhaled Nitric Oxide Research Group. N Engl J Med 2000; 342:469–74

107. Ricciardi MJ, Knight BP, Martinez FJ, Rubenfire M: Inhaled nitric oxide in primary pulmonary hypertension: A safe and effective agent for predicting response to nifedipine. J Am Coll Cardiol 1998; 32:1068–73

108. Petros AJ, Turner SC, Nunn AJ: Cost implications of using inhaled nitric oxide compared with epoprostenol for pulmonary hypertension. J Pharm Technol 1995; 11:163-6

109. Higenbottam T, Stenmark K, Simonneau G: Treatments for severe pulmonary hypertension. Lancet 1999; 353:338-40

110. Katayama Y, Higenbottam TW, Cremona G, Akamine S, Demoncheaux EA, Smith AP, Siddons TE: Minimizing the inhaled dose of NO with breath-bybreath delivery of spikes of concentrated gas. Circulation 1998; 98:2429-32

111. Perez-Penate G, Julia-Serda G, Pulido-Duque JM, Gorriz-Gomez E, Cabrera-Navarro P: One-year continuous inhaled nitric oxide for primary pulmonary hypertension. Chest 2001; 119:970-3

112. Roberts JD Jr, Chiche JD, Weimann J, Steudel W, Zapol WM, Bloch KD: Nitric oxide inhalation decreases pulmonary artery remodeling in the injured lungs of rat pups. Circ Res 2000; 87:140-5

113. Maruyama J, Maruyama K, Mitani Y, Kitabatake M, Yamauchi T, Miyasaka K: Continuous low-dose NO inhalation does not prevent monocrotaline-induced pulmonary hypertension in rats. Am J Physiol 1997; 272:H517-24

114. Horstman DJ, Frank DU, Rich GF: Prolonged inhaled NO attenuates hypoxic, but not monocrotaline-induced, pulmonary vascular remodeling in rats. Anesth Analg 1998; 86:74-81

115. Jeffery TK, Wanstall JC: Pulmonary vascular remodeling: A target for therapeutic intervention in pulmonary hypertension. Pharmacol Ther 2001; 92:1-20

116. Kourembanas S, McQuillan LP, Leung GK, Faller DV: Nitric oxide regulates the expression of vasoconstrictors and growth factors by vascular endothelium under both normoxia and hypoxia. J Clin Invest 1993; 92:99-104

117. Mitani Y, Zaidi SH, Dufourcq P, Thompson K, Rabinovitch M: Nitric oxide reduces vascular smooth muscle cell elastase activity through cGMPmediated suppression of ERK phosphorylation and AML1B nuclear partitioning. FASEB J 2000; 14:805–14

118. Jiang ZY, Costachescu T, Derouin M, Blaise G: Treatment of pulmonary hypertension during surgery with nitric oxide and vasodilators. Can J Anaesth 2000; 47:552-5

119. Fullerton DA, Jones SD, Jaggers J, Piedalue F, Grover FL, McIntyre RC Jr: Effective control of pulmonary vascular resistance with inhaled nitric oxide after cardiac operation. J Thorac Cardiovasc Surg 1996; 111:753-62

120. Wilkens H, Guth A, Konig J, Forestier N, Cremers B, Hennen B, Bohm M,

Sybrecht GW: Effect of inhaled iloprost plus oral sildenafil in patients with primary pulmonary hypertension. Circulation 2001; 104:1218-22

121. Michelakis E, Tymchak W, Lien D, Webster L, Hashimoto K, Archer S: Oral sildenafil is an effective and specific pulmonary vasodilator in patients with pulmonary arterial hypertension: Comparison with inhaled nitric oxide. Circulation 2002; 105:2398-403

122. Atz AM, Lefler AK, Fairbrother DL, Uber WE, Bradley SM: Sildenafil augments the effect of inhaled nitric oxide for postoperative pulmonary hypertensive crises. J Thorac Cardiovasc Surg 2002; 124:628-9

123. Ghofrani HA, Wiedemann R, Rose F, Schermuly RT, Olschewski H, Weissmann N, Gunther A, Walmrath D, Seeger W, Grimminger F: Sildenafil for treatment of lung fibrosis and pulmonary hypertension: A randomised controlled trial. Lancet 2002; 360:895–900

124. Shekerdemian LS, Ravn HB, Penny DJ: Intravenous sildenafil lowers pulmonary vascular resistance in a model of neonatal pulmonary hypertension. Am J Respir Crit Care Med 2002; 165:1098-102

125. Abrams D, Schulze-Neick I, Magee AG: Sildenafil as a selective pulmonary vasodilator in childhood primary pulmonary hypertension. Heart 2000; 84:E4

126. Dweik RA, Laskowski D, Abu-Soud HM, Kaneko F, Hutte R, Stuehr DJ, Erzurum SC: Nitric oxide synthesis in the lung: Regulation by oxygen through a kinetic mechanism. J Clin Invest 1998; 101:660-6

127. Prasad S, Wilkinson J, Gatzoulis MA: Sildenafil in primary pulmonary hypertension (letter). N Engl J Med 2000; 343:1342

128. Medina P, Segarra G, Torondel B, Chuan P, Domenech C, Vila JM, Lluch S: Inhibition of neuroeffector transmission in human vas deferens by sildenafil. Br J Pharmacol 2000: 131:871-4

129. Rich S, McLaughlin VV: The effects of chronic prostacyclin therapy on cardiac output and symptoms in primary pulmonary hypertension. J Am Coll Cardiol 1999; 34:1184-7

130. Rubin LJ, Groves BM, Reeves JT, Frosolono M, Handel F, Cato AE: Prostacyclin-induced acute pulmonary vasodilation in primary pulmonary hypertension. Circulation 1982; 66:334-8

131. Friedman R, Mears JG, Barst RJ: Continuous infusion of prostacyclin normalizes plasma markers of endothelial cell injury and platelet aggregation in primary pulmonary hypertension. Circulation 1997; 96:2782-4

132. Langleben D, Barst RJ, Badesch D, Groves BM, Tapson VF, Murali S, Bourge RC, Ettinger N, Shalit E, Clayton LM, Jobsis MM, Blackburn SD, Crow JW, Stewart DJ, Long W: Continuous infusion of epoprostenol improves the net balance between pulmonary endothelin-1 clearance and release in primary pulmonary hypertension. Circulation 1999; 99:3266-71

133. Barst RJ, Rubin LJ, Long WA, McGoon MD, Rich S, Badesch DB, Groves BM, Tapson VF, Bourge RC, Brundage BH: A comparison of continuous intravenous epoprostenol (prostacyclin) with conventional therapy for primary pulmonary hypertension. The Primary Pulmonary Hypertension Study Group. N Engl J Med 1996; 334:296–302

134. Barst RJ, Rubin LJ, McGoon MD, Caldwell EJ, Long WA, Levy PS: Survival in primary pulmonary hypertension with long-term continuous intravenous prostacyclin. Ann Intern Med 1994; 121:409-15

135. Wax D, Garofano R, Barst RJ: Effects of long-term infusion of prostacyclin on exercise performance in patients with primary pulmonary hypertension. Chest 1999; 116:914-20

136. Badesch DB, Tapson VF, McGoon MD, Brundage BH, Rubin LJ, Wigley FM, Rich S, Barst RJ, Barrett PS, Kral KM, Jobsis MM, Loyd JE, Murali S, Frost A, Girgis R, Bourge RC, Ralph DD, Elliott CG, Hill NS, Langleben D, Schilz RJ, McLaughlin VV, Robbins IM, Groves BM, Shapiro S, Medsger TA Jr: Continuous intravenous epoprostenol for pulmonary hypertension due to the scleroderma spectrum of disease: A randomized, controlled trial. Ann Intern Med 2000; 132:425–34

137. McLaughlin VV, Gaine SP, Barst RJ, Oudiz RJ, Bourge RC, Frost A, Robbins IM, Tapson VF, McGoon MD, Badesch DB, Sigman J, Roscigno R, Blackburn SD, Arneson C, Rubin IJ, Rich S: Efficacy and safety of treprostinil: An epoprostenol analog for primary pulmonary hypertension. J Cardiovasc Pharmacol 2003; 41: 293–9

138. Simonneau G, Barst RJ, Galie N, Naeije R, Rich S, Bourge RC, Keogh A, Oudiz R, Frost A, Blackburn SD, Crow JW, Rubin LJ: Continuous subcutaneous infusion of treprostinil, a prostacyclin analogue, in patients with pulmonary arterial hypertension: A double-blind, randomized, placebo-controlled trial. Am J Respir Crit Care Med 2002; 165:800-4

139. Olschewski H, Walmrath D, Schermuly R, Ghofrani A, Grimminger F, Seeger W: Aerosolized prostacyclin and iloprost in severe pulmonary hypertension. Ann Intern Med 1996; 124:820-4

140. Hoeper MM, Olschewski H, Ghofrani HA, Wilkens H, Winkler J, Borst MM, Niedermeyer J, Fabel H, Seeger W: A comparison of the acute hemodynamic effects of inhaled nitric oxide and aerosolized iloprost in primary pulmonary hypertension. German PPH Study Group. J Am Coll Cardiol 2000; 35:176-82

141. Schermuly RT, Krupnik E, Tenor H, Schudt C, Weissmann N, Rose F, Grimminger F, Seeger W, Walmrath D, Ghofrani HA: Coaerosolization of phosphodiesterase inhibitors markedly enhances the pulmonary vasodilatory response to inhaled iloprost in experimental pulmonary hypertension: Maintenance of lung selectivity. Am J Respir Crit Care Med 2001; 164:1694-700

142. Weimann J, Ullrich R, Hromi J, Fujino Y, Clark MW, Bloch KD, Zapol WM: Sildenafil is a pulmonary vasodilator in awake lambs with acute pulmonary hypertension. ANESTHESIOLOGY 2000; 92:1702-12 143. Zhao L, Mason NA, Morrell NW, Kojonazarov B, Sadykov A, Maripov A, Mirrakhimov MM, Aldashev A, Wilkins MR: Sildenafil inhibits hypoxia-induced pulmonary hypertension. Circulation 2001; 104:424-8

144. Lodato RF: Viagra for impotence of pulmonary vasodilator therapy? Am J Respir Crit Care Med 2001; 163:312-3

145. Nagaya N, Uematsu M, Okano Y, Satoh T, Kyotani S, Sakamaki F, Nakanishi N, Miyatake K, Kunieda T: Effect of orally active prostacyclin analogue on survival of outpatients with primary pulmonary hypertension. J Am Coll Cardiol 1999; 34:1188–92

146. Chen SJ, Chen YF, Opgenorth TJ, Wessale JL, Meng QC, Durand J, DiCarlo VS, Oparil S: The orally active nonpeptide endothelin A-receptor antagonist A-127722 prevents and reverses hypoxia-induced pulmonary hypertension and pulmonary vascular remodeling in Sprague-Dawley rats. J Cardiovasc Pharmacol 1997; 29:713–25

147. Dupuis J, Cernacek P, Tardif JC, Stewart DJ, Gosselin G, Dyrda I, Bonan R, Crepeau J: Reduced pulmonary clearance of endothelin-1 in pulmonary hypertension. Am Heart J 1998; 135:614-20

148. Dupuis J, Jasmin JF, Prie S, Cernacek P: Importance of local production of endothelin-1 and of the ET(B)receptor in the regulation of pulmonary vascular tone. Pulm Pharmacol Ther 2000; 13:135-40

149. de Nucci G, Thomas R, D'Orleans-Juste P, Antunes E, Walder C, Warner TD, Vane JR: Pressor effects of circulating endothelin are limited by its removal in the pulmonary circulation and by the release of prostacyclin and endothelium-derived relaxing factor. Proc Natl Acad Sci U S A 1988; 85:9797–800

150. Channick RN, Simonneau G, Sitbon O, Robbins IM, Frost A, Tapson VF, Badesch DB, Roux S, Rainisio M, Bodin F, Rubin LJ: Effects of the dual endothelinreceptor antagonist bosentan in patients with pulmonary hypertension: A randomised placebo-controlled study. Lancet 2001; 358:1119–23

151. Newman JH: Treatment of primary pulmonary hypertension: The next generation. N Engl J Med 2002; 346:933-5

152. Rubin LJ, Badesch DB, Barst RJ, Galie N, Black CM, Keogh A, Pulido T, Frost A, Roux S, Leconte I, Landzberg M, Simonneau G: Bosentan therapy for pulmonary arterial hypertension. N Engl J Med 2002; 346:896-903

153. Food and Drug Administration Office of Public Affairs: FDA approves first oral medication for pulmonary arterial hypertension. T01-60. U.S. Food and Drug Administration, October 20, 2001

154. Faul JL, Nishimura T, Berry GJ, Benson GV, Pearl RG, Kao PN: Triptolide attenuates pulmonary arterial hypertension and neointimal formation in rats. Am J Respir Crit Care Med 2000; 162:2252-8

155. Cowan KN, Heilbut A, Humpl T, Lam C, Ito S, Rabinovitch M: Complete reversal of fatal pulmonary hypertension in rats by a serine elastase inhibitor. Nat Med 2000; 6:698-702

156. Cowan KN, Jones PL, Rabinovitch M: Elastase and matrix metalloproteinase inhibitors induce regression, and tenascin-C antisense prevents progression, of vascular disease. J Clin Invest 2000; 105:21-34

157. Cowan KN, Jones PL, Rabinovitch M: Regression of hypertrophied rat pulmonary arteries in organ culture is associated with suppression of proteolytic activity, inhibition of tenascin-C, and smooth muscle cell apoptosis. Circ Res 1999; 84:1223-33

158. Champion HC, Bivalacqua TJ, D'Souza FM, Ortiz LA, Jeter JR, Toyoda K, Heistad DD, Hyman AL, Kadowitz PJ: Gene transfer of endothelial nitric oxide synthase to the lung of the mouse in vivo: Effect on agonist-induced and flow-mediated vascular responses. Circ Res 1999; 84:1422-32

159. Champion HC, Bivalacqua TJ, Toyoda K, Heistad DD, Hyman AL, Kadowitz PJ: In vivo gene transfer of prepro-calcitonin gene-related peptide to the lung attenuates chronic hypoxia-induced pulmonary hypertension in the mouse. Circulation 2000; 101:923–30

160. Campbell AI, Kuliszewski MA, Stewart DJ: Cell-based gene transfer to the pulmonary vasculature: Endothelial nitric oxide synthase overexpression inhibits monocrotaline-induced pulmonary hypertension. Am J Respir Cell Mol Biol 1999; 21:567–75

161. Cuenco J, Tzeng G, Wittels B: Anesthetic management of the parturient with systemic lupus erythematosus, pulmonary hypertension, and pulmonary edema. ANESTHESIOLOGY 1999; 91:568-70

162. Tay SM, Ong BC, Tan SA: Cesarean section in a mother with uncorrected congenital coronary to pulmonary artery fistula. Can J Anaesth 1999; 46:368-71

163. Weiss BM, Maggiorini M, Jenni R, Lauper U, Popov V, Bombeli T, Spahn DR: Pregnant patient with primary pulmonary hypertension: Inhaled pulmonary vasodilators and epidural anesthesia for cesarean delivery. ANESTHESIOLOGY 2000; 92:1191-4

164. Medical Research Council Working Party: Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema: Report of the Medical Research Council Working Party. Lancet 1981; 1:681-6

165. Rich S, Seidlitz M, Dodin E, Osimani D, Judd D, Genthner D, McLaughlin V, Francis G: The short-term effects of digoxin in patients with right ventricular dysfunction from pulmonary hypertension. Chest 1998; 114:787-92

166. Rudolph AM, Yuan S: Response of the pulmonary vasculature to hypoxia and $\rm H+$ ion concentration changes. J Clin Invest 1966; 45:399–411

167. Reves JG, Kissin I, Lell WA, Tosone S: Calcium entry blockers: Uses and implications for anesthesiologists. ANESTHESIOLOGY 1982; 57:504-18

168. Lynch C III: Are volatile anesthetics really calcium entry blockers? ANES-THESIOLOGY 1984; 61:644-6 169. Bradford KK, Deb B, Pearl RG: Combination therapy with inhaled nitric oxide and intravenous dobutamine during pulmonary hypertension in the rabbit. J Cardiovasc Pharmacol 2000; 36:146–51

170. Tanaka H, Tajimi K, Moritsune O, Kobayashi K, Okada K: Effects of milrinone on pulmonary vasculature in normal dogs and in dogs with pulmonary hypertension. Crit Care Med 1991; 19:68–74

171. Angle MR, Molloy DW, Penner B, Jones D, Prewitt RM: The cardiopulmonary and renal hemodynamic effects of norepinephrine in canine pulmonary embolism. Chest 1989; 95:1333-7

172. Hirsch LJ, Rooney MW, Wat SS, Kleinmann B, Mathru M: Norepinephrine and phenylephrine effects on right ventricular function in experimental canine pulmonary embolism. Chest 1991; 100:796-801

173. Kwak YL, Lee CS, Park YH, Hong YW: The effect of phenylephrine and norepinephrine in patients with chronic pulmonary hypertension. Anaesthesia 2002; 57:9-14

174. Tritapepe L, Voci P, Cogliati AA, Pasotti E, Papalia U, Menichetti A: Successful weaning from cardiopulmonary bypass with central venous prostaglandin E1 and left atrial norepinephrine infusion in patients with acute pulmonary hypertension. Crit Care Med 1999; 27:2180-3

175. Bichel T, Spahr-Schopfer I, Berner M, Jaeggi E, Velkovski Y, Friedli B, Kalangos A, Faidutti B, Rouge JC: Successful weaning from cardiopulmonary bypass after cardiac surgery using inhaled nitric oxide. Paediatr Anaesth 1997; 7:335-9

176. Schroeder RA, Wood GL, Plotkin JS, Kuo PC: Intraoperative use of inhaled PGI(2) for acute pulmonary hypertension and right ventricular failure. Anesth Analg 2000; 91:291-5

177. Walmrath D, Schneider T, Schermuly R, Olschewski H, Grimminger F, Seeger W: Direct comparison of inhaled nitric oxide and aerosolized prostacyclin in acute respiratory distress syndrome. Am J Respir Crit Care Med 1996; 153: 991-6

178. Troncy E, Francoeur M, Salazkin I, Yang F, Charbonneau M, Leclerc G, Vinay P, Blaise G: Extra-pulmonary effects of inhaled nitric oxide in swine with and without phenylephrine. Br J Anaesth 1997; 79:631-40

179. Troncy E, Blaise G: Phenylephrine and inhaled nitric oxide. ANESTHESIOL-OGY 1998; 89:538-40

180. Veering BT, Cousins MJ: Cardiovascular and pulmonary effects of epidural anaesthesia. Anaesth Intensive Care 2000; 28:620-35

181. Pollard JB: Common mechanisms and strategies for prevention and treatment of cardiac arrest during epidural anesthesia. J Clin Anesth 2002; 14:52-6

182. Burrows FA, Klinck JR, Rabinovitch M, Bohn DJ: Pulmonary hypertension in children: Perioperative management. Can Anaesth Soc J 1986; 33:606-28

183. Rouby JJ, Andreev A, Leger P, Arthaud M, Landault C, Vicaut E, Maistre G, Eurin J, Gandjbakch I, Viars P: Peripheral vascular effects of thiopental and propofol in humans with artificial hearts. ANESTHESIOLOGY 1991; 75:32-42

184. Claeys MA, Gepts E, Camu F: Haemodynamic changes during anaesthesia induced and maintained with propofol. Br J Anaesth 1988; 60:3-9

185. Horibe M, Ogawa K, Sohn JT, Murray PA: Propofol attenuates acetylcholine-induced pulmonary vasorelaxation: Role of nitric oxide and endotheliumderived hyperpolarizing factors. ANESTHESIOLOGY 2000; 93:447-55

186. Kondo U, Kim SO, Murray PA: Propofol selectively attenuates endothelium-dependent pulmonary vasodilation in chronically instrumented dogs. Anes-THESIOLOGY 2000; 93:437-46

187. Kondo U, Kim SO, Nakayama M, Murray PA: Pulmonary vascular effects of propofol at baseline, during elevated vasomotor tone, and in response to a state of the second second

sympathetic α - and β -adrenoreceptor activation. ANESTHESIOLOGY 2001; 94:815–23 188. Mertens E, Saldien V, Coppejans H, Bettens K, Vercauteren M: Target controlled infusion of remiferitanil and propofol for cesarean section in a patient with multivalvular disease and severe pulmonary hypertension. Acta Anaesthesiol Belg 2001: 52:207–9

189. White PF, Way WL, Trevor AJ: Ketamine: Its pharmacology and the rapeutic uses. Anesthesiology 1982; 56:119-36

190. Rich GF, Roos CM, Anderson SM, Daugherty MO, Uncles DR: Direct effects of intravenous anesthetics on pulmonary vascular resistance in the isolated rat lung. Anesth Analg 1994; 78:961-6

191. Lee TS, Hou X: Vasoactive effects of ketamine on isolated rabbit pulmonary arteries. Chest 1995; 107:1152-5

192. Miyawaki I, Nakamura K, Terasako K, Toda H, Kakuyama M, Mori K: Modification of endothelium-dependent relaxation by propofol, ketamine, and midazolam. Anesth Analg 1995; 81:474-9

193. Ogawa K, Tanaka S, Murray PA: Inhibitory effects of etomidate and ketamine on endothelium-dependent relaxation in canine pulmonary artery. ANESTHESIOLOGY 2001; 94:668-77

194. Fulton D, Mcgiff JC, Quilley J: Pharmacological evaluation of an epoxide as the putative hyperpolarizing factor mediating the nitric oxide-independent vasodilator effect of bradykinin in the rat heart. J Pharmacol Exp Ther 1998; 287:497-503

195. Lennon PF, Murray PA: Attenuated hypoxic pulmonary vasoconstriction during isoflurane anesthesia is abolished by cyclooxygenase inhibition in chronically instrumented dogs. ANESTHESIOLOGY 1996; 84:404-14

196. Lennon PF, Murray PA: Isoflurane and the pulmonary vascular pressureflow relation at baseline and during sympathetic α - and β -adrenoreceptor activation in chronically instrumented dogs. ANESTHESIOLOGY 1995; 82:723-33

Anesthesiology, V 99, No 6, Dec 2003

Copyright © by the American Society of Anesthesiologists. Unauthorized reproduction of this article is prohibited.

197. Nichols CG, Lederer WJ: Adenosine triphosphate-sensitive potassium channels in the cardiovascular system. Am J Physiol 1991; 261:H1675-86

198. Gambone LM, Murray PA, Flavahan NA: Isoflurane anesthesia attenuates endothelium-dependent pulmonary vasorelaxation by inhibiting the synergistic interaction between nitric oxide and prostacyclin. ANESTHESIOLOGY 1997; 86: 936-44

199. Fujiwara Y, Murray PA: Effects of isoflurane anesthesia on pulmonary vascular response to K^+ ATP channel activation and circulatory hypotension in chronically instrumented dogs. ANESTHESIOLOGY 1999; 90:799–811

200. Seki S, Sato K, Nakayama M, Murray PA: Halothane and enflurane attenuate pulmonary vasodilation mediated by adenosine triphosphate-sensitive potassium channels compared to the conscious state. ANESTHESIOLOGY 1997; 86: 923-35

201. Nakayama M, Kondo U, Murray PA: Pulmonary vasodilator response to adenosine triphosphate-sensitive potassium channel activation is attenuated during desflurane but preserved during sevoflurane anesthesia compared with the conscious state. ANESTHESIOLOGY 1998: 88:1023-35

202. Sykes MK, Hurtig JB, Tait AR, Chakrabarti MK: Reduction of hypoxic pulmonary vasoconstriction in the dog during administration of nitrous oxide. Br J Anaesth 1977; 49:301–7

203. Benumof JL, Wahrenbrock EA: Local effects of anesthetics on regional hypoxic pulmonary vasoconstriction. ANESTHESIOLOGY 1975; 43:525-32

204. Pagel PS, Kampine JP, Schmeling WT, Warltier DC: Effects of nitrous oxide on myocardial contractility as evaluated by the preload recruitable stroke work relationship in chronically instrumented dogs. ANESTHESIOLOGY 1990; 73: 1148-57

205. Hickey PR, Hansen DD, Strafford M, Thompson JE, Jonas RE, Mayer JE: Pulmonary and systemic hemodynamic effects of nitrous oxide in infants with normal and elevated pulmonary vascular resistance. ANESTHESIOLOGY 1986; 65: 374-8

206. Schulte-Sasse U, Hess W, Tarnow J: Pulmonary vascular responses to nitrous oxide in patients with normal and high pulmonary vascular resistance. Anesthesiology 1982; 57:9-13

207. Heerdt PM, Caldwell RW: The mechanism of nitrous oxide-induced changes in pulmonary vascular resistance in a dog model of left atrial outflow obstruction. J Cardiothorac Anesth 1989; 3:568-73

208. Hilgenberg JC, McCammon RL, Stoelting RK: Pulmonary and systemic vascular responses to nitrous oxide in patients with mitral stenosis and pulmonary hypertension. Anesth Analg 1980; 59:323-6

209. Konstadt SN, Reich DL, Thys DM: Nitrous oxide does not exacerbate pulmonary hypertension or ventricular dysfunction in patients with mitral valvular disease. Can J Anaesth 1990; 37:613-7

210. Robinson RJ, Shennib H, Noirclerc M: Slow-rate, high-pressure ventilation: A method of management of difficult transplant recipients during sequential double lung transplantation for cystic fibrosis. J Heart Lung Transplant 1994; 13:779-84

211. Ardehali A, Laks H, Levine M, Shpiner R, Ross D, Watson LD, Shvartz O, Sangwan S, Waters PF: A prospective trial of inhaled nitric oxide in clinical lung transplantation. Transplantation 2001; 72:112-5

212. Kemming GI, Merkel MJ, Schallerer A, Habler OP, Kleen MS, Haller M, Briegel J, Vogelmeier C, Furst H, Reichart B, Zwissler B: Inhaled nitric oxide (NO) for the treatment of early allograft failure after lung transplantation. Munich Lung Transplant Group. Intensive Care Med 1998: 24:1173–80

213. Thabut G, Brugiere O, Leseche G, Stern JB, Fradj K, Herve P, Jebrak G, Marty J, Fournier M, Mal H: Preventive effect of inhaled nitric oxide and pentoxifylline on ischemia/reperfusion injury after lung transplantation. Transplantation 2001; 71:1295-300

214. Miller OI, Tang SF, Keech A, Pigott NB, Beller E, Celermajer DS: Inhaled nitric oxide and prevention of pulmonary hypertension after congenital heart surgery: A randomised double-blind study. Lancet 2000; 356:1464-9

215. Hubert BP, Blaise GA: Inhaled NO given perioperatively improves oxygenation and decreases pulmonary arterial resistance following cardiopulmonary bypass in a pig model (abstract). ANESTHENOLOGY 2001; 95:A-366

216. Hubert BP, Radomski A, Blaise GA: Does inhaled nitric oxide (inhNO) affect matrix-metalloproteinase (MMP) concentration in bronchoalveolar lavage fluid (BAL) after cardiopulmonary bypass (CPB)? (abstract). ANESTHESIOLOGY 2001; 95:A-440

217. Radomski MW, Palmer RM, Moncada S: Comparative pharmacology of endothelium-derived relaxing factor, nitric oxide and prostacyclin in platelets. Br J Pharmacol 1987; 92:181-7

218. van Heerden PV, Gibbs NM, Michalopoulos N: Effect of low concentra-

tions of prostacyclin on platelet function in vitro. Anaesth Intensive Care 1997; 25:343-6

219. O'Hare R, McLoughlin C, Milligan K, McNamee D, Sidhu H: Anaesthesia for caesarean section in the presence of severe primary pulmonary hypertension. Br J Anaesth 1998; 81:790–2

220. van Heerden PV, Barden A, Michalopoulos N, Bulsara MK, Roberts BL: Dose-response to inhaled aerosolized prostacyclin for hypoxemia due to ARDS. Chest 2000; 117:819-27

221. Wood KE: Major pulmonary embolism: Review of a pathophysiologic approach to the golden hour of hemodynamically significant pulmonary embolism. Chest 2002; 121:877-905

222. Easterling TR, Ralph DD, Schmucker BC: Pulmonary hypertension in pregnancy: Treatment with pulmonary vasodilators. Obstet Gynecol 1999; 93: 494-8

223. Nelson DM, Main E, Crafford W, Ahumada GG: Peripartum heart failure due to primary pulmonary hypertension. Obstet Gynecol 1983; 62:58s-63s

224. Metcalfe J, McAnulty J, Ueland K: Burivell and Metcalfe's Heart Disease and Pregnancy: Physiology and Management, 2nd edition. Boston, Little, Brown, 1984, pp 265-77

225. Olofsson C, Bremme K, Forssell G, Ohqvist G: Cesarean section under epidural ropivacaine 0.75% in a parturient with severe pulmonary hypertension. Acta Anaesthesiol Scand 2001; 45:258-60

226. Hoeper MM, Maier R, Tongers J, Niedermeyer J, Hohlfeld JM, Hamm M, Fabel H: Determination of cardiac output by the Fick method, thermodilution, and acetylene rebreathing in pulmonary hypertension. Am J Respir Crit Care Med 1999; 160:535-41

227. Sitbon O, Brenot F, Denjean A, Bergeron A, Parent F, Azarian R, Herve P, Raffestin B, Simonneau G: Inhaled nitric oxide as a screening vasodilator agent in primary pulmonary hypertension: A dose-response study and comparison with prostacyclin. Am J Respir Crit Care Med 1995; 151:384-9

228. al Alaiyan S, al Omran A, Dyer D: The use of phosphodiesterase inhibitor (dipyridamole) to wean from inhaled nitric oxide. Intensive Care Med 1996; 22:1093-5

229. Buysse C, Fonteyne C, Dessy H, De Laet MH, Biarent D: The use of dipyridamole to wean from inhaled nitric oxide in congenital diaphragmatic hernia. J Pediatr Surg 2001; 36:1864-5

230. Fullerton DA, Jaggers J, Piedalue F, Grover FL, McIntyre RC Jr: Effective control of refractory pulmonary hypertension after cardiac operations. J Thorac Cardiovasc Surg 1997; 113:363-8

231. Ivy DD, Kinsella JP, Ziegler JW, Abman SH: Dipyridamole attenuates rebound pulmonary hypertension after inhaled nitric oxide withdrawal in post-operative congenital heart disease. J Thorac Cardiovasc Surg 1998; 115:875-82

232. Jiang ZY, Costachescu T, Derouin M, Blaise G: Treatment of pulmonary hypertension during surgery with nitric oxide and vasodilators. Can J Anaesth 2000; 47:552-5

233. Hoeper MM, Schwarze M, Ehlerding S, Adler-Schuermeyer A, Spiekerkoetter E, Niedermeyer J, Hamm M, Fabel H: Long-term treatment of primary pulmonary hypertension with aerosolized iloprost, a prostacyclin analogue. N Engl J Med 2000; 342:1866–70

234. Wilkens H, Guth A, Konig J, Forestier N, Cremers B, Hennen B, Bohm M, Sybrecht GW: Effect of inhaled iloprost plus oral sildenafil in patients with primary pulmonary hypertension. Circulation 2001; 104:1218-22

235. Rosner B: Fundamental of Biostatistics, 4th edition. Belmont, California, Duxburry, 1995

236. Fiser SM, Cope JT, Kron IL, Kaza AK, Long SM, Kern JA, Tribble CG, Lowson SM: Aerosolized prostacyclin (epoprostenol) as an alternative to inhaled nitric oxide for patients with reperfusion injury after lung transplantation. J Thorac Cardiovasc Surg 2001; 121:981-2

237. Hache M, Denault AY, Belisle S, Couture P, Babin D, Tetrault F, Guimond JG: Inhaled prostacyclin (PGI2) is an effective addition to the treatment of pulmonary hypertension and hypoxia in the operating room and intensive care unit. Can J Anaesth 2001; 48:924-9

238. Stewart R, Tuazon D, Olson G, Duarte AG: Pregnancy and primary pulmonary hypertension: Successful outcome with epoprostenol therapy. Chest 2001; 119:973-5

239. Martin JT, Tautz TJ, Antognini JF: Safety of regional anesthesia in Eisenmenger's syndrome. Reg Anesth Pain Med 2002; 27:509-13

240. Hohn L, Schweizer A, Morel DR, Spiliopoulos A, Licker M: Circulatory failure after anesthesia induction in a patient with severe primary pulmonary hypertension. ANESTHESIOLOGY 1999; 91:1943-5

241. Smedstad KG, Cramb R, Morison DH: Pulmonary hypertension and pregnancy: A series of eight cases. Can J Anaesth 1994; 41:502-12