

Will This Patient Be Difficult to Intubate?

The Rational Clinical Examination Systematic Review

Michael E. Detsky, MD, MSHP; Naheed Jivraj, MBBS, MSc; Neill K. Adhikari, MDCM, MSc; Jan O. Friedrich, MD, MSc, DPhil; Ruxandra Pinto, PhD; David L. Simel, MD; Duminda N. Wijeyesundera, MD, PhD; Damon C. Scales, MD, PhD

IMPORTANCE Recognizing patients in whom endotracheal intubation is likely to be difficult can help alert physicians to the need for assistance from a clinician with airway training and having advanced airway management equipment available.

OBJECTIVE To identify risk factors and physical findings that predict difficult intubation.

DATA SOURCES The databases of MEDLINE and EMBASE were searched from 1946 to June 2018 and from 1947 to June 2018, respectively, and the reference lists from the retrieved articles and previous reviews were searched for additional studies.

STUDY SELECTION Sixty-two studies with high (level 1-3) methodological quality that evaluated the accuracy of clinical findings for identifying difficult intubation were reviewed.

DATA EXTRACTION AND SYNTHESIS Two authors independently abstracted data. Bivariate random-effects meta-analyses were used to calculate summary positive likelihood ratios across studies or univariate random-effects models when bivariate models failed to converge.

RESULTS Among the 62 high-quality studies involving 33 559 patients, 10% (95% CI, 8.2%-12%) of patients were difficult to intubate. The physical examination findings that best predicted a difficult intubation included a grade of class 3 on the upper lip bite test (lower incisors cannot extend to reach the upper lip; positive likelihood ratio, 14 [95% CI, 8.9-22]; specificity, 0.96 [95% CI, 0.93-0.97]), shorter hyomental distance (range of <3-5.5 cm; positive likelihood ratio, 6.4 [95% CI, 4.1-10]; specificity, 0.97 [95% CI, 0.94-0.98]), retrognathia (mandible measuring <9 cm from the angle of the jaw to the tip of the chin or subjectively short; positive likelihood ratio, 6.0 [95% CI, 3.1-11]; specificity, 0.98 [95% CI, 0.90-1.0]), and a combination of physical findings based on the Wilson score (positive likelihood ratio, 9.1 [95% CI, 5.1-16]; specificity, 0.95 [95% CI, 0.90-0.98]). The widely used modified Mallampati score (≥ 3) had a positive likelihood ratio of 4.1 (95% CI, 3.0-5.6; specificity, 0.87 [95% CI, 0.81-0.91]).

CONCLUSIONS AND RELEVANCE Although several simple clinical findings are useful for predicting a higher likelihood of difficult endotracheal intubation, no clinical finding reliably excludes a difficult intubation. An abnormal upper lip bite test, which is easily assessed by clinicians, raises the probability of difficult intubation from 10% to greater than 60% for the average-risk patient.

JAMA. 2019;321(5):493-503. doi:10.1001/jama.2018.21413

+ Video and Supplemental content

+ CME Quiz at
jamanetwork.com/learning
 and **CME Questions** page 508

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Damon C. Scales, MD, PhD, 2075 Bayview Ave, Room D108, Toronto, ON M4N 3M5, Canada (damon.scales@sunnybrook.ca).

Section Editors: David L. Simel, MD, MHS, Durham Veterans Affairs Medical Center and Duke University Medical Center, Durham, NC; Edward H. Livingston, MD, Deputy Editor.

Clinical Scenario

Case 1

A previously healthy 27-year-old woman was scheduled for elective cholecystectomy. Examination of her airway demonstrated a modified Mallampati score of 2; however, she was unable to bite her upper lip with her lower incisors.

Case 2

A 68-year-old woman with pneumonia was seen on the medical ward for worsening hypoxemia and the need for mechanical ventilation. On initial inspection she was obese, breathing at a respiratory rate of 40 breaths per minute, and had retrognathia. She was confused and uncooperative. Her compromised clinical condition precluded a thorough oropharyngeal and neck examination.

Will endotracheal intubation be difficult in these patients?

Why Is This Question Important?

Endotracheal intubation is often required for major surgical procedures and for respiratory support in critically ill patients. Recognizing a potentially difficult intubation can help clinicians prepare for complications by getting assistance from clinicians with airway training and having advanced airway management equipment available.¹⁻³

Failure to predict and plan for a patient **with a difficult airway is** the most important factor contributing to the catastrophic **"cannot intubate, cannot ventilate"** scenario.^{2,4} Although this occurs **in fewer than 1/5000 elective general anesthetic procedures and requires surgical airway rescue** in fewer than **1/50 000 cases**, these situations can result in major complications associated with long-term morbidity and account for **25% of anesthesia-related deaths**.^{2,4-6} The ability to predict which patients have a high risk of difficult intubation may reduce the risk for "cannot intubate, cannot ventilate" scenarios. This study was performed to identify patient history, clinical features, and bedside tests predictive for difficult intubation.

What Is a Difficult Intubation?

The 2 most common definitions of difficult intubation used in published studies are the **Cormack-Lehane grading scale**^{7,8} and the Intubation Difficulty Scale.⁹ The Cormack-Lehane grading scale describes how visible the vocal cords are during laryngoscopy, ranging from 1 (full view of vocal cords) to 4 (cannot see the epiglottis). **The Intubation Difficulty Scale is a** scoring system that accounts for **the Cormack-Lehane grading scale and other** features including the **number of intubation attempts**, the **clinicians involved**, advanced **airway adjuncts** used, the need for increased **lifting force**, the requirement for **external laryngeal pressure**, and whether the **vocal cords are open** or **closed** during laryngoscopy.

Components of the Airway Examination

The American Society of Anesthesiologists has identified **11 anatomical features** that should be assessed prior to general anesthesia and

Key Points

Question Which risk factors and physical findings can help predict difficult endotracheal intubation?

Findings In this systematic review, several physical findings increased the likelihood of difficult intubation. The **best predictors** were an **inability to bite the upper lip with the lower incisors**, a **short hyomental distance**, **retrognathia**, or a **combination of findings based on the Wilson score**. **No risk factor** or physical finding **consistently ruled out** a potentially difficult intubation.

Meaning Although a variety of tests are helpful in identifying a potentially difficult intubation, the **inability to bite the upper lip with the lower teeth** was the **best predictor**.

endotracheal intubation to help identify patients at risk for difficult intubation.¹⁰ However, even during emergency situations when a thorough assessment of the oropharynx and neck is not feasible, experienced observers might recognize anthropometric features that increase the likelihood of a difficult intubation. Recognition of the potential for a difficult intubation is the purpose of this review. The factors associated with difficult bag-mask ventilation or establishment of an emergent surgical airway were not reviewed.

History

A comprehensive history begins with a review of **prior intubations** and factors that may have altered the anatomy of the airway or neck. Examples include previous neck injury, radiation, surgery, or medical conditions including ankylosing spondylitis and diabetes. A history or symptoms suggestive of **obstructive sleep apnea** should be elicited because this syndrome is **associated with upper airway obstruction during sedation**.^{11,12}

Physical Examination

Several physical signs and bedside tests have been assessed for predicting difficult endotracheal intubation.^{13,14} Physical examination should involve inspection of the oropharynx using a penlight and estimates of anthropometric distances and mobility of the cervical spine and mandible.

Upper Lip Bite Test, Retrognathia, and Mandibular Protrusion


The upper lip bite test assesses mandibular range of movement by asking patients to bite their upper lip with their lower incisors. The results of this test are described in terms of **3 grading classifications: class 1**, the lower incisors extend beyond the vermilion border of the upper lip; **class 2**, the lower incisors bite the lip but cannot extend above the vermilion border; and **class 3**, the lower incisors **cannot bite the upper lip at all**.¹⁵ (Figure 1). **Among patients without teeth**, the upper lip bite test can be replaced with the **upper lip catch test**, which evaluates whether the lower lip can be raised to cover the vermilion border of the upper lip.¹⁶

Retrognathia refers to either the mandible measuring less than **9 cm from the angle of the jaw to the tip of the chin** or the subjective appearance of a short mandible. **Mandibular protrusion** assesses the range of movement of the mandible by asking patients to move their **lower teeth past their upper teeth**.

Thyromental and Hyomental Distances

The thyromental distance is the distance between the upper-most border of the thyroid cartilage and the mentum measured with the

Figure 1. Upper Lip Bite Test



The **upper lip bite test** is performed by asking patients to bite their upper lip with their lower incisors. The results are classified as follows: class 1, the lower incisors extend beyond the vermilion border of the upper lip; class 2, the lower

incisors bite the lip but cannot extend above the vermilion border; and **class 3**, the lower incisors cannot bite the upper lip at all.

neck **extended**.¹⁷ Similarly, the **hyomental** distance is the distance between the hyoid bone and the mentum (Figure 2).¹⁸ Comparing the thyromental or hyomental distance with a **patient's height** can **adjust for the difference** in these measures in relation to a patient's overall size. For example, a **thyromental** distance of **6 cm** in a patient who is **200 cm tall** is **more predictive** of difficult intubation than a thyromental distance of **6 cm** in a patient who is **160 cm** tall. Using a tape measure reduces interobserver variability, but in practice clinicians may use the patient's fingerbreadths or their own as a surrogate.

Cervical Spine Mobility and Sternomental Distance

The degree of cervical spine flexion and extension as well as any neurological symptoms that arise from neck movement should be assessed prior to intubation.¹⁹ Patients **with better cervical spine mobility** will have a **longer sternomental distance**, which is the distance between the upper border of the sternum and the tip of the jaw with the neck fully extended.²⁰ Poor cervical spine mobility can make intubation more difficult.

Interincisor Gap and Modified Mallampati Score

The **maximal distance between the upper and lower incisors** is the **mouth opening capacity**, referred to as the interincisor gap. The modified Mallampati score is a grading system used to rate the visibility of the structures in the oropharynx, including the uvula, faucial pillars, and soft palate when the mouth is opened. The **original** Mallampati score used a **3-level** classification system²¹; however, a **modified** Mallampati score is more commonly used and has a **4-level** system to classify which oropharyngeal structures are visible (Figure 3).²²

Palm Print Sign and Prayer Sign

Among patients with diabetes, collagen glycosylation can lead to limitations in mobility of the small joints of the **hands** and other ana-

tomical regions, including the **cervical spine**. One method to measure mobility of the interphalangeal joints is the palm print sign.²³ An impression of the dominant hand is stamped on a piece of paper and graded based on the proportion of the hand seen on the paper. Another method is the **prayer sign**,²⁴ which tests if the patient is able to **press his or her 2 palms together**.

Abnormal Teeth

Abnormalities in teeth can make it difficult to visualize the vocal cords.^{25,26} This includes subjective assessments of prominent, loose, or missing teeth.²⁷⁻³⁰

Composite Scores


Combining findings from the history and physical examinations can improve the predictive accuracy for difficult intubation. Composite scores include the El Ganzouri score¹⁷ (which incorporates the modified Mallampati score, interincisor gap, thyromental distance, and cervical spine mobility) and the **Wilson Score** (which incorporates **weight, cervical spine mobility, jaw mobility, degree of retrognathia, and the appearance of the incisors**) (Table 1).³⁰

Methods

Search Strategy

We conducted a computerized search using OVID versions of MEDLINE (1946-June 2018) and EMBASE Classic and EMBASE (1947-June 2018). The search strategy used was (difficult\$ or awkward\$ or challeng\$ or fail\$ or ease or easy or success\$ or complicat\$ or uncomplicat\$) adj2 (intubat\$ or airway or laryngoscop\$), limited to human. We also searched the reference lists of included studies. Each citation was reviewed in duplicate by 2 of the reviewers, with full-text retrieval of any citation that either reviewer considered potentially relevant for assessing risk factors or clinical tests

Figure 2. Measurements for Thyromental, Sternomental, and Hyomental Distances



The thyromental distance is the distance between the thyroid notch and the mentum measured with the neck extended. The hyomental distance is the distance between the hyoid bone and the mentum and can be measured with the head in the neutral position (Table 2) or with the neck extended (eTable 4 in the Supplement).

One method to assess cervical spine mobility involves placing a marker on the

forehead in the vertical plane when the neck is fully extended, and then measuring the change in marker orientation as the neck is brought into full flexion. Patients with better cervical spine mobility have a longer sternomental distance, which is the distance between the upper border of the sternum and the mentum with the neck fully extended.


that predict difficult intubation. Additional details appear in eFigure 1 in the Supplement.

Study Selection

Two reviewers independently assessed the full text of each retrieved citation. The following criteria were used for study inclusion: (1) cohort study design and a minimum of 10 patients or a clinical trial, (2) population of adult patients aged 18 years or older,

(3) intervention of endotracheal intubation performed by direct laryngoscopy, (4) any element of medical history or physical examination, and (5) outcome of difficult laryngoscopy or endotracheal intubation that was measured in the same manner for all patients in each individual study. We excluded studies that were not written in English, were review articles, or if we were unable to abstract relevant data. Studies that used advanced airway devices for endotracheal intubation also were excluded.

Figure 3. Modified Mallampati Score and Mouth-Opening Capacity



The interincisor gap is the maximal distance between the upper and lower incisors. The modified Mallampati classification assesses the visibility of oropharyngeal structures when the mouth is maximally opened and

tongue protruded: class 1, soft palate, fauces, uvula, pillars; class 2, soft palate, fauces, uvula; class 3, soft palate, base of uvula; and class 4, soft palate not visible at all.²²

Table 1. Wilson Score

Parameter	Score (Range, 0-10)		
	0	1	2
Weight, kg	<90	90-110	>110
Cervical spine mobility	>90°	90°	<90°
Impaired jaw mobility	Interincisor gap ≥ 5 cm or able to protrude lower teeth past the upper teeth	Interincisor gap <5 cm and only able to protrude lower teeth to meet upper teeth	Interincisor gap <5 cm and unable to protrude lower teeth to meet upper teeth
Retrognathia	Normal	Moderate	Severe
Prominent incisors	Normal	Moderate	Severe

Assessment of Study Quality

Study quality was summarized using a quality checklist designed for the Rational Clinical Examination series.³¹ Level 1 studies included 100 or more consecutive patients, clinical features were assessed and categorized independently, and the person who intubated the patient was blinded to the assessment. Level 2 studies included less than 100 patients. Level 3 studies included nonconsecutive patients. The study characteristics of level 1 to 3 studies appear in eTable 1 in the Supplement. We excluded level 4 and 5 studies. All studies were graded independently and in duplicate.

Statistical Methods

Two reviewers independently abstracted data to construct 2×2 tables for each risk factor and clinical test. Disagreements were ar-

bitrated and resolved by a third reviewer. The 2×2 tables were used to calculate sensitivity, specificity, and positive and negative likelihood ratios (LRs). We summarized the sensitivities, specificities, and LRs using a bivariate model³² when 3 or more studies were available for each topic. When bivariate random-effects models failed to converge, we used a random-effects generic inverse variance method on (1) the logit scale for sensitivity and specificity and (2) the log scale for the LR. In Table 2, we highlight the results of risk factors and clinical tests that were derived from 3 or more studies and had a summary positive LR of 3 or greater or a summary negative LR less than 0.33 and corresponding 95% CI that exclude 1.0.

When there were only 2 studies for a risk factor or clinical test, the results appear as a range in the Supplement. When the predictive test was described only in a single study, the results

Table 2. Summary of Diagnostic Accuracy for Clinical Tests of Difficult Intubation Evaluated in 3 or More Studies

Predictor	Type of Analysis	No. of Patients	No. of Studies	Threshold	Sensitivity (95% CI)	Specificity (95% CI)	Positive LR (95% CI)	Negative LR (95% CI)
Risk factor: snoring ³³⁻³⁵	Bivariate	3866	3	History of snoring	0.43 (0.34-0.53)	0.87 (0.71-0.95)	3.4 (1.6-7.3)	0.65 (0.58-0.72)
Clinical Tests								
Upper lip bite test grading classification								
Class 3 ^{15,36-47}	Bivariate	5005	13	Inability of lower incisors to reach lower border of upper lip	0.60 (0.42-0.76)	0.96 (0.93-0.97)	14 (8.9-22)	0.42 (0.27-0.65)
Class 2 or 3 ^{15,33,36-50}	Bivariate	8091	17	Inability of lower incisors to reach upper lip or vermillion border of upper lip	0.60 (0.46-0.73)	0.95 (0.91-0.97)	12 (6.9-20)	0.42 (0.30-0.59)
Wilson score ⁵¹⁻⁵⁸	Bivariate	6520	8	≥2 used by 7 studies; ≥3 used by 1 study	0.43 (0.26-0.62)	0.95 (0.90-0.98)	9.1 (5.1-16)	0.60 (0.44-0.82)
Hyomental distance (measured in neutral position) ^{18,42,59}	Univariate	1245	3	Range, <3-≤5.5 cm	0.20 (0.11-0.34)	0.97 (0.94-0.98)	6.4 (4.1-10)	0.84 (0.73-0.96)
Retrogathia ^{25,34,35,40,42}	Univariate	4017	5	Mandible <9 cm ^a or subjectively short	0.19 (0.07-0.42)	0.98 (0.90-1.0)	6.0 (3.1-11)	0.85 (0.76-0.94)
Impaired mandibular protrusion ^{25,35,53,60-63}	Bivariate	4229	7	Cannot move lower teeth past upper teeth	0.25 (0.06-0.63)	0.95 (0.86-0.99)	5.5 (2.1-15)	0.78 (0.54-1.1)
Ratio of height to thyromental distance ^{36,40,44,47,48,54}	Bivariate	3497	6	Range, ≥17-≥25	0.69 (0.57-0.78)	0.87 (0.67-0.95)	5.2 (1.9-14)	0.36 (0.25-0.52)
Impaired neck mobility ^{23,25,26,33-35,40,47,60,64-66}	Bivariate	8061	12	Significant variability from subjective assessment to <30°-90° of flexion or extension	0.28 (0.13-0.51)	0.93 (0.85-0.97)	4.2 (1.9-9.5)	0.77 (0.60-0.99)
Sternomental distance ^{20,33,40,43,50,52,53,60,63,67-72}	Bivariate	6187	15	Range, <12-15 cm	0.41 (0.27-0.57)	0.90 (0.83-0.94)	4.1 (2.7-6.1)	0.65 (0.52-0.82)
Modified Mallampati score ^{15,16,18,23,25,26,33,35-38,41,44,46,47,49-57,59,60,62-66,69-84}	Bivariate	23396	47	Score ≥3	0.55 (0.48-0.62)	0.87 (0.81-0.91)	4.1 (3.0-5.6)	0.52 (0.45-0.60)
Impaired mouth opening ^{20,25,33,35,40,43,44,47,50,53,59-61,63,64,73,75,76}	Bivariate	9549	18	Interincisor gap <2-5 cm	0.36 (0.20-0.56)	0.90 (0.80-0.95)	3.6 (2.1-6.1)	0.71 (0.55-0.92)
Thyromental distance ^{18,23,25,33,40,43,45-47,50,52-54,60,63-66,69-73,75,76,85}	Bivariate	10596	26	Range, <4-≤7 cm	0.45 (0.36-0.55)	0.86 (0.80-0.91)	3.3 (2.4-4.4)	0.63 (0.55-0.73)
Palm print ^{23,26,86,87}	Univariate	695	4	Based on scoring system from each article	0.77 (0.22-0.98)	0.84 (0.55-0.96)	3.0 (1.9-4.7)	0.28 (0.08-0.97)

Abbreviation: LR, likelihood ratio.

^a Measured from the angle of the jaw to the tip of the chin.

appear as a point estimate and 95% CI (eTable 2 in the [Supplement](#)). We summarized the pooled incidence of difficult endotracheal intubation on the logit scale using a random-effects generic inverse variance method.

Because standard measures of between-study statistical heterogeneity are not available from bivariate random-effects models, we assessed the consistency of the results using the following sensitivity analyses: (1) restricting the analyses to studies that had a minimum of 30 difficult intubations; (2) excluding higher-risk populations (ie, obstetrical patients, head and neck surgeries, etc) from the analyses; (3) restricting the analyses to studies that used the Cormack-Lehane grading scale as the definition for difficult intubation; and (4) restricting the analyses to studies that fell within first and third quartile of incidence of difficult intubation (ie, 5.7%-15%). For the sensitivity analyses, we calculated summary point estimates and 95% CIs for sensitivities, specificities, and LRs using the same approach as the primary analysis, but restricted the analyses to predictors that could be summarized using bivariate random-effects models.

When at least 10 studies were available for the same predictor, we evaluated for publication bias that might have favored findings with higher diagnostic accuracy using a weighted regression of the logarithm of the diagnostic odds ratio on the inverse root of the effective sample size.⁸⁸ We used SAS version 9.4 (SAS Institute Inc) for the bivariate models and R version 3.4.0 (R Foundation for Statistical Computing) for the univariate analysis.

Results

After removal of duplicate studies, the search retrieved 12 394 articles and 62 studies (N = 33 559 patients) met criteria of level 1, 2, or 3 (eTable 1 and eFigure 1 in the [Supplement](#)). All studies that were level 1, 2, or 3 were operating room investigations, and some of these studies were restricted to specific patient populations such as obstetric (4 studies),^{36,51,67,68} patients with diabetes (2 studies),^{23,86} obese patients (1 study),⁸⁹ or those undergoing head and neck surgery (3 studies).^{25,64,73}

Incidence of Difficult Intubation

The overall proportion of patients having a [difficult intubation](#) was [10%](#) (95% CI, 8.2%-12%). Difficult intubation was most commonly defined as [a Cormack-Lehane grade of 3 or 4](#) (47 studies). Other definitions included the Cormack-Lehane grade with additional requirements (such as the number of intubation attempts, time, or use of bougie; 6 studies), percentage of glottis open (n = 1 study), an Intubation Difficulty Scale score greater than 5 (3 studies), or a minimum intubation time requirement or number of attempts (5 studies) to achieve successful endotracheal intubation.

Risk Factors for Difficult Intubation

[A history of difficult intubation](#) (2 studies) was the [risk factor most predictive](#) for a difficult intubation (positive LR range, 16-19; negative LR range, 0.72-0.82).^{52,85} Other risk factors included [snoring](#) (3 studies; positive LR, [3.4](#) [95% CI, 1.6-7.3]; negative LR, 0.65 [95% CI, 0.58-0.72]),³³⁻³⁵ [difficulty with bag-mask ventilation prior to intubation](#) (1 study; positive LR, 3.5 [95% CI, 2.6-4.7]; negative LR, 0.67 [95% CI, 0.55-0.80]),⁶⁰ and overweight or [obesity](#) (defined as a

[body mass index >27-35](#)) (5 studies; positive LR, 2.2 [95% CI, 1.6-3.1]; negative LR, 0.70 [95% CI, 0.46-1.1]).^{23,25,33,34,52} Compared with women, men were slightly more difficult to intubate (21 studies; positive LR, 1.2 [95% CI, 1.0-1.3]; negative LR, 0.87 [95% CI, 0.76-0.99]).^{18,20,26,33-35,37-39,48,52,53,59-62,69,74-76,85} (Table 2 and eTables 2 and 3 in the [Supplement](#)).

Accuracy of Clinical Examination

Upper Lip Bite Test, Retrognathia, and Mandibular Protrusion

[The upper lip bite test](#) (class 3, an inability to bite any part of the upper lip with the lower incisors) [strongly predicted a difficult intubation](#) (13 studies; positive LR, [14](#) [95% CI, 8.9-22]), whereas the ability to extend the teeth above the lower border of the upper lip was predictive of a reduced risk of difficult intubation (negative LR, 0.42 [95% CI, 0.27-0.65]).^{15,36-47} When including both class 2 and 3 upper lip bite test as a positive test, the results were similar (17 studies; positive LR, 12 [95% CI, 6.9-20]; negative LR, 0.42 [95% CI, 0.30-0.59]).^{15,33,36-50} The upper lip catch test, which is used in people with edentulism, had slightly lower predictive accuracy in a single study (positive LR, 7.2 [95% CI, 4.8-11]; negative LR, 0.28 [95% CI, 0.10-0.74]).¹⁶

[Retrognathia](#) (ie, a receding chin [2 studies] or chin length <9 cm [3 studies]) was a [good predictor](#) of difficult intubation (positive LR, 6.0 [95% CI, 3.1-11]; negative LR, 0.85 [95% CI, 0.76-0.94]).^{25,34,35,40,42} Impaired mandibular protrusion (defined as an inability to bring the lower teeth to the upper teeth [2 studies] or past the upper teeth [4 studies]; 1 study defined it as low protraction of lower jaw) was also a useful predictor (positive LR, 5.5 [95% CI, 2.1-15]; negative LR, 0.78 [95% CI, 0.54-1.1]).^{25,35,53,60-63} (Table 2 and eTable 4 in the [Supplement](#)).

Ratio of Height to Thyromental or Hyomental Distance in a Neutral Neck Position vs Neck Extension

A [high ratio of height to thyromental distance](#) (6 studies; thresholds ranging from ≥ 17 to ≥ 25) [was predictive of a difficult intubation](#) (positive LR, 5.2 [95% CI, 1.9-14]) and a [lower ratio made difficult intubation less likely](#) (negative LR, 0.36 [95% CI, 0.25-0.52]).^{36,40,44,47,48,54} (Table 2 and eTable 4 in the [Supplement](#)). A normal ratio of the hyomental distance measured when the neck is extended compared with when the neck is in a neutral position (1 study; normal is ≥ 1.2) was useful in identifying patients who had an easier intubation (negative LR, 0.19 [95% CI, 0.07-0.56]).¹⁸ (eTables 2 and 4 in the [Supplement](#)).

Thyromental and Hyomental Distance

[A shorter thyromental distance](#) (thresholds ranging from <4-7 cm; 26 studies) [increased the likelihood of a difficult intubation](#) (positive LR, 3.3 [95% CI, 2.4-4.4]), whereas a longer thyromental distance made a difficult intubation less likely (negative LR, 0.63 [95% CI, 0.55-0.73]).^{18,23,25,33,40,43,45-47,50,52-54,60,63-66,69-73,75,76,85} A shorter [hyomental distance](#) (thresholds ranging from <3-5.5 cm; 3 studies) also was helpful [in predicting difficult intubation](#) (positive LR, 6.4 [95% CI, 4.1-10]; negative LR, 0.84 [95% CI, 0.73-0.96]).^{18,42,59} (Table 2 and eTables 2 and 4 in the [Supplement](#)).

Cervical Spine Mobility and Sternomental Distance

The approach to assessing neck mobility (12 studies) was variable. Definitions included total neck extension of less than 80° (4 studies) or 90° (1 study), atlantooccipital extension of less than

35° (2 studies), or other definitions (5 studies).^{23,25,26,33-35,40,47,60,64-66} Overall, the presence of impaired neck mobility had modest predictive accuracy (positive LR, 4.2 [95% CI, 1.9-9.5]; negative LR, 0.77 [95% CI, 0.60-0.99]). Sternomental distance (thresholds ranging from <12-15 cm; 15 studies) provided similar results (positive LR, 4.1 [95% CI, 2.7-6.1]; negative LR, 0.65 [95% CI, 0.52-0.82])^{20,33,40,43,50,52,53,60,63,67-72} (Table 2 and eTable 4 in the Supplement).

Impaired Mouth Opening

A short interincisor gap (thresholds ranging from <2-5 cm; 18 studies) had moderate accuracy for predicting a difficult intubation (positive LR, 3.6 [95% CI, 2.1-6.1]; negative LR, 0.71 [95% CI, 0.55-0.92])^{20,25,33,35,40,43,44,47,50,53,59-61,63,64,73,75,76} (Table 2 and eTable 4 in the Supplement).

Modified Mallampati Score

The modified Mallampati score was the most frequently assessed clinical test in our analysis (47 studies).^{15,16,18,23,25,26,33,35-38,41,44,46,47,49-57,59,60,62-66,69-84} A modified Mallampati score of 3 or 4 had moderate accuracy for predicting a difficult intubation (positive LR, 4.1 [95% CI, 3.0-5.6]). However, a lower Mallampati score (1 or 2) did not rule out a difficult intubation (negative LR, 0.52 [95% CI, 0.45-0.60]; Table 2 and eTable 4 in the Supplement).

Palm Print Sign and Prayer Sign

A positive palm print test result (4 studies) was modestly predictive of a difficult intubation (positive LR, 3.0 [95% CI, 1.9-4.7]), whereas a normal test result made a difficult intubation less likely (negative LR, 0.28 [95% CI, 0.08-0.97])^{23,26,86,87} (Table 2 and eTable 4 in the Supplement). The prayer sign (defined as no contact between the fourth and fifth metacarpals; 1 study) provided similar results (positive LR, 4.9 [95% CI, 2.8-8.7]; negative LR, 0.75 [95% CI, 0.67-0.84])⁷⁹ (eTables 2 and 4 in the Supplement).

Accuracy of Composite Scores

The Wilson score (8 studies) was the only composite score evaluated in multiple studies in our primary analysis.⁵¹⁻⁵⁸ A Wilson score (≥ 2 in 7 studies and ≥ 3 in 1 study) was strongly predictive of a difficult intubation (positive LR, 9.1 [95% CI, 5.1-16]), but a lower score did not exclude difficulty (negative LR, 0.60 [95% CI, 0.44-0.82]) (Table 2 and eTable 5 in the Supplement). A combination of the modified Mallampati score, thyromental distance, anatomical abnormality, and cervical mobility (ie, M-TAC score; 1 study) score of 4 or greater increased the likelihood of a difficult intubation (positive LR, 6.7 [95% CI, 5.3-8.5]), whereas a score of less than 4 was useful for excluding difficult intubation (negative LR, 0.04 [95% CI, 0.01-0.17]; eTables 2 and 5 in the Supplement).⁷⁸

In addition to composite measures, investigators have assessed the usefulness of combining various clinical tests. Particularly useful combinations for ruling in difficult intubation included thyromental distance and modified Mallampati score⁶⁰ (positive LR, 6.0 [95% CI, 3.1-12]); thyromental distance and impaired mandibular protrusion⁶⁰ (positive LR, 7.3 [95% CI, 3.2-17]); thyromental distance, sternomental distance, and modified Mallampati score⁶⁹ (positive LR, 120 [95% CI, 7.0-2000]; eTables 2 and 4 in the Supplement).

Sensitivity Analyses

For each of the 4 sensitivity analyses of the bivariate results, the point estimates did not qualitatively change the interpretation of the primary results and the 95% CIs tended to be wider given the smaller sample sizes (eFigure 2 in the Supplement).

Publication Bias

For topics with at least 10 studies, there was no evidence of publication bias (ie, suspected unpublished studies with diagnostic odds ratios closer to 1 vs the summary diagnostic odds ratio of published studies) for any of the tests, including sternomental distance ($P = .07$), impaired mouth opening ($P = .71$), impaired neck mobility ($P = .65$), modified Mallampati score ($P = .48$), sex being male ($P = .83$), thyromental distance ($P = .20$), and grade of class 3 on the upper lip bite test ($P = .21$).

Discussion

An evidence-based approach to predict difficult airway situations should help identify patients who are more likely to be difficult to intubate. Sixty-two high-quality studies were found investigating the accuracy of various risk factors and physical examination findings to predict difficult intubation. The strongest risk factor for difficult intubation is a prior history of difficult intubation; however, the absence of this finding does not rule out difficult intubation. The best bedside test for predicting difficult intubation was the upper lip bite test. Other tests with modest accuracy include low thyromental distance, retrognathia, and impaired mandibular protrusion. The Wilson score was the most widely studied composite score and when the score was 2 or greater, it was predictive of a difficult intubation (Table 1). No clinical tests reliably excluded all cases of difficult intubation.

Limitations

First, there was significant variability in the reference standard used among the studies to identify a difficult airway. The Cormack-Lehane grading scale was the most commonly used definition, but it only identifies a difficult view of the vocal cords during direct laryngoscopy rather than a difficult tracheal intubation. Studies that use the number of intubation attempts are vulnerable to differences in clinician ability. Nevertheless, in clinical practice, these definitions are commonly used.

Second, some predictors such as retrognathia and impaired spine mobility require subjective assessments and may be more vulnerable to interobserver variability. There was also significant variation among the studies in how the predictors were defined, thresholds for the various measurements, and in clinician ability.

Third, all level 1 to 3 studies included in this review were conducted in the operating room, which limits applicability to emergency situations.⁴ Predictors for difficult intubations in nonemergency situations may still be predictive for emergency situations; however, assessing patients for the risk factors may not be feasible if patients are clinically unstable or unable to follow simple instructions. We restricted our analysis to studies that had independent assessments of predictors and outcomes to minimize bias, but this led to the exclusion of large studies in emergency situations, like the MACHOCA score study.⁹⁰

Fourth, our analysis considered the predictors independently of each other; however, patients may have several factors that increase the risk of difficult intubation.

Fifth, **contemporary airway management** is **less reliant** on direct **laryngoscopy** because there is now greater use of extraglottic airway devices, **video laryngoscopy**, and advanced airway techniques.^{10,91,92}

Scenario Resolution

Case 1

Reflecting the prevalence of difficult intubation, this patient's pretest probability of difficult intubation was 10%. Her modified Mallampati score of 2 (negative LR, 0.52) did not suggest she would be difficult to intubate. However, her upper lip bite test grade was class 3 and that grade is associated with a higher likelihood of difficulty (positive LR, 14). The posttest probability of difficulty was 60% based on the upper lip bite test. A video laryngoscope and bougie were made available in the operating room and a second anesthesiologist was present during the intubation attempt. Even though the anesthesiologist's view of the vocal cords on direct laryngoscopy was a Cormack-Lehane grade of 3, the endotracheal intubation was successful on the first attempt.

Case 2

The patient's pretest probability of difficult intubation was 10%. Based on the cursory physical examination (obese; positive LR, 2.2) and retrognathia (positive LR, 6.0), it was estimated that her posttest probability of a difficult intubation was between 25% and 40%. The patient was transferred to the intensive care unit and a member of the anesthesiology department was called to assist with a plan for the intubation using video laryngoscopy with topical xylocaine and minimal sedation. The patient was intubated successfully on the first attempt with a Cormack-Lehane grade of 2 for the view of the larynx.

Clinical Bottom Line

Several individual physical examination findings are predictive but do not reliably exclude the likelihood for a difficult intubation. The most accurate individual bedside clinical assessment is the easily performed upper lip bite test. Given the **prevalence of a difficult intubation of 10%**, the **inability to bite the upper lip with the lower incisors raises the probability** of experiencing a difficult intubation to **more than 60%**. Other individual tests that are helpful include hyomental distance, retrognathia, and impaired mandibular protrusion. The Wilson score is also helpful for predicting which patients will have a difficult intubation.

ARTICLE INFORMATION

Accepted for Publication: December 13, 2018.

Author Affiliations: Department of Medicine, Sinai Health System, Toronto, Ontario, Canada (Detsky); Interdepartmental Division of Critical Care Medicine and Department of Medicine, University of Toronto, Toronto, Ontario, Canada (Detsky, Adhikari, Friedrich, Scales); Institute for Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada (Jivraj, Adhikari, Wijeyesundera, Scales); Department of Anesthesia, University of Toronto, Toronto, Ontario, Canada (Jivraj, Wijeyesundera); Department of Critical Care Medicine, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada (Adhikari, Pinto, Scales); Departments of Critical Care Medicine and Medicine, St Michael's Hospital, Toronto, Ontario, Canada (Friedrich); Li Ka Shing Knowledge Institute, St Michael's Hospital, Toronto, Ontario, Canada (Friedrich, Wijeyesundera); Department of Medicine, Duke University School of Medicine, Durham, North Carolina (Simel); Durham Veterans Affairs Medical Center, Durham, North Carolina (Simel); Department of Anesthesia and Pain Management, University Health Network, Toronto, Ontario, Canada (Wijeyesundera).

Author Contributions: Drs Detsky and Scales had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Detsky, Adhikari, Friedrich, Wijeyesundera, Scales.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Detsky, Jivraj, Simel, Scales.

Critical revision of the manuscript for important intellectual content: Jivraj, Adhikari, Friedrich, Pinto, Simel, Wijeyesundera, Scales.

Statistical analysis: Detsky, Pinto, Simel, Scales.

Administrative, technical, or material support: Jivraj, Scales.

Supervision: Adhikari, Simel, Scales.

Conflict of Interest Disclosures: Dr Simel reported receiving honoraria for contributions to JAMAEvidence.com; and is supported by the Durham Center of Innovation to Accelerate Discovery and Practice Transformation (CIN 13-410) at the Durham VA Health Care System. Dr Wijeyesundera is supported in part by a New Investigator Award from the Canadian Institutes of Health Research and an Excellence in Research Award from the Department of Anesthesia at the University of Toronto. No other disclosures were reported.

Additional Contributions: We acknowledge Jamie Spiegelman, MD (Humber River Hospital, Toronto, Ontario, Canada), for help with a preliminary search and related data abstraction and we thank Daniel Nishijima, MD (University of California-Davis School of Medicine), Karen Welty-Wolf, MD (Durham Veterans Affairs Medical Center and Duke University), and Jonathan Mark, MD (Durham Veterans Affairs Medical Center and Duke University, Durham, NC) for helpful comments on an earlier version of the manuscript. None of those acknowledged were compensated for contributing.

REFERENCES

- Caplan RA, Posner KL, Ward RJ, Cheney FW. Adverse respiratory events in anesthesia: a closed claims analysis. *Anesthesiology*. 1990;72(5):828-831. doi:10.1097/0000542-199005000-00010
- Cook TM, Woodall N, Frerk C; Fourth National Audit Project. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society, part 1: anaesthesia. *Br J Anaesth*. 2011;106(5):617-631. doi:10.1093/bja/aer058
- Law JA, Broemling N, Cooper RM, et al; Canadian Airway Focus Group. The difficult airway with recommendations for management—part 1—difficult tracheal intubation encountered in an unconscious/induced patient. *Can J Anaesth*. 2013; 60(11):1089-1118. doi:10.1007/s12630-013-0019-3
- Cook TM, Woodall N, Harper J, Benger J; Fourth National Audit Project. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society, part 2: intensive care and emergency departments. *Br J Anaesth*. 2011;106(5):632-642.
- Cook TM, MacDougall-Davis SR. Complications and failure of airway management. *Br J Anaesth*. 2012;109(suppl 1):i68-i85. doi:10.1093/bja/aes393
- Nagaro T, Yorozuya T, Sotani M, et al. Survey of patients whose lungs could not be ventilated and whose trachea could not be intubated in university hospitals in Japan. *J Anesth*. 2003;17(4):232-240.
- Cormack RS, Lehane J. Difficult tracheal intubation in obstetrics. *Anaesthesia*. 1984;39(11):1105-1111. doi:10.1111/j.1365-2044.1984.tb08932.x
- Yentis SM, Lee DJ. Evaluation of an improved scoring system for the grading of direct laryngoscopy. *Anaesthesia*. 1998;53(11):1041-1044. doi:10.1046/j.1365-2044.1998.00605.x
- Adnet F, Borron SW, Racine SX, et al. The Intubation Difficulty Scale (IDS): proposal and evaluation of a new score characterizing the complexity of endotracheal intubation. *Anesthesiology*. 1997;87(6):1290-1297.
- Apfelbaum JL, Hagberg CA, Caplan RA, et al; American Society of Anesthesiologists Task Force on Management of the Difficult Airway. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of

- the Difficult Airway. *Anesthesiology*. 2013;118(2): 251-270. doi:10.1097/ALN.0b013e31827773b2
11. Myers KA, Mrkobrada M, Simel DL. Does this patient have obstructive sleep apnea?: the Rational Clinical Examination systematic review. *JAMA*. 2013;310(7):731-741. doi:10.1001/jama.2013.276185
 12. Joshi GP, Ankichetty SP, Gan TJ, Chung F. Society for Ambulatory Anesthesia consensus statement on preoperative selection of adult patients with obstructive sleep apnea scheduled for ambulatory surgery. *Anesth Analg*. 2012;115(5): 1060-1068. doi:10.1213/ANE.0b013e318269cfd7
 13. Roth D, Pace NL, Lee A, et al. Airway physical examination tests for detection of difficult airway management in apparently normal adult patients. *Cochrane Database Syst Rev*. 2018;5(5):CD008874.
 14. Vannucci A, Cavallone LF. Bedside predictors of difficult intubation: a systematic review. *Minerva Anesthesiol*. 2016;82(1):69-83.
 15. Khan ZH, Kashfi A, Ebrahimkhani E. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *Anesth Analg*. 2003;96(2):595-599.
 16. Khan ZH, Arbabi S, Yekaninejad MS, Khan RH. Application of the upper lip catch test for airway evaluation in edentulous patients: an observational study. *Saudi J Anaesth*. 2014;8(1):73-77. doi:10.4103/1658-354X.125942
 17. el-Ganzouri AR, McCarthy RJ, Tuman KJ, Tanck EN, Ivankovich AD. Preoperative airway assessment: predictive value of a multivariate risk index. *Anesth Analg*. 1996;82(6):1197-1204.
 18. Huh J, Shin HY, Kim SH, Yoon TK, Kim DK. Diagnostic predictor of difficult laryngoscopy: the hyomental distance ratio. *Anesth Analg*. 2009;108(2):544-548. doi:10.1213/ane.0b013e3181818fc347
 19. Mashour GA, Stallmer ML, Kheterpal S, Shanks A. Predictors of difficult intubation in patients with cervical spine limitations. *J Neurosurg Anesthesiol*. 2008;20(2):110-115. doi:10.1097/ANA.0b013e318166dd00
 20. Savva D. Prediction of difficult tracheal intubation. *Br J Anaesth*. 1994;73(2):149-153. doi:10.1093/bja/73.2.149
 21. Mallampati SR, Gatt SP, Gugino LD, et al. A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J*. 1985;32(4):429-434. doi:10.1007/BF03011357
 22. Samsoon GL, Young JR. Difficult tracheal intubation: a retrospective study. *Anaesthesia*. 1987;42(5):487-490. doi:10.1111/j.1365-2044.1987.tb04039.x
 23. Nadal JL, Fernandez BG, Escobar IC, Black M, Rosenblatt WH. The palm print as a sensitive predictor of difficult laryngoscopy in diabetics. *Acta Anaesthesiol Scand*. 1998;42(2):199-203. doi:10.1111/j.1399-6576.1998.tb05109.x
 24. Erden V, Basaranoglu G, Delatiglu H, Hamzaoglu NS. Relationship of difficult laryngoscopy to long-term non-insulin-dependent diabetes and hand abnormality detected using the 'prayer sign'. *Br J Anaesth*. 2003;91(1):159-160. doi:10.1093/bja/aeg583
 25. Ayuso MA, Sala X, Luis M, Carbó JM. Predicting difficult orotracheal intubation in pharyngo-laryngeal disease: preliminary results of a composite index. *Can J Anaesth*. 2003;50(1): 81-85. doi:10.1007/BF03020193
 26. Mahmoodpoor A, Soleimanpour H, Nia KS, et al. Sensitivity of palm print, modified Mallampati score and 3-3-2 rule in prediction of difficult intubation. *Int J Prev Med*. 2013;4(9):1063-1069.
 27. Ezri T, Weisenberg M, Khazin V, et al. Difficult laryngoscopy: incidence and predictors in patients undergoing coronary artery bypass surgery versus general surgery patients. *J Cardiothorac Vasc Anesth*. 2003;17(3):321-324. doi:10.1016/S1053-0770(03)00052-1
 28. Juvin P, Lavaut E, Dupont H, et al. Difficult tracheal intubation is more common in obese than in lean patients. *Anesth Analg*. 2003;97(2):595-600.
 29. Nath G, Sekar M. Predicting difficult intubation—a comprehensive scoring system. *Anaesth Intensive Care*. 1997;25(5):482-486.
 30. Wilson ME, Spiegelhalter D, Robertson JA, Lesser P. Predicting difficult intubation. *Br J Anaesth*. 1988;61(2):211-216. doi:10.1093/bja/61.2.211
 31. Simel D. Update: primer on precision and accuracy. In: Simel D, ed. *The Rational Clinical Examination*. New York, NY: McGraw Hill; 2009.
 32. Takwoingi Y, Deeks JJ, MetaDAS: a SAS macro for meta-analysis of diagnostic accuracy studies. In: User Guide, version 1.3; 2010. <https://methods.cochrane.org/sdt/software-meta-analysis-dta-studies>. Accessed December 20, 2018.
 33. Chhina AK, Jain R, Gautam PL, Garg J, Singh N, Grewal A. Formulation of a multivariate predictive model for difficult intubation: a double blinded prospective study. *J Anaesthesiol Clin Pharmacol*. 2018;34(1):62-67.
 34. Hanouz J-L, Bonnet V, Buléon C, et al. Comparison of the Mallampati classification in sitting and supine position to predict difficult tracheal intubation: a prospective observational cohort study. *Anesth Analg*. 2018;126(1):161-169. doi:10.1213/ANE.0000000000002108
 35. Prakash S, Kumar A, Bhandari S, Mullick P, Singh R, Gogia AR. Difficult laryngoscopy and intubation in the Indian population: an assessment of anatomical and clinical risk factors. *Indian J Anaesth*. 2013;57(6):569-575. doi:10.4103/0019-5049.123329
 36. Honarmand A, Safavi MR. Prediction of difficult laryngoscopy in obstetric patients scheduled for Caesarean delivery. *Eur J Anaesthesiol*. 2008;25(9): 714-720. doi:10.1017/S026502150800433X
 37. Ali MA, Qamar-ul-Hoda M, Samad K. Comparison of upper lip bite test with Mallampati test in the prediction of difficult intubation at a tertiary care hospital of Pakistan. *J Pak Med Assoc*. 2012;62(10):1012-1015.
 38. Eberhart LH, Arndt C, Cierpka T, Schwaneckamp J, Wulf H, Putzke C. The reliability and validity of the upper lip bite test compared with the Mallampati classification to predict difficult laryngoscopy: an external prospective evaluation. *Anesth Analg*. 2005;101(1):284-289.
 39. Shah AA, Rafique K, Islam M. Can difficult intubation be accurately predicted using upper lip bite test? *J Postgrad Med Inst*. 2014;28(3):282-287.
 40. Badheka JP, Doshi PM, Vyas AM, Kacha NJ, Parmar VS. Comparison of upper lip bite test and ratio of height to thyromental distance with other airway assessment tests for predicting difficult endotracheal intubation. *Indian J Crit Care Med*. 2016;20(1):3-8. doi:10.4103/0972-5229.173678
 41. Hester CE, Dietrich SA, White SW, Secrest JA, Lindgren KR, Smith T. A comparison of preoperative airway assessment techniques: the modified Mallampati and the upper lip bite test. *AANA J*. 2007;75(3):177-182.
 42. Khan ZH, Maleki A, Makarem J, Mohammadi M, Khan RH, Zandieh A. A comparison of the upper lip bite test with hyomental/thyrosternal distances and mandible length in predicting difficulty in intubation: a prospective study. *Indian J Anaesth*. 2011;55(1):43-46. doi:10.4103/0019-5049.76603
 43. Khan ZH, Mohammadi M, Rasouli MR, Farrokhnia F, Khan RH. The diagnostic value of the upper lip bite test combined with sternomental distance, thyromental distance, and interincisor distance for prediction of easy laryngoscopy and intubation: a prospective study. *Anesth Analg*. 2009;109(3):822-824. doi:10.1213/ane.0b013e3181af7f0d
 44. Min JJ, Kim G, Kim E, Lee J-H. The diagnostic validity of clinical airway assessments for predicting difficult laryngoscopy using a grey zone approach. *J Int Med Res*. 2016;44(4):893-904. doi:10.1177/0300060516642647
 45. Salimi A, Farzanegan B, Rastegarpour A, Kolahi AA. Comparison of the upper lip bite test with measurement of thyromental distance for prediction of difficult intubations. *Acta Anaesthesiol Taiwan*. 2008;46(2):61-65. doi:10.1016/S1875-4597(08)60027-2
 46. Selvi O, Kahraman T, Senturk O, Tulgar S, Serifsoy E, Ozer Z. Evaluation of the reliability of preoperative descriptive airway assessment tests in prediction of the Cormack-Lehane score: a prospective randomized clinical study. *J Clin Anesth*. 2017;36:21-26. doi:10.1016/j.jclinane.2016.08.006
 47. Shah PJ, Dubey KP, Yadav JP. Predictive value of upper lip bite test and ratio of height to thyromental distance compared to other multivariate airway assessment tests for difficult laryngoscopy in apparently normal patients. *J Anaesthesiol Clin Pharmacol*. 2013;29(2):191-195.
 48. Balakrishnan KP, Chockalingam PA. Ethnicity and upper airway measurements: a study in South Indian population. *Indian J Anaesth*. 2017;61(8): 622-628. doi:10.4103/ijana.IJA_247_17
 49. Mahmoodpoor A, Soleimanpour H, Golzari SE, et al. Determination of the diagnostic value of the Modified Mallampati score, upper lip bite test and facial angle in predicting difficult intubation: a prospective descriptive study. *J Clin Anesth*. 2017; 37:99-102. doi:10.1016/j.jclinane.2016.12.010
 50. Mehta T, Jayaprakash J, Shah V. Diagnostic value of different screening tests in isolation or combination for predicting difficult intubation: a prospective study. *Indian J Anaesth*. 2014;58(6): 754-757. doi:10.4103/0019-5049.147176
 51. Gupta S, Pareek S, Dulara SC. Comparison of two methods for predicting difficult intubation in obstetric patients. *Middle East J Anaesthesiol*. 2003;17(2):275-285.
 52. Kim WH, Ahn HJ, Lee CJ, et al. Neck circumference to thyromental distance ratio: a new predictor of difficult intubation in obese patients. *Br J Anaesth*. 2011;106(5):743-748. doi:10.1093/bja/aer024

53. Topcu I, Ovali GY, Yentur EA, Kefi A, Tuncyurek O, Pabuscu Y. Clinical and radiologic evaluations for predicting difficult tracheal intubation. *Acta Anaesth Italica*. 2006;57:285-300.
54. Krishna HM, Munisha A, Dali JS, Prashant R, Dua CK. Prediction of difficult laryngoscopy in indian population: role of ratio of patient's height to thyromental distance. *J Anaesthesiol Clin Pharmacol*. 2005;21(3):257-260.
55. Oates JD, Macleod AD, Oates PD, Pearsall FJ, Howie JC, Murray GD. Comparison of two methods for predicting difficult intubation. *Br J Anaesth*. 1991;66(3):305-309. doi:10.1093/bja/66.3.305
56. Siddiqi R, Kazi WA. Predicting difficult intubation—a comparison between Mallampati classification and Wilson risk-sum. *J Coll Physicians Surg Pak*. 2005;15(5):253-256.
57. Yamamoto K, Tsubokawa T, Shibata K, Ohmura S, Nitta S, Kobayashi T. Predicting difficult intubation with indirect laryngoscopy. *Anesthesiology*. 1997;86(2):316-321. doi:10.1097/00000542-199702000-00007
58. Guo Y, Feng Y, Liang H, Zhang R, Cai X, Pan X. Role of flexible fiberoptic laryngoscopy in predicting difficult intubation. *Minerva Anestesiol*. 2018;84(3):337-345.
59. Yu T, Wang B, Jin XJ, et al. Predicting difficult airways: 3-3-2 rule or 3-3 rule? *Ir J Med Sci*. 2015;184(3):677-683. doi:10.1007/s11845-015-1276-7
60. Yildiz TS, Korkmaz F, Solak M, et al. Prediction of difficult tracheal intubation in Turkish patients: a multi-center methodological study. *Eur J Anaesthesiol*. 2007;24(12):1034-1040. doi:10.1017/S026502150700052X
61. Sahin SH, Yilmaz A, Gunday I, et al. Using temporomandibular joint mobility to predict difficult tracheal intubation. *J Anesth*. 2011;25(3):457-461. doi:10.1007/s00540-011-1126-3
62. Ul Haq MI, Ullah H. Comparison of Mallampati test with lower jaw protrusion maneuver in predicting difficult laryngoscopy and intubation. *J Anaesthesiol Clin Pharmacol*. 2013;29(3):313-317.
63. Khatiwada S, Bhattarai B, Pokharel K, Acharya R. Prediction of difficult airway among patients requiring endotracheal intubation in a tertiary care hospital in Eastern Nepal. *JNMA J Nepal Med Assoc*. 2017;56(207):314-318. doi:10.31729/jnma.2918
64. Kuriakose R, Mathew A, Koshy CR. Screening tests for predicting difficult endotracheal intubation—a clinical assessment in facio-oro-maxillary and neck malignancy patients. *J Anaesthesiol Clin Pharmacol*. 2003;19(1):37-44.
65. Tse JC, Rimm EB, Hussain A. Predicting difficult endotracheal intubation in surgical patients scheduled for general anesthesia: a prospective blind study. *Anesth Analg*. 1995;81(2):254-258.
66. Wong SH, Hung CT. Prevalence and prediction of difficult intubation in Chinese women. *Anaesth Intensive Care*. 1999;27(1):49-52.
67. Al Ramadhani S, Mohamed LA, Rocke DA, Gouws E. Sternomental distance as the sole predictor of difficult laryngoscopy in obstetric anaesthesia [published correction appears in *Br J Anaesth*. 1996;77(5):701]. *Br J Anaesth*. 1996;77(3):312-316. doi:10.1093/bja/77.3.312
68. Eiamcharoenwit J, Itthisompaiaboon N, Limpawattana P, Suwanpratheep A, Siriussawakul A. The performance of neck circumference and other airway assessment tests for the prediction of difficult intubation in obese parturients undergoing cesarean delivery. *Int J Obstet Anesth*. 2017;31:45-50. doi:10.1016/j.ijoa.2017.01.011
69. Iohom G, Ronayne M, Cunningham AJ. Prediction of difficult tracheal intubation. *Eur J Anaesthesiol*. 2003;20(1):31-36. doi:10.1097/00003643-200301000-00006
70. Badhe VK, Deogaonkar SG, Tambe MV, Singla A, Shidhaye RV. Clinical comparison of five different predictor tests for difficult intubation. *Anaesth Pain Intensive Care*. 2014;18(1):31-37.
71. Etezadi F, Ahangari A, Shokri H, et al. Thyromental height: a new clinical test for prediction of difficult laryngoscopy. *Anesth Analg*. 2013;117(6):1347-1351. doi:10.1213/ANE.0b013e3182a8c734
72. Reddy PB, Punetha P, Chalam KS. Ultrasonography—a viable tool for airway assessment. *Indian J Anaesth*. 2016;60(11):807-813. doi:10.4103/0019-5049.193660
73. Han Y-Z, Tian Y, Xu M, et al. Neck circumference to inter-incisor gap ratio: a new predictor of difficult laryngoscopy in cervical spondylosis patients. *BMC Anesthesiol*. 2017;17(1):55. doi:10.1186/s12871-017-0346-y
74. Choi JW, Kim JA, Kim HK, Oh MS, Kim DK. Chest anteroposterior diameter affects difficulty of laryngoscopy for non-morbidly obese patients. *J Anesth*. 2013;27(4):563-568. doi:10.1007/s00540-013-1572-1
75. Pinto J, Cordeiro L, Pereira C, Gama R, Fernandes HL, Assunção J. Predicting difficult laryngoscopy using ultrasound measurement of distance from skin to epiglottis. *J Crit Care*. 2016;33:26-31. doi:10.1016/j.jccr.2016.01.029
76. Yao W, Wang B. Can tongue thickness measured by ultrasonography predict difficult tracheal intubation? *Br J Anaesth*. 2017;118(4):601-609. doi:10.1093/bja/aeo051
77. Aktas S, Atalay YO, Tugrul M. Predictive value of bedside tests for difficult intubations. *Eur Rev Med Pharmacol Sci*. 2015;19(9):1595-1599.
78. Ambesh SP, Singh N, Rao PB, Gupta D, Singh PK, Singh U. A combination of the modified Mallampati score, thyromental distance, anatomical abnormality, and cervical mobility (M-TAC) predicts difficult laryngoscopy better than Mallampati classification. *Acta Anaesthesiol Taiwan*. 2013;51(2):58-62. doi:10.1016/j.aat.2013.06.005
79. Baig MM, Khan FH. To compare the accuracy of prayer's sign and Mallampatti test in predicting difficult intubation in diabetic patients. *J Pak Med Assoc*. 2014;64(8):879-883.
80. Bindra A, Prabhakar H, Singh GP, Ali Z, Singhal V. Is the modified Mallampati test performed in supine position a reliable predictor of difficult tracheal intubation? *J Anesth*. 2010;24(3):482-485. doi:10.1007/s00540-010-0905-6
81. Kamalipour H, Bagheri M, Kamali K, Taleie A, Yarmohammadi H. Lateral neck radiography for prediction of difficult orotracheal intubation. *Eur J Anaesthesiol*. 2005;22(9):689-693. doi:10.1017/S0265021505001146
82. Khan ZH, Eskandari S, Yekaninejad MS. A comparison of the Mallampati test in supine and upright positions with and without phonation in predicting difficult laryngoscopy and intubation: a prospective study. *J Anaesthesiol Clin Pharmacol*. 2015;31(2):207-211. doi:10.4103/0970-9185.155150
83. Savva D. Sternomental distance—a useful predictor of difficult intubation in patients with cervical spine disease? *Anaesthesia*. 1996;51(3):284-285. doi:10.1111/j.1365-2044.1996.tb13653.x
84. Gao YX, Song YB, Gu ZJ, et al. Video versus direct laryngoscopy on successful first-pass endotracheal intubation in ICU patients. *World J Emerg Med*. 2018;9(2):99-104.
85. Qudaisat IY, Al-Ghanem SM. Short thyromental distance is a surrogate for inadequate head extension, rather than small submandibular space, when indicating possible difficult direct laryngoscopy. *Eur J Anaesthesiol*. 2011;28(8):600-606. doi:10.1097/EJA.0b013e328347cdd9
86. Reissell E, Orko R, Maunukela EL, Lindgren L. Predictability of difficult laryngoscopy in patients with long-term diabetes mellitus. *Anaesthesia*. 1990;45(12):1024-1027. doi:10.1111/j.1365-2044.1990.tb14879.x
87. Sachdeva KP, Singh A, Kathuria S, et al. Prediction of difficult laryngoscopy in diabetics by palm print and interphalangeal gap. *J Anaesthesiol Clin Pharmacol*. 2005;21(3):261-264.
88. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol*. 2005;58(9):882-893. doi:10.1016/j.jclinepi.2005.01.016
89. Toshniwal G, McKelvey GM, Wang H. STOP-Bang and prediction of difficult airway in obese patients. *J Clin Anesth*. 2014;26(5):360-367.
90. De Jong A, Molinari N, Terzi N, et al; AzuRéa Network for the Frida-Réa Study Group. Early identification of patients at risk for difficult intubation in the intensive care unit: development and validation of the MACOCHA score in a multicenter cohort study. *Am J Respir Crit Care Med*. 2013;187(8):832-839.
91. Frerk C, Mitchell VS, McNarry AF, et al; Difficult Airway Society Intubation Guidelines Working Group. Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults. *Br J Anaesth*. 2015;115(6):827-848.
92. Law JA, Broemling N, Cooper RM, et al; Canadian Airway Focus Group. The difficult airway with recommendations for management—part 2—the anticipated difficult airway. *Can J Anaesth*. 2013;60(11):1119-1138. doi:10.1007/s12630-013-0020-x