20. Kurnatowski P, Putynski L, Lapienis M, Kowalska B: Neurocognitive abilities in children with adenotonsillar hypertrophy. Int J Ped Otorhinolaryngol 2006; 70:419-24 21. Nixon GM, Kermack AS, McGregor CD, Davis GM, Manoukian JJ, Brown KA, Brouillette RT: Sleep and breathing on the first night after adenotonsillectomy for obstructive sleep apnea. Ped Pulmonology 2005; 39:332-8

Anesthesiology 2006; 105:647-8

Copyright © 2006, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

# **Regional Anesthesia, Intraneural Injection, and** Nerve Injury

### Beyond the Epineurium

NERVE damage after regional anesthesia is appropriately regarded as a major complication and, when the injury is severe, may take weeks or even months to recover completely.<sup>1,2</sup> There are many possible causes for such injuries.<sup>3,4</sup> These include stretching, compression, ischemia, surgical trauma, and local anesthetic toxicity.<sup>5-7</sup> One causative factor that has been the subject of intense discussion involves the direct intraneural injection of local anesthetics. The deleterious effect of such injections was demonstrated by Selander et al.<sup>3</sup> nearly 30 yr ago. Since that time, we have been advised to avoid direct contact between the needle and nerve and to think of the epineurium as a barrier that we should not cross. One consequence of this advice has been a move away from "seeking paresthesias" during the performance of blocks and the use of electrical stimulation and evoked motor responses to estimate proximity to the nerve. However, in this issue of ANESTHESIOLOGY, Dr. Bigeleisen<sup>8</sup> has challenged the idea that intraneural injection is uniformly damaging and is to be avoided at all costs.

In this study, videography and ultrasonography were used to assess local anesthetic distribution when axillary brachial plexus block was performed according to his usual practice, which was seeking paresthesia by needle manipulation. When paresthesia was established, 2–3 ml local anesthetic was administered. If the injection appeared intraneurally, the needle was withdrawn until it appeared outside the nerve, the injection was continued, and the block was completed. The patients were checked 6 months later for the occurrence of neuropathy. The results of the study were surprising: 22 of 26 patients (85%) had nerve puncture of at least one nerve, and 21 of 26 patients (81%) had an intraneural injection

This Editorial View accompanies the following article: Bigeleisen PE: Nerve puncture and apparent intraneural injection during ultrasound-guided axillary block do not invariably result in neurologic injury. ANESTHESIOLOGY 2006; 105:779-83. of at least one nerve. Assessment 6 months later showed no clinical evidence of nerve damage. Two important new considerations emerge from this investigation: First, intraneural injection of local anesthetic, at least in a small volume, does not seem to result in nerve damage, and second, performance of the paresthesia technique does result in frequent intraneural injection.

The belief that administration of local anesthetic inside the epineurium uniformly results in nerve damage should be reconsidered in view of Bigeleisen's results.<sup>8</sup> The study showed that injection of local anesthetic (2-3 ml) inside the epineurium does not result in severe nerve damage. Some minor, transient neurologic symptoms may have occurred between block performance and neurologic assessment at 6 months and may have been unrecognized, but the occurrence of severe nerve damage would most likely have been brought to the attention of the author or detected by the surgeon.

Ultrasonographic resolution does not allow us to differentiate between an injection into the subepineurium or subperineurium. The perineurium, in contrast to the epineurium, is a tough and resistant tissue withstanding very high pressure.<sup>3</sup> The ability to expand the nerve, as shown in figure 2B in Bigeleisen's article,<sup>8</sup> suggests that the needle lies in a compliant space between the epineurium and perineurium. However, the main issue coming from this investigation is that the barrier that should not be penetrated to avoid severe neural damage is likely the perineurium. The next question, which cannot be answered, is how much volume can be placed in this space until the pressure increases and adversely affects the blood supply. A study will be needed to clarify this question.

A more recent study by Hadzic *et al.*<sup>9</sup> further evaluated the consequences of either subepineurium or subperineurium injections in dogs. In this study, the authors placed the tip of the needle under microscopic control either around the epineurium or intraneurally by piercing the epineurium. In the control group, injection pressures were low (< 4 psi) in all animals. In the intraneural group, the authors were able to distinguish two subgroups: one with a moderately increased injection pressure, and the other with a very high injection pressure (25-45 psi). After the dogs awakened from general an-

Accepted for publication June 21, 2006. The author is not supported by, nor maintains any financial interest in, any commercial activity that may be associated with the topic of this article.

esthesia, motor function returned to normal within 3 h in all animals, except for those with very high injection pressures. In this subgroup, severe and persistent motor deficits were recorded, with varying degrees of damage to the neural architecture. The weakness of this investigation resides in the absence of proof that the needle was effectively placed subperineurally. Similar studies using electronic microscopy and injection of dye should be able to confirm these suppositions.

Another interesting point made by Bigeleisen<sup>8</sup> is the apparent high frequency of subepineurium local anesthetic deposition when using the paresthesia technique for performing peripheral nerve block. These findings give support to those promoting the use of electrical nerve stimulation. However, the volume of the injection that enters subepineurally is unknown in this context, but this observation may explain the greater incidence of minor neurologic symptoms observed by some authors using the paresthesia technique<sup>10</sup> and the observation that the incidence of severe neurologic complication is not greater when using the paresthesia technique compared with electrical nerve stimulation.<sup>10</sup> Therefore, intraneural injection may not cause severe neurologic deficits and might be explained by the relatively good tolerance of low or moderate volume of local anesthetics between the epineurium and perineurium. Another interesting finding reported by Bigeleisen<sup>8</sup> is the heterogeneous description of symptoms observed after eliciting paresthesia-a phenomenon that is poorly explained. It must be emphasized that the possibility to "contact" the nerve without eliciting any paresthesia or dysesthesia may occur. This phenomenon has occasionally been reported in the literature.<sup>11</sup>

This investigation has some limitations. Detractors will criticize the current study because of its relatively small sample size and the lack of any neurologic assessment until 6 months after the injection. It has been demonstrated that most peripheral nerve injuries are transient after regional blocks and resolve within a few weeks after the injury.<sup>1,2</sup> However, this study raises pertinent questions about the

importance of penetrating the epineurium and nerve damage during regional anesthesia.

In summary, for neurologic complications from regional anesthesia, the belief that the epineurium as the last barrier should be balanced—local anesthetics should be injected outside of it—but we should recognize that some local anesthetics can be injected without uniformly damaging the nerve. Evidence is growing that the key barrier is the perineurium. The work performed by Bigeleisen<sup>8</sup> contributes to this understanding. However, this new information should not yet change our clinical practice: Nerves should be treated with care, and the basic rule not to inject local anesthetics into the nerve remains.

Alain Borgeat, M.D., Department of Anesthesiology, Orthopedic University Hospital Balgrist, Zurich, Switzerland. alain.borgeat@ balgrist.ch

#### References

1. Borgeat A, Ekatodramis G, Kalberer F, Benz C: Acute and nonacute complications associated with interscalene block and shoulder surgery: A prospective study. ANESTHESIOLOGY 2001; 95:875-80

2. Borgeat A, Dullenkopf A, Ekatodramis G, Nagy L: Evaluation of the lateral modified approach for continuous interscalene block after shoulder surgery. ANESTHESIOLOGY 2003; 99:436-42

3. Selander D, Brattsand R, Lundborg G, Nordborg C, Olsson Y: Local anesthetics: Importance of mode of application, concentration and adrenaline for the appearance of nerve lesions. An experimental study of axonal degeneration and barrier damage after intrafascicular injection or topical application of bupivacaine (Marcain). Acta Anaesthesiol Scand 1979; 23:127-36

4. Moore DC, Mulroy MF, Thompson GE: Peripheral nerve damage and regional anaesthesia. Br J Anaesth 1994; 73:435-6

 Pitman MI, Nainzadeh N, Ergas E, Springer S: The use of somatosensory evoked potentials for detection of neuropraxia during shoulder arthroscopy. Arthroscopy 1988; 4:250-5

6. Ikeda K, Yokoyama M, Tomita K, Tanaka S: Vulnerability of the gradually elongated nerve to compression injury. Hand Surg 2001; 6:29-35

7. Selander D: Peripheral nerve injury after regional anesthesia, Complications of Regional Anesthesia, 1st edition. Edited by Finuncane BT. Philadelphia, Churchill Livingstone, 1999, pp 105-15

8. Bigeleisen P: Nerve puncture and apparent intraneural injection during ultrasound-guided axillary block do not invariably result in neurologic injury. ANESTHESIOLOGY 2006; 105:779-83

9. Hadzic A, Dilberovic F, Shah S, Kulenovic A, Kapur E, Zaciragic A, Cosovic E, Vuckovic I, Divanovic KA, Mornjakovic Z, Thys DM, Santos AC: Combination of intraneural injection and high injection pressure leads to fascicular injury and neurologic deficits in dogs. Reg Anesth Pain Med 2004; 29:417-23

10. Selander D, Edshage S, Wolff T: Paresthesiae or no paresthesiae? Nerve lesions after axillary blocks. Acta Anaesthesiol Scand 1979; 23:27-33

11. Sala-Blanch X, Pomes J, Matute P, Valls-Sole J, Carrera A, Tomas X, Garcia-Diez AI: Intraneural injection during anterior approach for sciatic nerve block. ANESTHESIOLOGY 2004; 101:1027-30 Anesthesiology 2006; 105:779-83

## Nerve Puncture and Apparent Intraneural Injection during Ultrasound-guided Axillary Block Does Not Invariably Result in Neurologic Injury

Paul E. Bigeleisen, M.D.\*

*Background:* Nerve puncture by the block needle and intraneural injection of local anesthetic are thought to be major risk factors leading to neurologic injury after peripheral nerve blocks. In this study, the author sought to determine the needle–nerve relation and location of the injectate during ultrasound-guided axillary plexus block.

*Metbods:* Using ultrasound-guided axillary plexus block (10-MHz linear transducer, SonoSite, Bothel, WA; 22-gauge B-bevel needle, Becton Dickinson, Franklin Parks, NJ), the incidence of apparent nerve puncture and intraneural injection of local anesthetic was prospectively studied in 26 patients. To determine the onset, success rate, and any residual neurologic deficit, qualitative sensory and quantitative motor testing were performed before and 5 and 20 min after block placement. At a follow-up 6 months after the blocks, the patients were examined for any neurologic deficit.

*Results:* Twenty-two of 26 patients had nerve puncture of at least one nerve, and 21 of 26 patients had intraneural injection of at least one nerve. In the entire cohort, 72 of a total of 104 nerves had intraneural injection. Sensory and motor testing before and 6 months after the nerve injections were unchanged.

*Conclusions:* Under the conditions of this study, puncturing of the peripheral nerves and apparent intraneural injection during axillary plexus block did not lead to a neurologic injury.

ULTRASOUND is a useful aid for the performance of peripheral nerve block. Its use has resulted in a decreased incidence of vascular puncture, faster onset times, decreased dose requirements, and higher success rates.<sup>1-4</sup> The practice of ultrasound-guided nerve block is still evolving. Most practitioners continue to use a nerve stimulator even while performing the procedure under real-time viewing. There is also considerable debate about what the images mean. Most publications have shown a black (hypoechoic) local anesthetic ring around the nerve after injection is done (fig. 1). This is consistent with a nerve that is surrounded by a tissue

This article is accompanied by an Editorial View. Please see: Borgeat A: Regional anesthesia, intraneural injection, and nerve injury: Beyond the epineurium. ANESTHESIOLOGY 2006; 105:647-8. plane that is separated from the epineurium. Local anesthetic is injected into this potential space.

In the author's experience, as the needle touches the nerve, the nerve moves 1 or 2 cm before the needle pierces an anatomical structure that may be the fascia. As the needle pierces this anatomical structure/fascia, the practitioner may feel a pop, and the patient often reports a paresthesia or dysesthesia. Simultaneously, the needle seems to enter the substance of the nerve. Injection of 2-3 ml of local anesthetic usually proceeds with minimal pain and resistance. When compared with the undisturbed nerve (fig. 2A), the injected nerve seems to swell with a uniform stippled image and little or no black ring around it (fig. 2B). Some of the local anesthetic forms a black hypoechoic shadow in the nerve, and some of the local anesthetic leaks out of the nerve, forming a small hypoechoic ring around the nerve (fig. 2C). These observations are intriguing because they contradict the common assumptions that needle-nerve contact and intraneural injection are invariably associated with neurologic injury. A retrospective review of 50 ultrasoundguided blocks in the author's archive showed that one or more nerves in each patient was punctured and followed by an intraneural injection with local anesthetic. Because these patients did not experience any known sequelae, the author decided to perform a prospective study of his ultrasound-guided axillary blocks to determine the incidence of needle entry into the nerve as well as the images formed if local anesthetic was injected into the nerve. The author also studied any transient or permanent injuries to the nerves.

### Materials and Methods

After institutional review board approval at Lindsay House Surgery Center and written patient consent, 50 American Society of Anesthesiologists physical status I-III patients who were older than 17 yr were enrolled in the study. All patients had previously consented to an ultrasound-guided axillary block for surgical repair of the basal joint of the thumb. An anesthesiologist with experience performing ultrasound-guided nerve blocks conducted the study. Before the performance of the block and before sedation, each patient was given a sensory and motor examination of the musculocutaneous nerve, median nerve, radial nerve, and ulnar nerve. The sensory examination was performed as follows: musculocutane-

<sup>\*</sup> Associate Professor.

Received from the Department of Anesthesiology, Lindsay House Surgery Center at Linden Oaks, Rochester, New York. Submitted for publication December 28, 2005. Accepted for publication April 19, 2006. Supported by Sleeping Gorilla Studios, Pittsford, New York.

Address correspondence to Dr. Bigeleisen: Department of Anesthesiology, University of Pittsburgh School of Medicine, 2300 Lothrop Street, C Wing, Room C-203, Pittsburgh, Pennsylvania 15213. bigeleisenpe@upmc.edu. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.



Fig. 1. Injection of local anesthetic around peripheral nerve.

ous nerve in the distribution of the lateral antebrachial nerve, median nerve in the palmar tip of the middle finger, radial nerve on the dorsum of the wrist proximal to the incision site, ulnar nerve in the palmar surface of the fifth finger. The following modalities were used: pinprick, alcohol wipe (cold), and a soft brush (light touch). A score of 1 was given if the patient could identify the modality, and a score of 0 was given if the patient could not identify the modality.

A qualitative muscular examination using the Medical Research Council Scale<sup> $\dagger$ </sup> (5 = full strength, 0 = no movement) was conducted by asking the patient to perform the following maneuvers: musculocutaneous nerve (flex forearm), median nerve (flex distal interphalangeal joint of second finger), radial nerve (extend wrist), ulnar nerve (abduct third and fourth fingers). Any patient who had deficiencies in any of the sensory or motor tests was excluded from the study. There were no other exclusionary criteria. A quantitative muscular examination was also conducted using grip strength and a force meter. With the forearm pronated and the wrist extended, the force meter was squeezed between the index finger and thumb. With the forearm supinated, the force meter was squeezed between the index finger and thumb. These maneuvers collectively test the median (index finger flexion), radial (wrist extension), and ulnar nerves (thumb apposition).

All axillary blocks were performed with the arm abducted 120°, the shoulder externally rotated, and the forearm flexed 90°. Patients received premedication with 1–2 mg midazolam and 50–100  $\mu$ g fentanyl. A 22-gauge, short-bevel needle designed for nerve blocks (Becton Dickinson, Franklin Parks, NJ) was used for all injections. A 10-MHz linear transducer was sited in the axilla over the axillary artery. The axillary artery, vein,



Fig. 2. Intraneural puncture and injection of peripheral nerve.

pectoralis major muscle, biceps muscle, and coracobrachialis muscle were identified on ultrasound. To the best of the author's ability, the musculocutaneous, median, ulnar, and radial nerves were also identified on ultra-

sound as well. Each nerve was anesthetized in the order described above by eliciting a paresthesia or piercing the fascia around the nerve by the sensation of a pop. After a paresthesia had been elicited or pop had been felt, 2-3 ml of anesthetic mixture (2.5 mg/ml bupivacaine, 10 mg/ml lidocaine, 3  $\mu$ g/ml epinephrine) was injected. If the injection appeared intraneural, the needle was withdrawn and 2-3 ml of local anesthetic mixture was injected around the nerve. If the initial injection produced a halo, an additional 2-3 ml of local anesthetic was injected around the nerve. If neither an intraneural injection nor a halo appeared, the process was repeated one more time with the exception of the musculocutaneous nerve. In the case of the musculocutaneous nerve, if neither an intraneural injection nor a halo could be achieved on the first injection, a total of 10 ml of local anesthetic was injected into the coracobrachialis muscle. The musculocutaneous and median nerves were anesthetized using an approach superior to the axillary artery. The ulnar and radial nerves were anesthetized using an approach inferior to the artery. The total dose of local anesthetic injected was recorded for each patient.

The ultrasound sequence was recorded on a digital recorder, and a nurse recorded the clinical activities of the author and patient response during the injections with a digital video recorder. The ultrasound tape and clinical video were then spliced together using Adobe Premiere (San Jose, CA) so that both ultrasound and clinical sequences could be viewed simultaneously. A licensed sonographer, trained in musculoskeletal imaging, reviewed each spliced tape to ascertain whether the needle appeared to enter the substance of the nerve during injection. The sonographer also noted whether the nerve appeared to swell, to become surrounded by a dark ring, or both. Finally, the sonographer noted whether the patient appeared to have a paresthesia or dysesthesia during the fascial puncture or injection by studying whether the patient moved or reported tingling or pain. Any occurrence of vascular puncture or hematoma formation was also tallied.

Qualitative sensory and muscle testing was repeated on each patient using the same maneuvers as described

Table 1. Mean Sensory Block at 5 and 20 Minutes

n = 26	Mu	М	R	U
5 min				
Sharp	0.2	0.4	0.3	0.3
Light touch	0	0.5	0.3	0.2
Cold	0	0.3	0.4	0.3
20 min				
Sharp	0	0.1	0	0.1
Light touch	0	0	0	0
Cold	0	0	0	0

M = median nerve; Mu = musculocutaneous nerve; R = radial nerve; U = ulnar nerve.

#### Table 2. Mean Motor Block at 5 and 20 Minutes

n = 26	Mu	М	R	U
5 min	1.1	2.7	2.2	2.3
20 min	0.1	0.2	0.1	0.2

5 =full strength, 0 =akinetic.

M = median nerve; Mu = musculocutaneous nerve; R = radial nerve; U = ulnar nerve.

above. These tests were repeated at 5 min, and 20 minutes after the block was completed by the author. The same examination, including quantitative muscle testing, was repeated at a follow-up visit 6 months after the patient's surgery. The 6-month follow-up examinations were performed by a nurse practitioner or physical therapist who was blinded to the results of the preoperative examinations. A tally was also made of the number of patients who reported surgical pain during the surgery or in the recovery room. Statistical analysis of quantitative strength was done using a one-sided t test. A value of 0.05 was assumed to be statistically significant. All patients were contacted 24-48 h after surgery to determine whether they had residual anesthesia in their surgical limb. They were also questioned about whether they had paresthesias, dysesthesias, weakness in the surgical limb, or pain at the injection site. All patients were seen within 96 h after the surgery, at 3 weeks, and again at 6 months by the surgeon. She was asked to report any abnormalities in the patients' sensory or motor examinations to the author.

#### Results

Twenty-two patients were excluded from the study because of preoperative abnormalities in their qualitative sensory or motor examinations. Two patients were lost to follow-up. The musculocutaneous and radial nerves were the easiest to block (tables 1 and 2). None of the patients reported surgical pain during the procedure, and all of the patients completed the surgery without local anesthetic supplementation by the surgeon or the need for general anesthesia. None of the patients required analgesia in the recovery room. All 26 patients who completed the study had normal qualitative sensory and motor examinations at 6-month follow-up. There was no significant difference in quantitative muscle testing preoperatively and at 6-month follow-up (table 3). The incidence of nerve puncture, halo around the nerve, nerve swelling, halo and nerve swelling, and paresthesias or dysesthesias are listed in table 4.

Tal	ble	3.	Force	M	le	ter
-----	-----	----	-------	---	----	-----

n = 26	Supination	Pronation
Preoperatively, mean, kg 6 Months postoperatively, mean, kg	$\begin{array}{c} 5.7 \pm 0.6 \\ 5.7 \pm 0.8 \end{array}$	$\begin{array}{l} 4.4\pm0.8\\ 4.5\pm0.5\end{array}$

Anesthesiology, V 105, No 4, Oct 2006 Copyright © by the American Society of Anesthesiologists. Unauthorized reproduction of this article is prohibited.

n = 26	Puncture	Halo	Swelling	Halo and Swelling	Paresthesia or Dysesthesia
Mu	8	18	5	3	4
Μ	22	4	4	18	23
R	23	3	3	20	24
U	19	7	1	18	15

Table 4. Incidence of Nerve Puncture, Halo, Swelling, orParesthesia

M= median nerve; Mu= musculocutaneous nerve; R= radial nerve; U= ulnar nerve.

Puncture, intraneural injection, or halo around the musculocutaneous nerve did not produce any paresthesias or dysesthesias. Injection into the coracobrachialis muscle produced a paresthesia in 4 patients. Puncture of the median nerve produced a paresthesia or dysesthesia in 18 patients. In 15 patients, this paresthesia diminished during intraneural injection. In 3 patients, the paresthesia intensified during injection in to the median nerve. In 4 patients, puncture or injection of the median nerve did not produce any sensation. Halo alone of the median nerve produced a paresthesia when the nerve was initially contacted in 1 patient. Augmentation of the paresthesia occurred in this patient during injection. In 10 patients, puncture of the ulnar nerve produced a paresthesia or dysesthesia, which was diminished on intraneural injection. Halo alone produced a paresthesia in 5 patients during contact of the ulnar nerve. Three of these patients had a paresthesia or dysesthesia during injection around the ulnar nerve. Puncture of the radial nerve produced a paresthesia or dysesthesia in 21 patients. In 14 patients, this paresthesia diminished during intraneural injection. In 7 patients, the paresthesia increased with injection of the radial nerve. Halo alone produced a paresthesia or dysesthesia on radial nerve contact in 3 patients but no sensation during injection. Three patients reported tenderness at the injection site in the axilla on the day after surgery. This complication resolved by 3 weeks in all 3 patients without treatment. None of the patients reported paresthesias or dysesthesias in the distribution of the four injected nerves after surgery. The mean dose of local anesthetic used was 26  $\pm$  12 ml (range, 17-45 ml). There were no vascular punctures or hematomas.

#### Discussion

The incidence of nerve puncture was lowest for the musculocutaneous nerve. This is most likely because the musculocutaneous nerve is the smallest in diameter and therefore the most difficult to puncture. The incidence of halo or swelling alone after injection was very low for all nerves. In most cases, there was both a halo and swelling because some of the local anesthetic infiltrated the nerve itself, whereas some of the local anesthetic filled a potential space around the nerve. The absence of measurable neurologic injury (qualitative sensory and qualitative and quantitative motor testing) in any of the patients at 6-month follow-up suggests that neural puncture and or injection, *per se*, are not the immediate or most likely cause of neural injury after nerve block using the techniques described in this study. Unfortunately, the author did not have access to diagnostic electrophysiologic tools such as sensory nerve conduction thresholds, nerve conduction velocity, or electromyography, which may have added useful information to the study. It is also possible that the surgeon missed transient injuries in the early postoperative period, which may have resolved by the 6-month follow-up.

This study does not explain the cause of neural injury after nerve block, and the study was likely underpowered to elucidate the cause of neural injury because the incidence of nerve injury after neural block is so low.<sup>5</sup> The large number of patients excluded from the study because of neurologic abnormalities also decreased the robustness of the study. Other mechanisms, however, may be the cause of neural injury. Ultrasound does not have the resolution to differentiate between injection into the stroma of the nerve or a fascicle of the nerve. Intrafascicular injection may be one cause of nerve injury. High injection pressure is also thought to be one cause of neural injury.<sup>6</sup> Unfortunately, the author did not have the equipment to perform manometry. However, small aliquots of local were injected into or around the nerve; therefore, high injection pressure was less likely. In traditional nerve block techniques, the nerve is localized by paresthesia or electrolocation. When the nerve or plexus is located, many practitioners inject 30-40 ml of local anesthetic at one site. This is more likely to create a high pressure if the injection is intraneural. Finally, dysesthesia without motor deficit after a nerve block may be due to injury of the nervi nervorum, which innervate the epineurium and mesoneurium. If this is the mechanism of injury, puncture or injection of the epineurium may be the cause of injury. Again, this study does not have the spatial resolution to identify this type of injury.

Bevel type and needle gauge may also influence the incidence of nerve injury. In this study, we used a shortbevel, 22-gauge needle. One may not assume that needles of different gauge or with different bevel types, used in the same manner, will result in the same safety record. Inglis *et al.*<sup>7</sup> found that conduction deficits occurred after individual axon impalement by microelectrodes. Finally, the type and location of the nerve may influence the way ultrasound-guided injections proceed. Nerves in the axilla have little or no fascia surrounding them and are frequently septated with large amounts of stroma between the fascicles (fig. 3). A blunt needle that pierces these small nerves may be less likely to puncture a fascicle. Moreover, these small nerves are freer to swell



Fig. 3. Cross-section of peripheral nerve.

because they are not constrained by dense fascia. Finally, the cross section of a peripheral nerve is comprised of approximately 50% neurons and 50% fat and connective tissue.<sup>8,9</sup> Thus, there is a significant probability of puncturing a peripheral nerve without contacting a fascicle or damaging the neurons. Case reports describe similar scenarios. Impalement of the femoral nerve under ultrasound guidance resulted in temporary injury,<sup>10</sup> whereas the placement of a catheter within the sciatic nerve under computed tomography guidance did not cause any clinical injury.<sup>11</sup> These may be reasons why puncture and injection of small doses of local anesthetic into nerves did not cause injury. Nerve roots and nerve trunks have larger fascicles with virtually no intraneural stroma and are much more heavily invested with fascia.<sup>9</sup> Injection into these structures, even with small doses of local anesthetic, may produce higher pressures and permanent injury.

In summary, the author has used ultrasound to study, in real time, the injection of local anesthetic during nerve block in the axilla. Nerves were frequently punctured and infiltrated with local anesthetic. This did not seem to cause any long-term damage to the nerves under the conditions used in this study. Injection into the nerve after eliciting a paresthesia/dysesthesia produced a heterogeneous set of sensations. Sometimes the paresthesia/dysesthesia was augmented by injection, whereas in other instances, the paresthesia/dysesthesia was diminished or absent during intraneural injection. This study is underpowered to assert that intraneural injections or nerve puncture are inconsequential. However, these data and recent reports suggest that nerve puncture or intraneural injection does not inevitably lead to neurologic injury.

#### References

1. Marhofer P, Schrogendorfer K, Koinig H, Kapral S, Weinstabl C, Mayer N: Ultrasonographic guidance improves sensory block and onset time of three-inone blocks. Anesth Analg 1997; 85:854-7

2. Marhofer P, Schrogendorfer K, Wallner T, Konig H, Mayer N, Kapral S: Ultrasonic guidance reduces the amount of local anesthetic for 3-in-1 blocks. Reg Anesth Pain Med 1998; 23:584–8

 Sandu NS, Capan LM: Ultrasound-guided infractavicular brachial plexus block. Br J Anaesth 2002; 89:254–9

4. Chan VWS: Applying ultrasound imaging to interscalene brachial plexus block. Reg Anesth Pain Med 2003; 28:340-3

5. Auroy Y, Benhamou D, Bargues L, Ecoffey C, Falissard B, Mercier F, Bouaziz H, Kamran S: Major complications of regional anesthesia in France: The SOS, Regional Anesthesia Hotline Service. ANESTHESIOLOGY 2002; 97:1274-80

6. Hadzic A, Dilberovic F, Shah S, Kulenovic A, Kapur E, Zaciragic A, Cosovic E, Vuckovic I, Divanovic KA, Mornjakovic Z, Thys DM, Santos AC: Combination of intraneural injections and high injection pressure leads to fascicular injury and neurological deficits in dogs. Reg Anesth Pain Med 2004; 29:417-23

7. Inglis JT, Leeper JB, Wilson LR, Gandevia SC, Burke D: The development of conduction block in single human axons following a focal nerve injury. J Physiol 1988; 518:127-33

8. Reina MA, Lopez A, De Andres JA: Adipose tissue within peripheral nerves: Study of human sciatic nerve [in Spanish]. Rev Esp Anestesiol Reanim 2002; 49:397-402

9. Sunderland S: Nerve Injuries and Their Repair: A Critical Appraisal. New York, Churchill Livingstone, 1991, pp 31-40

10. Schafhalter-Zoppoth I, Gray AT: Inadvertent femoral nerve impalement and intraneural injection visualized by ultrasound. Anesth Analg 2004; 99:627-8

11. Sala-Blanch X, Pomes J, Matute P, Valls-Sole J, Carrera A, Tomas X, Garcia-Diez A: Intraneural injection during anterior approach for sciatic nerve block. ANESTHESIOLOGY 2004; 101:1027-30