

COMMENTARY

Early goal-directed therapy: what do we do now?

Mitchell M Levy

See related research by Gu *et al.*, <http://ccforum.com/content/18/5/570>

Abstract

The meta-analysis of early goal-directed therapy (EGDT) by Gu and colleagues in the previous issue of *Critical Care* adds to the ongoing controversy about the value of EGDT for resuscitating patients with severe sepsis and septic shock. The results of the ProCESS (protocolized care for early septic shock) and ARISE (Australasian resuscitation in sepsis evaluation) trials failed to demonstrate any benefit of EGDT or protocolized resuscitation when compared with 'usual care'. The questions are the following: What is 'usual' care? What is 'real world' care? Do the results of a robust and well-conducted randomized controlled trial - in which many patients may be excluded for a variety of reasons - reflect the care given to patients on a daily basis in our emergency departments and intensive care units? Of course, there are no obvious answers to these questions, and many clinicians look forward to managing these patients without protocols. For now, the data do seem to support the management of patients with septic shock without mandated central lines or protocols. Does this mean we should go back to the era of 'do whatever you want'? No consensus exists among clinicians regarding optimal hemodynamic monitoring, and to date no method has been proven to be superior. Given the amount of fluids given prior to randomization in the ProCESS and ARISE trials, 'usual care' appears to now include aggressive, early fluid resuscitation with at least 20 mL/kg of crystalloid and rapid administration of appropriate antibiotics. Certainly, this reflects the impact of the original trial by Rivers and colleagues and the broad-based implementation of the Surviving Sepsis Campaign Guidelines and bundles. If this continues to define 'usual care', then perhaps it is no longer necessary to mandate specific protocols for resuscitation, as it appears that standard sepsis management has evolved to be consistent with published protocols.

Introduction

The meta-analysis of early goal-directed therapy (EGDT) by Gu and colleagues [1] in the previous issue of *Critical Care* adds to the ongoing controversy about the value of EGDT for resuscitating patients with severe sepsis and septic shock. The results of the ProCESS (protocolized care for early septic shock) [2] and ARISE (Australasian resuscitation in sepsis evaluation) [3] trials have been interpreted by some to sound the death knell for EGDT and the value of protocol-driven resuscitation in these critically ill patients. Both of these trials failed to demonstrate any benefit of EGDT or protocolized resuscitation when compared with 'usual care'. However, aspects of the two trials provoke concerns about comparing them with the original study by Rivers and colleagues [4], including population severity, administration of fluids prior to randomization, and the overall mortality rate.

Early goal-directed therapy: interpreting the recent data

In the meta-analysis by Gu and colleagues (which included the ProCESS but not the ARISE trial), a significant survival benefit from EGDT was found in seven trials - all randomized controlled trials (RCTs) - in the subgroup of 'early' intervention trials (that is, within 6 hours). However, by the authors' own admission, the meta-analysis was flawed because of a high risk of bias due to the uncertain methodology of some of the trials. Though weakened by the fact that four of the included studies were published only in Chinese-language journals, this meta-analysis and another recently published study [5] raise questions about how to generalize the results of the ProCESS and ARISE trials.

The challenge for practicing clinicians is how to understand 'usual care' in the settings of these large RCTs. In both the ProCESS and ARISE trials, the usual care mortality was 18%. That is a remarkably low mortality for a population with septic shock. Most clinicians

Correspondence: mitchell_levy@brown.edu
Rhode Island Hospital, Brown University, 593 Eddy Street, Providence, RI 02903, USA

would ask whether this population reflects their clinical practice, given the low acuity - Acute Physiology and Chronic Health Evaluation II scores - and short intensive care unit length of stay in these two trials, and whether these results can be generalized to a sicker population seen outside the confines of an RCT in which 'real world' patients are excluded.

The questions are the following: What is 'usual' care? What is 'real world' care? Do the results of a robust and well-conducted RCT - in which many patients may be excluded for a variety of reasons - reflect the care given to patients on a daily basis in our emergency departments and intensive care units? Does a mortality rate of 18% represent a realistic goal in 2014? Are clinicians and hospitals prepared to be held to that mortality rate by the public and third-party payers? In the US, mandated public reporting of sepsis is now a reality in one state [6], and preparations are under way nationally to use sepsis measures for public reporting in just a few years. Will regulatory agencies hold clinicians to this mortality rate?

Of course, there are no obvious answers to these questions, and many clinicians look forward to managing these patients without protocols. For now, the data do seem to support the management of patients with septic shock without mandated central lines or protocols. However, almost two thirds of the patients in the two large RCTs were on pressors; therefore, the large majority of patients in both the ProCESS and ARISE trials were managed with the use of central lines.

Fortunately for patients and clinicians, the opportunity to address the question of what comprises 'usual care' will emerge over the next several years as state and national reporting takes hold [6,7]. Metrics will be collected from large populations of patients with severe sepsis and septic shock on the various modalities of monitoring available for clinicians to choose from, including physical examination. Over time, analyses of these large databases should create clarity about the use of hemodynamic monitoring and protocolized care as 'routine care' in guiding clinicians to improve outcomes of patients with this common illness. Meanwhile, clinicians and hospitals should continue to dedicate time and resources to establish systems to identify these patients early and begin treatment rapidly after identification. Most still agree that when it comes to sepsis, the earlier we treat, the better.

Conclusions

Does this mean we should go back to the era of 'do whatever you want'? No consensus exists among clinicians regarding optimal hemodynamic monitoring, and to date no method has been proven to be superior. Given the amount of fluids given prior to randomization in the ProCESS and ARISE trials, 'usual care' appears to

now include aggressive, early fluid resuscitation with at least 20 mL/kg of crystalloid and rapid administration of appropriate antibiotics. Certainly, this reflects the impact of the original trial by Rivers and colleagues [4] and the broad-based implementation of the Surviving Sepsis Campaign Guidelines and bundles [8]. If this continues to define 'usual care', then perhaps it is no longer necessary to mandate specific protocols for resuscitation, as it appears that standard sepsis management has evolved to be consistent with published protocols. That would be good news indeed for patients with severe sepsis and septic shock.

Abbreviations

ARISE: Australasian resuscitation in sepsis evaluation; EGDT: Early goal-directed therapy; ProCESS: Protocolized care for early septic shock; RCT: Randomized controlled trial.

Competing interests

The author is a founding member of the Surviving Sepsis Campaign.

Published online: 29 December 2014

References

1. Gu WJ, Wang F, Bakker J, Tang L, Liu JC: The effect of goal-directed therapy on mortality in patients with sepsis - earlier is better: a meta-analysis of randomized controlled trials. *Crit Care* 2014, **18**:570.
2. Investigators ProCESS, Yealy DM, Kellum JA, Huang DT, Barnato AE, Weissfeld LA, Pike F, Terndrup T, Wang HE, Hou PC, LoVecchio F, Filbin MR, Shapiro NI, Angus DC: A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 2014, **370**:1683-1693.
3. ARISE Investigators; ANZICS Clinical Trials Group, Peake SL, Delaney A, Bailey M, Bellomo R, Cameron PA, Cooper DJ, Higgins AM, Holdgate A, Howe BD, Webb SA, Williams P: Goal-directed resuscitation for patients with early septic shock. *N Engl J Med* 2014, **371**:1496-1506.
4. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, Peterson E, Tomlanovich M, for the Early Goal-Directed Therapy Collaborative Group: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 2001, **345**:1368-1377.
5. Levy MM, Rhodes A, Phillips GS, Townsend SR, Schorr CA, Beale R, Osborn T, Lemeshow S, Chiche JD, Artigas A, Dellinger RP: Surviving Sepsis Campaign: association between performance metrics and outcomes in a 7.5-year study. *Crit Care Med* 2014 [Epub ahead of print].
6. New York State regulations for Sepsis reporting [http://www.health.ny.gov/facilities/public_health_and_health_planning_council/meetings/2013-02-07/docs/13-01.pdf]
7. Proposed rule - Federal Register [https://www.federalregister.gov/articles/2014/05/15/2014-10067/medicare-program-hospital-inpatient-prospective-payment-systems-for-acute-care-hospitals-and-the]
8. Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, Sevransky JE, Sprung CL, Douglas IS, Jaeschke R, Osborn TM, Nunnally ME, Townsend SR, Reinhart K, Kleinpell RM, Angus DC, Deutschman CS, Machado FR, Rubenfeld GD, Webb SA, Beale RJ, Vincent JL, Moreno R: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2012. *Int Care Med* 2013, **39**:165-228.

doi:10.1186/s13054-014-0705-8

Cite this article as: Levy: Early goal-directed therapy: what do we do now? *Critical Care* 2014 **18**:705.

RESEARCH

Open Access

The effect of goal-directed therapy on mortality in patients with sepsis - earlier is better: a meta-analysis of randomized controlled trials

Wan-Jie Gu¹, Fei Wang², Jan Bakker³, Lu Tang² and Jing-Chen Liu^{1*}

Abstract

Introduction: The Surviving Sepsis Campaign guidelines recommend goal-directed therapy (GDT) for the early resuscitation of patients with sepsis. However, the findings of the ProCESS (Protocolized Care for Early Septic Shock) trial showed no benefit from GDT for reducing mortality rates in early septic shock. We performed a meta-analysis to integrate these findings with existing literature on this topic and evaluate the effect of GDT on mortality due to sepsis.

Methods: We searched the PubMed, Embase and CENTRAL (Cochrane Central Register of Controlled Trials) databases and reference lists of extracted articles. Randomized controlled trials comparing GDT with standard therapy or usual care in patients with sepsis were included. The prespecified primary outcome was overall mortality.

Results: In total, 13 trials involving 2,525 adult patients were included. GDT significantly reduced overall mortality in the random-effects model (relative risk (RR), 0.83; 95% confidence interval (CI), 0.71 to 0.96; $P = 0.01$; $I^2 = 56\%$). Predefined subgroup analysis according to the timing of GDT for resuscitation suggested that a mortality benefit was seen only in the subgroup of early GDT within the first 6 hours (seven trials; RR, 0.77; 95% CI, 0.67 to 0.89; $P = 0.0004$; $I^2 = 40\%$), but not in the subgroup with late or unclear timing of GDT (six trials; RR, 0.92; 95% CI, 0.69 to 1.24; $P = 0.59$; $I^2 = 56\%$). GDT was significantly associated with the use of dobutamine (five trials; RR, 2.71; 95% CI, 1.20 to 6.10; $P = 0.02$).

Conclusions: The results of the present meta-analysis suggest that GDT significantly reduces overall mortality in patients with sepsis, especially when initiated early. However, owing to the variable quality of the studies, strong and definitive recommendations cannot be made.

Introduction

Sepsis is a systemic response to infection, which may progress to severe sepsis and septic shock [1]. Severe sepsis and septic shock represent global problems with a high economic burden. In the United States, more than 750,000 people experience severe sepsis each year, with a short-term mortality of 20% to 30%, reaching up to 50% when shock is present [2,3]. Therefore, numerous therapeutic strategies that aimed at reducing mortality in these patients have been investigated. However, most of them have not led to significant reductions in mortality [4]. Goal-directed therapy (GDT) has been shown to substantially improve clinical outcomes in surgical patients [5]. Two important aspects of a GDT protocol include early

initiation of the therapeutic measures, together with specific (hemodynamic) targets. The Surviving Sepsis Campaign guidelines recommend GDT for the early resuscitation of patients with sepsis [1], which is based largely upon the results of the Rivers *et al.* trial, in which the researchers reported a 16% absolute reduction in mortality among patients with severe sepsis or septic shock who received early GDT compared to standard therapy [6]. However, in a recent study, this approach was challenged [7], with no benefit shown when GDT was compared to standard therapy. However, initiating therapy early rather than late in the course of critical illness remains a logical clinical goal. In the context of this situation, we systematically reviewed all trials of GDT in patients with sepsis and performed a meta-analysis, focusing on the early initiation of the protocol and its effect on mortality.

* Correspondence: jingchenliu1964@sina.cn

¹Department of Anaesthesiology, the First Affiliated Hospital, Guangxi Medical University, 22 Shuangyong Road, Nanning 530021, China
Full list of author information is available at the end of the article

Methods

Ethical approval and patient consent were not required, because we conducted a meta-analysis of previously published studies. We followed the recommendations of the *Cochrane Handbook for Systematic Reviews of Interventions* to carry out the study [8] and follow the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) statement to report our meta-analysis [9].

Search strategy

Electronic searches were conducted in the PubMed, Embase and Cochrane Central Register of Controlled Trials (CENTRAL) databases. Search terms included “goal directed,” “goal oriented,” “goal target,” “cardiac output,” “cardiac index,” “oxygen delivery,” “oxygen consumption,” “cardiac volume,” “stroke volume,” “fluid therapy,” “fluid loading,” “fluid administration,” “optimization,” “optimization,” “supranormal” and “sepsis,” “severe sepsis,” “septic shock,” “septicemia,” “septicaemia,” “pyohemia,” “pyaemia” and “pyemia.” There was no language restriction placed on the searches. Each database was searched from inception to April 2014. Additionally, reference lists in the articles chosen for inclusion, and the reference lists of previous reviews were screened to identify other potentially eligible trials.

Inclusion criteria

We included trials with the following characteristics:

1. **Population:** Adult patients with one or more of the following characteristics were eligible for inclusion: sepsis, severe sepsis or septic shock. *Adults* were defined as being of a legal age for consent in the country where the trial was conducted. Studies that included sepsis secondary to noninfectious causes were excluded.
2. **Intervention:** The intervention had to be *GDT*, defined as an explicit protocol encompassing the use of hemodynamic monitoring and manipulation of hemodynamic parameters to achieve predetermined hemodynamic endpoints.
3. **Control:** The control group had to have received standard therapy or usual care.
4. **Outcomes:** The overall mortality rate had to be the outcome measured.
5. **Type of study:** The studies had to be randomized controlled trials (RCTs).

We included studies that randomized a mixed population of critically ill patients when a septic subpopulation that met our inclusion criteria was defined; that is, the patients with sepsis constituted a subgroup of the trial population.

Data extraction

We extracted data using a standardized data collection form. Discrepancies in collected data were addressed through team consensus. The following information was extracted from each trial: first author, year of publication, number of patients (GDT and control), study population, clinical setting, goals in GDT and control groups, timing of GDT, mortality endpoint, study design (patient selection and concealment) and outcome data (overall mortality and dobutamine use).

Outcomes and definitions

The prespecified primary outcome was overall mortality. If the study authors reported mortality at one time point, we used the only data used for analysis. If the study authors reported mortality at more than one time point, we used hospital mortality preferentially. Secondary outcomes included overall mortality in the early-initiated GDT (that is, within the first 6 hours) versus GDT and dobutamine use initiated later.

Assessment of risk of bias

We used the Cochrane Collaboration tool to assess the risk of bias of individual study and with bias domains across studies [8,10]. Two investigators (WJG and FW) subjectively reviewed all studies and assigned a value of “high,” “low” or “unclear” to the following domains: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting and other bias. Trials with high risk of bias for any one or more key domains were considered to be at high risk of bias. Trials with low risk of bias for all key domains were considered to be at low risk of bias. Otherwise, they were considered to have an unclear risk of bias [10].

Statistical analysis

We estimated the relative risk (RR) with 95% confidence interval (CI) for dichotomous outcomes. Statistical heterogeneity across studies was tested by using the I^2 statistic [11,12]. Heterogeneity was suggested if the P -value was ≤ 0.10 . I^2 values of 0 to 24.9%, 25% to 49.9%, 50% to 74.9% and 75% to 100% were considered zero, low, moderate and high thresholds for statistical heterogeneity, respectively [11,12]. Clinical heterogeneity could not be excluded, so the more conservative random-effects model [13] (Mantel-Haenszel method) was used. Predefined subgroup analysis was conducted according to the timing of GDT for resuscitation (early being within the first 6 hours versus late or unclear being outside the first 6 hours or unclear timing). In addition, we performed *post hoc* subgroup analyses according to risk of bias (low versus unclear), sample size (≥ 100 versus < 100) and setting (emergency department versus

intensive care unit). We further investigated the influence of a single study on the overall pooled estimate by omitting one study in each step. The potential for bias was assessed by inspection of a funnel plot and Egger's test [14]. The results were considered statistically significant at two-sided P -values <0.05 . All statistical analyses were performed using RevMan 5.2 software (The Nordic Cochrane Centre, Copenhagen, Denmark).

Quality of evidence

We evaluated the quality of the evidence by using the Grades of Recommendation, Assessment, Development and Evaluation (GRADE) approach [15]. In addition, the GRADEprofiler 3.6 software (The Nordic Cochrane Centre) was used to create the evidence profile. The GRADE Working Group grades of evidence used were as follows:

- (1) *High quality*: Further research is very unlikely to change our confidence in the estimate of effect.
- (2) *Moderate quality*: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
- (3) *Low quality*: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

- (4) *Very low quality*: We are very uncertain about the estimate.

Results

In the initial search, we identified 1,263 records. After examination of the titles and abstracts, there were 30 potentially eligible studies assessed for inclusion. After application of the inclusion criteria, 13 RCTs [6,7,16-26] were included in the meta-analysis. The study flow diagram, including the reasons for exclusion of studies, is shown in Figure 1.

Study characteristics

The characteristics of the included trials are presented in Table 1. These trials were published between 1992 and 2014. The sample size of the trials ranged from 34 to 895, with a total of 2,525 patients comprising 1,299 in the GDT group and 1,226 in the control group. Eleven trials were conducted in the intensive care unit [16-26], and the remaining two were conducted in the emergency department [6,7]. Four trials were published in Chinese [23-26], and the other nine trials were in English [6,7,16-22]. Early GDT for resuscitation within the first 6 hours was reported in seven trials [6,7,16,21,22,24-26], late GDT for resuscitation outside the first 6 hours was assessed in one trial [23] and unclear timing of GDT was described in five trials [17-21]. Overall mortality was reported in all trials




Table 1 Characteristics of included randomized controlled trials^a

Study	Year	No. of patients with sepsis (GDT/control)	Study population	Clinical setting	Goals in GDT group	Goals in control group	Timing of GDT	Mortality endpoint
Tuchschmidt <i>et al.</i> [16]	1992	51 (26/25)	Adult patients with septic shock	ICU	CI ≥ 6 L/min/m ² SBP ≥ 90 mmHg	CI ≥ 3 L/min/m ² SBP ≥ 90 mmHg	Within the first 6 hr	14 days
Yu <i>et al.</i> [17]	1993	52 (30/22)	Adult patients with sepsis or septic shock (septic subpopulation)	ICU	DO ₂ I >600 ml/min/m ² SBP >100 mmHg	DO ₂ I 450 to 550 ml/min/m ² SBP >100 mmHg	Unclear	30 days
Hayes <i>et al.</i> [18]	1994	47 (24/23)	Adult patients with septic shock (septic subpopulation)	ICU	CI ≥ 4.5 L/min/m ² DO ₂ 600 ml/min/m ² VO ₂ > 170 ml/min/m ²	Usual care	Unclear	Hospital
Gattinoni <i>et al.</i> [19]	1995	181 (124/57)	Adult patients with septic shock or septic syndrome (septic subpopulation)	ICU	CI ≥ 4.5 L/min/m ² or SvO ₂ $\geq 70\%$ MAP ≥ 65 mmHg CVP 8 to 12 mmHg UO ≥ 0.5 ml/kg/h	CI 2.5 to 3.5 L/min/m ² MAP ≥ 65 mmHg CVP 8 to 12 mmHg UO ≥ 0.5 ml/kg/hr	Unclear	ICU
Yu <i>et al.</i> [20]	1998	87 (58/29)	Adult patients with sepsis, severe sepsis or septic shock (septic subpopulation)	ICU	DO ₂ I >600 ml/min/m ² SBP ≥ 100 mmHg SvO ₂ > 65% UO >50 ml/hr	DO ₂ I 450 to 550 ml/min/m ² SBP ≥ 100 mmHg SvO ₂ > 65% UO >50 ml/hr	Unclear	ICU
Alía <i>et al.</i> [21]	1999	63 (31/32)	Adult patients with severe sepsis or septic shock	ICU	DO ₂ I >600 ml/min/m ² MAP >60 mmHg	DO ₂ I >330 ml/min/m ² MAP >60 mmHg	Unclear	ICU
Rivers <i>et al.</i> [6]	2001	263 (130/133)	Adult patients with severe sepsis, septic shock or sepsis syndrome	ED	SvO ₂ $\geq 70\%$ CVP 8 to 12 mmHg MAP 65 to 90 mmHg UO ≥ 0.5 ml/kg/hr	CVP 8 to 12 mmHg MAP 65 to 90 mmHg UO ≥ 0.5 ml/kg/hr	Within the first 6 hr	Hospital
Lin <i>et al.</i> [22]	2006	224 (108/116)	Adult patients with septic shock	ICU	CVP 8 to 12 mmHg MAP ≥ 65 mmHg UO ≥ 0.5 ml/kg/hr	Usual care	Within the first 6 hr	Hospital
Wang <i>et al.</i> [23]	2006	34 (16/17)	Adult patients with septic shock	ICU	SvO ₂ $\geq 70\%$ CVP 8 to 12 mmHg MAP ≥ 65 mmHg UO ≥ 0.5 ml/kg/hr	MAP ≥ 65 mmHg UO ≥ 0.5 ml/kg/hr	Within the first 6 to 10 hr	14 days

Table 1 Characteristics of included randomized controlled trials^a (Continued)

Chen <i>et al.</i> [24]	2007	123 (58/65)	Adult patients with severe sepsis (septic subpopulation)	ICU	ScvO ₂ ≥ 70% CVP 8 to 12 mmHg MAP ≥65 mmHg UO ≥0.5 ml/kg/hr	CVP 8 to 12 mmHg MAP ≥65 mmHg UO ≥0.5 ml/kg/hr	Within the first 6 hr	ICU
He <i>et al.</i> [25]	2007	203 (98/105)	Adult patients with septic shock	ICU	ScvO ₂ or SvO ₂ ≥ 70% CVP 8 to 12 mmHg MAP ≥65 mmHg UO ≥0.5 ml/kg/hr	Usual care	Within the first 6 hr	Hospital
Yan <i>et al.</i> [26]	2010	303 (157/146)	Adult patients with severe sepsis or septic shock	ICU	ScvO ₂ ≥ 70% CVP 8 to 12 mmHg SBP >90 mmHg MAP ≥65 mmHg UO ≥0.5 ml/kg/hr	CVP 8 to 12 mmHg SBP >90 mmHg MAP ≥65 mmHg UO ≥0.5 ml/kg/hr	Within the first 6 hr	ICU
ProCESS [7]	2014	895 (439/456)	Adult patients with septic shock	ED	ScvO ₂ ≥ 70% CVP 8 to 12 mmHg MAP 65 to 90 mmHg UO ≥0.5 ml/kg/hr	Usual care	Within the first 6 hr	Hospital

^aCI, Cardiac index; CVP, Central venous pressure; DO₂, Oxygen delivery; DO₂I, Oxygen delivery index; ED, Emergency department; GDT, Goal-directed therapy; ICU, Intensive care unit; MAP, Mean arterial pressure; ProCESS, Protocolized Care for Early Septic Shock; SIRS, Systemic inflammatory response syndrome; SBP, Systolic blood pressure; ScvO₂, Central venous oxygen saturation; SvO₂, Mixed venous oxygen saturation; UO, Urine output; VO₂, Oxygen consumption.

[6,7,16-26], and dobutamine use was described in eight of them [6,7,16-19,21,22].

Risk of bias in included studies

The details of risk of bias are summarized in Figure 2. Five trials were judged to be at low risk of bias [6,7,19,21,22], and eight trials were judged to be at unclear risk of bias [16-18,20,23-26]. Adequate randomized sequences were generated in eight trials [6,7,17-19,21,22,26], and the investigators in five trials reported appropriate allocation concealment [6,7,19,21,22]. Among the 13 RCTs, none were double-blinded. However, blinding of patients and clinicians was extremely difficult in these trials to evaluate a complex intervention such as a GDT protocol, and we judged that the primary outcome (that is, overall mortality) was not likely to be influenced by lack of blinding.

Primary outcome: overall mortality

Mortality data were available in all 13 included trials [6,7,16-26]. The overall mortality data in the GDT and control groups were 474 (36.5%) of 1,299 and 520 (42.4%) of 1,226, respectively. Overall, GDT significantly reduced overall mortality in the random-effects model (RR, 0.83; 95% CI, 0.71 to 0.96; $P=0.01$; $I^2=56\%$) (Figure 3). Further exclusion of any single study did not alter the overall combined RR, which ranged from 0.80 (95% CI, 0.69 to 0.93) to 0.85 (95% CI, 0.73 to 0.98). The results of subgroup analyses are presented in Table 2.

Secondary outcomes

Predefined subgroup analysis according to the timing of GDT for resuscitation suggested that a mortality benefit was seen only in the subgroup in early GDT within the first 6 hours (seven trials; RR, 0.77; 95% CI, 0.67 to 0.89; $P=0.0004$; $I^2=40\%$) (Figure 4), but not in the subgroup with late or unclear timing of GDT (six trials; RR, 0.92; 95% CI, 0.69 to 1.24; $P=0.59$; $I^2=56\%$) (Figure 4).


In five trials, the investigators reported available data on dobutamine use [6,7,16,21,22]. In those trials, GDT was significantly associated with dobutamine use (RR, 2.71; 95% CI, 1.20 to 6.10; $P=0.02$; $I^2=86\%$).

Publication bias

We detected no evidence of publication bias by assessing funnel plot either visually (Figure 5) or statistically ($P=0.367$ by Egger test).

GRADE profile evidence

We found that GRADE Working Group grades of evidence were low for overall mortality, moderate for mortality in early GDT (within the first 6 hours for resuscitation)



and very low for dobutamine use. An additional .doc file shows these data in more detail (see Additional file 1).

Discussion

Our meta-analysis of 13 RCTs showed that GDT was associated with a 17% RR reduction on overall mortality in patients with sepsis. This mortality benefit was present in studies in which GDT was started early, but not when initiated late or when the timing of GDT was unclear. In addition, GDT was significantly associated with dobutamine use.

Despite these findings, the effect of GDT remains a matter of debate, as the most recent and largest trial




Figure 3 Forest plot of the effect goal-directed therapy on overall mortality. GDT, Goal-directed therapy.

included in this meta-analysis [7] did not show a difference in mortality, in contrast to many of the preceding studies. This could be due to the effect of the rapid acceptance of the principal interventions of the Rivers *et al.* study [6] and the subsequent Surviving Sepsis Campaign guidelines, which encompassed all elements of the Rivers study protocol. This is illustrated by the fact that, in the ProCESS (Protocolized Care for Early Septic Shock) trial, all groups received, on average, more than 2 L of fluid prior to randomization and more than 75% of patients received antibiotics before randomization. In addition, the mortality rate was much lower in the ProCESS trial than in preceding trials, possibly reflecting the effect of early diagnosis, fluid resuscitation and initiation of antibiotics on mortality. As the ProCESS trial, like the Rivers *et al.* study, enrolled patients with sepsis in the emergency department, this effect may have been prominent [27].

Although the current evidence supports the early use of GDT to improve outcomes in patients with sepsis, the


optimal goals remain uncertain. Currently, the Surviving Sepsis Campaign guidelines recommend the use of central venous pressure (CVP), mean arterial pressure, urine output and central venous oxygen saturation (ScvO₂) as resuscitation goals. However, many of these recommendations have been questioned in recent studies. A recent study [28] was designed to compare the use of lactate clearance to ScvO₂ as a goal of early (up to 6 hours) sepsis resuscitation. No significant difference in mortality was found (17% in the lactate clearance group versus 23% in ScvO₂ group). However, when both normalization of ScvO₂ and a rapid decrease in lactate levels were applied as therapeutic goals in the early resuscitation of a mixed group of critically ill patients, including a large subgroup of sepsis patients, mortality was significantly reduced [29]. In addition, in a recent retrospective study, researchers questioned the CVP endpoint in sepsis resuscitation [30].

Further research is needed before strong and definitive recommendations can be made regarding the effect of

Table 2 Subgroup analyses of overall mortality^a


Subgroups	No. of studies	No. of patients	RR (95% CI)	P-value	I ² (%)
All trials [6,7,16-26]	13	2,525	0.83 (0.71 to 0.96)	0.01	56
GDT timing					
Early [6,7,16,22,24-26]	7	2,062	0.77 (0.67 to 0.89)	0.0004	40
Late or unclear [17-21,23]	6	463	0.92 (0.69 to 1.24)	0.59	56
Risk of bias					
Low [6,7,19,21,22]	5	1,626	0.92 (0.75 to 1.13)	0.42	67
Unclear [16-18,20,23-26]	8	899	0.74 (0.62 to 0.89)	0.002	31
Sample size					
≥100 [6,7,19,22,24-26]	7	2,192	0.82 (0.70 to 0.95)	0.01	59
<100 [16-18,20,21,23]	6	333	0.81 (0.56 to 1.17)	0.27	61
Setting					
ED [6,7]	2	1,158	0.86 (0.52 to 1.44)	0.52	83
ICU [16-26]	11	1,367	0.81 (0.69 to 0.96)	0.01	53

^aCI, Confidence interval; ED, Emergency department; GDT, Goal-directed therapy; ICU, Intensive care medicine; RR, Relative risk.



GDT for resuscitation of patients with sepsis. There are currently at least two ongoing RCTs of GDT in patients with sepsis: the Australasian Resuscitation in Sepsis Evaluation (ARISE) trial in Australia (ClinicalTrials.gov ID: NCT00975793) and the Protocolised Management in Sepsis (ProMiSe) trial in the United Kingdom (Current Controlled Trials number: ISRCTN36307479) [31]. The results of these ongoing trials should provide further guidance as to the effect of GDT for resuscitation of patients with sepsis.

A major strength of the present meta-analysis is its compliance with the Cochrane handbook methodology recommendations. We conducted an exhaustive literature search that included non-English-language articles. Two authors independently screened all references, included eligible trials, extracted data information, assessed risk of bias and performed statistical analyses. Moreover, we followed the PRISMA statement to report this meta-analysis and evaluated the quality of the evidence by using the GRADE approach.



Because early fluid resuscitation is vital in patients with sepsis, we performed predefined subgroup analyses according to the timing of GDT. We also performed *post hoc* subgroup analyses according to risk of bias, disease severity, sample size and publication date. These subgroup analyses based on assessment of bias and clinically relevant groups may help health care professionals in clinical decision-making.

Our analysis also has several limitations that must be taken into consideration when interpreting the results. First, most of the included trials had a high risk of bias (Figure 2). The potential importance of this issue is highlighted by the fact that predefined subgroup analysis comparing mortality estimates between trials with low versus unclear risk of bias suggested the mortality benefit is not clearly apparent among the trials with low risk of bias, although this subgroup difference was not statistically significant ($P = 0.90$). Second, there were some differences in the target populations and protocols of GDT of each study. These factors may have a potential impact on our results and may preclude firm conclusions. Third, different endpoints were used for mortality evaluation. Because this study was a study-level meta-analysis, individual patient data were not included in the analysis; thus, we could not adjust for patient-level confounders.

Conclusions

The evidence suggests that GDT significantly reduces overall mortality in patients with sepsis, especially when initiated early (within the first 6 hours of admission). Until the results of ongoing randomized controlled trials are known, strong and definitive recommendations cannot be made regarding the effect of GDT for resuscitation of patients with sepsis.

Key messages

- The Surviving Sepsis Campaign guidelines recommend GDT for the early resuscitation of patients with sepsis, but controversies about its effect remain.
- The recent ProCESS trial has shown no mortality benefit from GDT in early septic shock.
- The current evidence, in the aggregate, suggests that GDT significantly reduces overall mortality in patients with sepsis, especially when initiated early (within the first 6 hours of admission).
- Further research is needed before strong and definitive recommendations can be made regarding the effect of GDT for resuscitation of patients with sepsis, and the optimal goals remain uncertain.

Additional file

Additional file 1: GRADE summary of findings.

Abbreviations

ARISE: Australasian Resuscitation in Sepsis Evaluation; CI: Confidence interval; CVP: Central venous pressure; GDT: Goal-directed therapy; GRADE: Grades of Recommendation, Assessment, Development and Evaluation; MAP: Mean arterial pressure; PRISMA: Preferred reporting items for systematic reviews and meta-analyses; ProMISe: Protocolised management in sepsis; RCT: Randomized controlled trial; RR: Relative risk; ScvO₂: Central venous oxygen saturation.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

WJG conceived of the study, participated in its design, collected the data, performed statistical analyses and drafted the manuscript. FW collected the data, performed statistical analyses and helped to draft the manuscript. JB participated in the study design, collected the data and helped to revise the manuscript critically for important intellectual content. LT collected the data, performed statistical analyses and helped to revise the manuscript critically for important intellectual content. JCL conceived of the study, participated in its design, collected the data and revised the manuscript critically for important intellectual content. All authors read and approved the final manuscript.

Authors' details

WJG is now studying as a master course student and majors in anesthesiology. His major research interests include systematic reviews and meta-analysis, evidence-based medicine, critical care medicine and clinical anesthesiology. As the first author, he has published more than 20 papers in peer-reviewed journals such as *Chest*, *European Respiratory Journal*, *Critical Care Medicine*, *Intensive Care Medicine* and *Critical Care*. He has authored chapters of a book on his research interests. He is one of section editors (Systematic Review and Meta-analysis) in the *Journal of Thoracic Disease*. He also acts as a forum moderator in <http://i.dxy.cn/profile/200750383>. FW is an anesthesiologist, and he has published more than 10 papers in peer-reviewed journals such as *Chest*, *Critical Care* and *American Journal of Emergency Medicine*. JB is a professor of medicine and vice-chair of the Department of Intensive Care Adults at Erasmus Medical Centre in Rotterdam, the Netherlands, and is a professor of medicine at the Pontificia Universidad Católica de Chile. He is also currently a visiting professor at Columbia University Medical Center-New York Presbyterian Hospital in New York City. He has published over 160 articles and edited 19 books, and he has made presentations at more than 450 conferences. He is the editor of *Intensive Care Medicine* and *Netherlands Journal of Critical Care*. He is a member of the editorial boards of *Reista Brasileira de Terapia Intensiva* (Brasil), the *Turkish Journal of Anaesthesiology and Reanimation*, *Journal of Critical Care* and *ICU Management*. He is former treasurer and president of the Dutch Society of Intensive Care and currently is a member of the organizing committee of the annual International Symposium on Intensive Care and Emergency Medicine in Brussels, Belgium. He has organized a 5-day workshop on intensive care annually for 11 years and annual meetings for ICU nurses for the past 9 years. LT is a nurse anesthetist. JCL is a professor of anaesthesiology and acts as the Director of the Department of Anaesthesiology in the First Affiliated Hospital of Guangxi Medical University.

Author details

¹Department of Anaesthesiology, the First Affiliated Hospital, Guangxi Medical University, 22 Shuangyong Road, Nanning 530021, China.

²Department of Anaesthesiology, General Hospital of Jinan Military Command, 25 Shifan Road, Jinan 250031, China. ³Department of Intensive Care Adults, Erasmus MC University Medical Centre, PO Box 2040, 3000 Rotterdam, CA, the Netherlands.

Received: 30 May 2014 Accepted: 3 October 2014

Published online: 20 October 2014

References

- Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, Sevransky JE, Sprung CL, Douglas IS, Jaeschke R, Osborn TM, Nunnally ME, Townsend SR, Reinhart K, Kleinpell RM, Angus DC, Deutschman CS, Machado FR, Rubenfeld GD, Webb SA, Beale RJ, Vincent JL, Moreno R, Surviving Sepsis Campaign Guidelines Committee including the Pediatric Subgroup: **Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2012.** *Crit Care Med* 2013, **41**:580–637.
- Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR: **Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care.** *Crit Care Med* 2001, **29**:1303–1310.
- Annane D, Bellissant E, Cavaillon JM: **Septic shock.** *Lancet* 2005, **365**:63–78.
- Centers for Disease Control and Prevention (CDC): **Current trends: increase in National Hospital Discharge Survey rates for septicemia—United States, 1979–1987.** *MMWR Morb Mortal Wkly Rep* 1990, **39**:31–34.
- Cecconi M, Corredor C, Arulkumaran N, Abuella G, Ball J, Grounds RM, Hamilton M, Rhodes A: **Clinical review: goal-directed therapy - What is the evidence in surgical patients? The effect on different risk groups.** *Crit Care* 2013, **17**:209.
- Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, Peterson E, Tomlanovich M, Early Goal-Directed Therapy Collaborative Group: **Early goal-directed therapy in the treatment of severe sepsis and septic shock.** *N Engl J Med* 2001, **345**:1368–1377.
- ProCESS Investigators: **A randomized trial of protocol-based care for early septic shock.** *N Engl J Med* 2014, **370**:1683–1693.
- Higgins JPT, Green S (Eds): *Cochrane Handbook for Systematic Reviews of Interventions (Version 5.1.0) [updated March 2011].* The Cochrane Collaboration; 2011. Available from www.cochrane-handbook.org.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *BMJ* 2009, **339**:b2535.
- Higgins JPT, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JA, Cochrane Bias Methods Group, Cochrane Statistical Methods Group: **The Cochrane Collaboration's tool for assessing risk of bias in randomised trials.** *BMJ* 2011, **343**:d5928.
- Higgins JPT, Thompson SG: **Quantifying heterogeneity in a meta-analysis.** *Stat Med* 2002, **21**:1539–1558.
- Higgins JPT, Thompson SG, Deeks JJ, Altman DG: **Measuring inconsistency in meta-analyses.** *BMJ* 2003, **327**:557–560.
- DerSimonian R, Laird N: **Meta-analysis in clinical trials.** *Control Clin Trials* 1986, **7**:177–188.
- Egger M, Davey Smith G, Schneider M, Minder C: **Bias in meta-analysis detected by a simple, graphical test.** *BMJ* 1997, **315**:629–634.
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ, GRADE Working Group: **GRADE: an emerging consensus on rating quality of evidence and strength of recommendations.** *BMJ* 2008, **336**:924–926.
- Tuchschmidt J, Fried J, Astiz M, Rackow E: **Elevation of cardiac output and oxygen delivery improves outcome in septic shock.** *Chest* 1992, **102**:216–220.
- Yu M, Levy MM, Smith P, Takiguchi SA, Miyasaki A, Myers SA: **Effect of maximizing oxygen delivery on morbidity and mortality rates in critically ill patients: a prospective, randomized, controlled study.** *Crit Care Med* 1993, **21**:830–838.
- Hayes MA, Timmins AC, Yau EH, Palazzo M, Hinds CJ, Watson D: **Elevation of systemic oxygen delivery in the treatment of critically ill patients.** *N Engl J Med* 1994, **330**:1717–1722.
- Gattinoni L, Brazzi L, Pelosi P, Latini R, Tognoni G, Pesenti A, Fumagalli R, SvO2 Collaborative Group: **A trial of goal-oriented hemodynamic therapy in critically ill patients.** *N Engl J Med* 1995, **333**:1025–1032.
- Yu M, Burchell S, Hasaniya NW, Takanishi DM, Myers SA, Takiguchi SA: **Relationship of mortality to increasing oxygen delivery in patients ≥50 years of age: a prospective, randomized trial.** *Crit Care Med* 1998, **26**:1011–1019.
- Alia I, Esteban A, Gordo F, Lorente JA, Diaz C, Rodriguez JA, Frutos F: **A randomized and controlled trial of the effect of treatment aimed at maximizing oxygen delivery in patients with severe sepsis or septic shock.** *Chest* 1999, **115**:453–461.
- Lin SM, Huang CD, Lin HC, Liu CY, Wang CH, Kuo HP: **A modified goal-directed protocol improves clinical outcomes in intensive care unit patients with septic shock: a randomized controlled trial.** *Shock* 2006, **26**:551–557.
- Wang XZ, Lü CJ, Gao FQ, Li XH, Yan WF, Ning FY: **[Efficacy of goal-directed therapy in the treatment of septic shock] [Article in Chinese].** *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 2006, **18**:661–664.
- Chen ZQ, Jin YH, Chen H, Fu WJ, Yang H, Wang RT: **[Early goal-directed therapy lowers the incidence, severity and mortality of multiple organ dysfunction syndrome] [Article in Chinese].** *Nan Fang Yi Ke Da Xue Xue Bao* 2007, **27**:1892–1895.
- He ZY, Gao Y, Wang XR, Hang YN: **[Clinical evaluation of execution of early goal directed therapy in septic shock] [Article in Chinese].** *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 2007, **19**:14–16.
- Early Goal-Directed Therapy Collaborative Group of Zhejiang Province: **[The effect of early goal-directed therapy on treatment of critical patients with severe sepsis/septic shock: a multi-center, prospective, randomized, controlled study] [Article in Chinese].** *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 2010, **22**:331–334.
- Wira CR, Dodge K, Sather J, Dziura J: **Meta-analysis of protocolized goal-directed hemodynamic optimization for the management of severe sepsis and septic shock in the emergency department.** *West J Emerg Med* 2014, **15**:51–59.
- Jones AE, Shapiro NI, Trzeciak S, Arnold RC, Claremont HA, Kline JA, Emergency Medicine Shock Research Network (EMShockNet) Investigators: **Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial.** *JAMA* 2010, **303**:739–746.
- Jansen TC, van Bommel J, Schoonderbeek FJ, Sleeswijk Visser SJ, van der Klooster JM, Lima AP, Willemsen SP, Bakker J, LACTATE study group: **Early lactate-guided therapy in intensive care unit patients: a multicenter, open-label, randomized controlled trial.** *Am J Respir Crit Care Med* 2010, **182**:752–761.
- Legrand M, Dupuis C, Simon C, Gayat E, Mateo J, Lukasiewicz AC, Payen D: **Association between systemic hemodynamics and septic acute kidney injury in critically ill patients: a retrospective observational study.** *Crit Care* 2013, **17**:R278.
- The ProCESS/ARISE/ProMISE Methodology Writing Committee: **Harmonizing international trials of early goal-directed resuscitation for severe sepsis and septic shock: methodology of ProCESS, ARISE, and ProMISE.** *Intensive Care Med* 2013, **39**:1760–1775.

doi:10.1186/s13054-014-0570-5

Cite this article as: Gu et al.: The effect of goal-directed therapy on mortality in patients with sepsis - earlier is better: a meta-analysis of randomized controlled trials. *Critical Care* 2014 **18**:570.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

