www.medscape.com

Non-invasive Detection of Hypovolemia or Fluid Responsiveness in Spontaneously Breathing Subjects

BMC Anesthesiology

Éva Zöllei, Viktória Bertalan, Andrea Németh, Péter Csábi, Ildikó László, József Kaszaki, László Rudas BMC Anesthesiol. 2013;13(40)

Abstract

Background In the assessment of hypovolemia the value of functional hemodynamic monitoring during spontaneous breathing is debated. The aim of our study was to investigate in spontaneously breathing subjects the changes in hemodynamic parameters during graded central hypovolemia and to test whether slow patterned breathing improved the discriminative value of stroke volume (SV), pulse pressure (PP), and their variations (SVV, PPV). In addition, we tested the alterations in labial microcirculation.

Methods 20 healthy volunteers participated in our study. Central hypovolemia was induced by lower body negative pressure (LBNP). Continuous signals of ECG, non-invasive blood pressure and central venous pressure were recorded. During baseline and each stage of LBNP the labial microcirculation was investigated by orthogonal polarization spectral imaging, 3 minute periods of patterned breathing at 6 and 15/min respiratory rate were performed, and central venous blood gas analysis was done. Data from baseline and those of different LBNP levels were compared by analysis of variance and those of different breathing rates by t-test. Finally, we performed ROC analysis to assess the discriminative values of SV, PP, SVV and PPV.

Results Moderate central hypovolemia induced by LBNP caused significant, clinically relevant falls in PP (p < 0.05) and SV and central venous oxygen saturation (ScvO ₂) (p < 0.001). The proportion of perfused vessels (p < 0.001) and microvascular flow index decreased (p < 0.05). PPV increased (p < 0.001), however the magnitude of fluctuations was greater during slow patterned breathing (p < 0.001). SVV increased only during slow patterned breathing (p < 0.001). ROC analysis confirmed the best predictive value for SV (at 56 ml cut-off AUC 0.97, sensitivity 94%, specificity 95%). Slow patterned breathing improved the discriminative value of SVV (p = 0.0023).

Conclusions Functional hemodynamic monitoring with slow patterned breathing to control spontaneous respiration may be worthy for further study in different populations for the assessment of hypovolemia and the prediction of volume responsiveness.

Background

The recognition of hypovolemia is of critical importance in the management of our patients. This is especially important in severe civil or military mass trauma situations where a parameter or a combination of parameters would be very useful to help to triage patients who are still "stable" but are approaching cardiovascular collapse due to haemorrhage. ^[1,2] Furthermore, in Intensive Care Units it is very often necessary to predict if patients with multiple organ failure are hypovolemic, need fluid loading, will tolerate fluid loading and will improve as a result.

Clinical signs and laboratory or blood gas data often help to establish the diagnosis of hypovolemia. Hemodynamic parameters such as cardiac filling pressures (central venous pressure, CVP, and pulmonary artery occlusion pressure, Paop) and static volumetric preload parameters (global end-diastolic volume index, GEDVI; intrathoracic blood volume index, ITBVI) have been used to approximate cardiac preload and to guide fluid therapy.

Recently the so-called dynamic hemodynamic parameters have been found to be more accurate in the

diagnosis of hypovolemia and fluid responsiveness. ^[3–5] For calculating these parameters we use preload alterations (most often induced by respiration) and the resulting changes in either right atrial pressure, blood pressure, pulse pressure, stroke volume or cardiac output. From these fluctuations we can calculate the systolic pressure variation (SPV), pulse pressure variation (PPV) and stroke volume variation (SVV). ^[5–15] However, it has been demonstrated in previous studies that these indices are of clinical value only if the patient is mechanically ventilated without spontaneous breathing activity. ^[16–19] We also know that the level of PEEP, respiratory rate and tidal volume also influence the value of SPV, PPV and SVV. ^[3,9,20–26] In addition, significant arrhythmias, pulmonary hypertension, increased intraabdominal pressure, right and left heart failure and the use of catecholamines also influence their predictive value. ^[19,27–29]

From investigations on cardiovascular autonomic regulation it is clear that respiration has a fundamental influence on heart rate and blood pressure fluctuations. To control the breathing of awake and cooperative subjects the method of patterned breathing is used. This means that the subjects synchronize inspiration and expiration and sometimes tidal volume to an audio or visual signal. The respiratory frequency can be centred either to the high frequency (respiratory, 0.25 Hz) or low frequency (around 0.1 Hz) bands of cardiovascular fluctuations, which means 15/min and 6/min rate. Doing so, the oscillations in these bands can be enhanced, especially at 6/min breathing rate when the respiratory oscillations and Mayer waves merge.

To induce central hypovolemia in healthy volunteers lower body negative pressure (LBNP) is an accepted, safe and widely used method. This results in the redistribution of blood volume, which generates central hypovolemia without actual volume loss. According to previous data, the volume shift generated by -10-20 mmHg LBNP is similar to mild (400–550 ml), by -20-40 mmHg to moderate (550–1000 ml), and by greater than -40 mmHg to severe (more than 1000 ml) acute blood loss. ^[30]

Experimental and clinical research suggests that in circulatory failure or shock it is the alteration of the microcirculation that finally leads to tissue oxygenation problems and multiple organ failure. ^[31–33] Furthermore, ultimately the improvement of microcirculatory function determines survival, therefore it is important to know its behaviour during different conditions. ^[32]

The aim of our study was to investigate the effects of mild and moderate LBNP-induced central hypovolemia on the behaviour of hemodynamic parameters, especially on PPV and SVV in spontaneously breathing subjects, and to test the hypothesis that slow patterned breathing improves their discriminative value. Furthermore, we studied the microvascular alteration caused by different levels of central hypovolemia.

Methods

Our study population consisted of 20 healthy young male volunteers (age 24 ± 6 years), who provided informed consent to participation. Firstly, the subjects were familiarized with the laboratory and equipment, and were trained to perform patterned breathing. In order to control their respiration, patterned breathing was used, so they synchronized inspiration and expiration to an audio signal. Afterwards we inserted a peripherally inserted central line. We ascertained the correct position of the catheter tip by the appearance of the classic central venous pressure (CVP) wave form on the monitor. After this the subjects assumed supine position in the LBNP chamber.

We monitored the ECG, CVP and arterial oxygen saturation continuously using a bedside monitor (Marquette Eagle, Marquette Electronics, Milwaukee, Wi, USA) and arterial blood pressure by the Finapres non-invasive monitor (Ohmeda, Englewood, CO, USA). During the patterned breathing periods we monitored the end tidal CO ₂ level through a mouth-piece connected to a capnograph (Novametrix Medical Systems, Inc., Wallingford, CT, USA) to confirm the subjects' adherence to the breathing commands. All these signals were digitized by 500 Hz and recorded on-line by the Dataq/Windaq data acquisition system (Dataq Instruments, Inc., Akron, OH, USA).

In order to study microcirculation, we investigated the labial microvessels. For this the orthogonal polarization

spectral (OPS) imaging technique was used (Cytoscan A/R, Cytometrix, Philadelphia, PA, USA). A 10× objective was placed onto the inner surface of the labia, and microscopic images were recorded on an S-VHS video recorder (Panasonic AG-TL 700, Matsushita Electric Ind. Co. Ltd, Osaka, Japan).

Central hypovolemia was induced by -20 and -40 mmHg LBNP. During baseline and each stage of LBNP first the labial microcirculation was investigated and afterwards 3 minute periods of patterned breathing at 6 and 15/min respiratory rate were done. In addition, central venous blood sampling was performed to measure pH, central venous oxygen tension (PcvO₂), central venous CO₂ tension (PcvCO₂), central venous oxygen saturation (ScvO₂) and haemoglobin (Hb) (Figure 1). The study protocol was approved by the Institutional Human Medical Biological Research Ethics Committee of the University of Szeged, and was carried out in accordance with the Declaration of Helsinki.

Medscape			
	baseline	- 20 mmHg LBNP	- 40 mmHg LBNP
breathing	6/min 15/min	6/min 15/min	6/min 15/min
ECG -			
BP —			•
CVP -			
0	PS	OPS (DPS
	VE	3G VBG	VBG
		Source: BMC Anesthes	iol © 2013 BioMed Central, Ltd

Figure 1.

Study protocol. (BP: blood pressure, CVP: central venous pressure, SaO ₂: arterial oxygen saturation, OPS: orthogonal polarisation spectral imaging, VBG: central venous blood gas analysis).

For the analysis of hemodynamic data the WinCPRS (Absolute aliens Ay, 2000, Turku, Finland) program was used. For SV calculation this program uses the pulse-contour method (modified Modelflow). First, all the signals were automatically peak-detected and manually checked. Then we calculated RR interval (RRI), systolic arterial pressure (SAP), mean arterial pressure (MAP), pulse pressure (PP) and SV for the entire 3 minute periods in each study condition. Afterwards, we calculated the minimum and maximum RRI, SAP, MAP, CVP, PP and SV for each respiratory cycle during 1 minute of the patterned breathing periods and calculated their differences (delta values) as the respiratory fluctuations (this involved 6 respiratory cycles during 6/min and 15 cycles during 15/min breathing). In addition, we calculated PPV and SVV according to accepted formulas. ^[26]

For the analysis of data from the microcirculation we used a computer-assisted image analysis system (IVM Pictron, Budapest, Hungary). Quantitative assessment of the microcirculatory parameters was performed off-line by frame-to-frame analysis of the videotaped images. We determined red blood cell velocity (RBCV), the mean perfused capillary lengths (MPCL), the proportion of perfused vessels (PRPV) and the microvascular flow index (MFI) in accordance with accepted methods. ^[34] All microcirculatory evaluations were performed by the same investigator, who was not blinded to the study conditions, but was not aware of the results of the hemodynamic assessment (IL).

Data from baseline and those of different LBNP levels were compared by analysis of variance and those of the two different breathing rates by t-test. In the case of non-normal distribution of data, analysis of variance on ranks and the Mann–Whitney rank sum test were used. (Sigmastat, Aspire Software International, Ashburn, VA, USA). In order to assess the discriminative value of data, ROC analysis was done. SV and PP during 6/min, and SVV and PPV during both respiratory rates were analysed by using data from baseline and -40 mmHg LBNP (Medcalc, Mariakerke, Belgium). Data are given as mean \pm SD. P value below 0.05 was considered to be statistically significant.

Results

The typical recording of one subject is shown in Figure 2. at different LBNP levels and at 6/min and 15/min patterned breathing, which demonstrates the enhanced fluctuations at moderate hypovolemia and slow patterned breathing in heart rate, arterial blood pressure and central venous pressure. The whole protocol was completed in 11 subjects. 4 of them had presyncopal symptoms, and -40 mmHg LBNP could not be performed (therefore during moderate central hypovolemia we analysed data from 16 volunteers), and in another 5 PIC line insertion was unsuccessful (as a result we have CVP data from 15 volunteers).



Figure 2.

The typical recording of one subject at the different LBNP levels and at 6/min and 15/min patterned breathing.

Hemodynamic, Central Venous Blood Gas and Microcirculatory Parameters at Different Levels of LBNP

From the hemodynamic parameters MAP did not change; but RRI, SAP, CVP, PP and SV significantly

decreased at -40 mmHg, and CVP and SV even at -20 mmHg LBNP. In the case of RRI, SAP, PP and SV there was also a significant difference between -20 and -40 mmHg LBNP ().

	Baseline	−20 mmHg LBNP	−40 mmHg LBNP
RRI (msec) 859 ± 159		844 ± 144	739 ± 115 ^{##§§}
SAP (mmHg)	130 ± 17	126 ± 15	123 ± 11 ##§
MAP (mmHg)	83 ± 13	84 ± 10	87 ± 8
CVP (mmHg)	<mark>7.2</mark> ±2.0	3.7 ± 2.0 **	<mark>2.6 ± 2.0</mark> ##
PP (mmHg)	65 ± 10	58 ± 10	<mark>50 ± 7 ^{#§}</mark>
SV (ml)	<mark>79 ±</mark> 12	<mark>63 ± 12</mark> **	<mark>50 ± 8</mark> ##§§
рН	7.38 ± 0.02	7.39 ± 0.03	7.39 ± 0.04 [#]
pCO ₂ (mmHg)	45 ± 3	43 ± 4	43 ± 5 [#]
pO ₂ (mmHg)	42 ± 5	37 ± 4 [*]	36 ± 4 ##
ScvO ₂ (%)	<mark>76 ±</mark> 6	70 ± 7 [*]	<mark>67 ± 9</mark> ##
Hb (g/dl)	13.7 ± 0.8	14 ± 0.8 **	14.5 ± 0.9 ##§§
RBCV (microm/s)	1035 ± 67	792 ± 153 ^{**}	696 ± 156 ^{##§}
MPCL (microm)	2696 ± 593	2194 ± 567 **	1973 ± 512 ##
PRPV (%)	99.7 ± 1.5	90.4 ± 13 [*]	87.8 ± 13.9 ##
MFI	3.0 ± 0	2.8 ± 0.4 *	2.7 ± 0.4 [#]

Table 1. Hemodynamic parameters, central venous blood gases and microcirculatory variables

Hemodynamic data are calculated from the whole 3 minute recordings at 6/min patterned breathing at baseline and during mild (-20 mmHg LBNP) and moderate (-40 mmHg LBNP) central hypovolemia and are given as mean \pm SD.

*baseline vs. -20 mmHg, p < 0.05.

**baseline vs. -20 mmHg, p < 0.001.

#baseline vs. −40 mmHg, p < 0.05.

##baseline vs. -40 mmHg, p < 0.001.

§-20 vs -40 mmHg, p < 0.05.

§§-20 vs -40 mmHg, p < 0.001.

Central venous blood gas analysis showed that pH and Hb significantly increased, pCO $_2$, pO $_2$ and ScvO $_2$ significantly deceased at -40 mmHg LBNP, and this change was even significant between baseline and -20 mmHg in Hb, pO $_2$ and ScvO $_2$, and between -20 mmHg and -40 mmHg in Hb ().

Table 1. Hemodynamic parameters, central venous blood gases and microcirculatory variables

	Baseline	−20 mmHg LBNP	−40 mmHg LBNP
RRI (msec)	859 ± 159	844 ± 144	739 ± 115 ^{##§§}
SAP (mmHg)	130 ± 17	126 ± 15	123 ± 11 ^{##§}
MAP (mmHg)	83 ± 13	84 ± 10	87 ± 8
CVP (mmHg)	7.2 ± 2.0	3.7 ± 2.0 **	2.6 ± 2.0 ##

PP (mmHg)	65 ± 10	58 ± 10	50 ± 7 ^{#§}
SV (ml)	79 ± 12	63 ± 12 **	50 ± 8 ^{##§§}
рН	7.38 ± 0.02	7.39 ± 0.03	7.39 ± 0.04 [#]
pCO ₂ (mmHg)	45 ± 3	43 ± 4	43 ± 5 [#]
pO ₂ (mmHg)	42 ± 5	37 ± 4 [*]	36 ± 4 ^{##}
ScvO ₂ (%)	76 ± 6	70 ± 7 [*]	67 ± 9 ##
Hb (g/dl)	13.7 ± 0.8	14 ± 0.8 **	14.5 ± 0.9 ##§§
RBCV (microm/s)	1035 ± 67	792 ± 153 ^{**}	696 ± 156 ^{##§}
MPCL (microm)	2696 ± 593	2194 ± 567 **	1973 ± 512 ##
PRPV (%)	99.7 ± 1.5	90.4 ± 13 [*]	87.8 ± 13.9 ##
MFI	3.0 ± 0	2.8 ± 0.4 *	2.7 ± 0.4 [#]

Hemodynamic data are calculated from the whole 3 minute recordings at 6/min patterned breathing at baseline and during mild (-20 mmHg LBNP) and moderate (-40 mmHg LBNP) central hypovolemia and are given as mean \pm SD.

*baseline vs. -20 mmHg, p < 0.05.

**baseline vs. -20 mmHg, p < 0.001.

#baseline vs. -40 mmHg, p &It; 0.05.

##baseline vs. -40 mmHg, p < 0.001.

§-20 vs -40 mmHg, p < 0.05.

§§-20 vs -40 mmHg, p < 0.001.

All the microcirculatory variables (RBCV, MPCL, PRPV and MFI) decreased significantly between baseline and both LBNP levels, and RBCV between -20 and -40 mmHg LBNP, too ().

	Baseline	−20 mmHg LBNP	−40 mmHg LBNP
RRI (msec)	859 ± 159	844 ± 144	739 ± 115 ^{##§§}
SAP (mmHg)	130 ± 17	126 ± 15	123 ± 11 ^{##§}
MAP (mmHg)	83 ± 13	84 ± 10	87 ± 8
CVP (mmHg)	7.2 ± 2.0	3.7 ± 2.0 **	2.6 ± 2.0 ##
PP (mmHg)	65 ± 10	58 ± 10	50 ± 7 ^{#§}
SV (ml)	79 ± 12	63 ± 12 **	50 ± 8 ^{##§§}
рН	7.38 ± 0.02	7.39 ± 0.03	7.39 ± 0.04 [#]
pCO ₂ (mmHg)	45 ± 3	43 ± 4	43 ± 5 [#]
pO ₂ (mmHg)	42 ± 5	37 ± 4 [*]	36 ± 4 ##
ScvO ₂ (%)	76 ± 6	70 ± 7 [*]	67 ± 9 ##
Hb (g/dl)	13.7 ± 0.8	14 ± 0.8 **	14.5 ± 0.9 ##§§
RBCV (microm/s)	1035 ± 67	792 ± 153 ^{**}	696 ± 156 ^{##§}

Table 1.	Hemodynamic parameters,	central venous blood gase	es and microcirculatory variables
----------	-------------------------	---------------------------	-----------------------------------

MPCL (microm)	2696 ± 593	2194 ± 567 **	1973 ± 512 ^{##}
PRPV (%)	99.7 ± 1.5	90.4 ± 13 [*]	87.8 ± 13.9 ##
MFI	3.0 ± 0	2.8 ± 0.4 *	2.7 ± 0.4 [#]

Hemodynamic data are calculated from the whole 3 minute recordings at 6/min patterned breathing at baseline and during mild (-20 mmHg LBNP) and moderate (-40 mmHg LBNP) central hypovolemia and are given as mean \pm SD.

*baseline vs. -20 mmHg, p < 0.05.

**baseline vs. -20 mmHg, p < 0.001.

#baseline vs. -40 mmHg, p < 0.05.

##baseline vs. -40 mmHg, p < 0.001.

§-20 vs -40 mmHg, p < 0.05.

§§-20 vs -40 mmHg, p < 0.001.

Respiratory Fluctuations at Different Levels of LBNP and During Different Respiratory Rates

RRI, SAP, MAP, CVP, SV and PP data at different levels of LBNP and at different respiratory rates are shown in

	6/min patterned breathing			15/min patterned breathing		
	Baseline	−20 mmHg	−40 mmHg	Baseline	−20 mmHg	−40 mmHg
RRI (msec) min	767 ± 130	742 ± 130	651 ± 90 ^{#§}	823 ± 151	793 ± 164	674 ± 127 ^{##§§}
max	952 ± 157	950 ± 177	823 ± 151 ^{##§§}	889 ± 171	848 ± 177 [*]	726 ± 145 ^{##§§}
delta	185 ± 86	208 ± 103	173 ± 84	65 ± 32	54 ± 24	52 ± 25
SAP (mmHg) min	123 ± 17	117 ± 16	113 ± 11 ^{#§}	125 ± 14	122 ± 14	118 ± 11 ^{#§}
max	135 ± 18	133 ± 15	131 ± 12 [§]	131 ± 15	128 ± 13	125 ± 11 ^{#§}
delta	12 ± 4	16 ± 6 **	18 ± 8 ^{##}	6 ± 2	6 ± 2	7 ± 3 [#]
MAP (mmHg) min	78 ± 12	77 ± 11	78 ± 9	81 ± 11	83 ± 10	86 ± 8
max	87 ± 13	90 ± 10	94 ± 9	85 ± 12	87 ± 10	91 ± 8
delta	9±4	12 ± 5 **	15 ± 6 ^{##§}	3 ± 1	4 ± 1	5 ± 2 [#]
CVP (mmHg) min	2.7 ± 2.4	-0.2 ± 2.3 **	-0.7 ± 2.5 ^{##}	3 ± 2.7	1.5 ± 2.4 [*]	-0.35 ± 2.8 ^{##§}
max	12 ± 3.1	6.7 ± 2.8 **	5.5 ± 2.6 ^{##}	10 ± 2.8	5.8 ± 2.3 **	4.6 ± 2.2 ##
<mark>delta</mark>	9.3 ± 1.8	6.9 ± 1.7 **	6.2 ± 1.8 ^{##}	7 ± 2.2	4.4 ± 1.5 **	4.9 ± 1.7 [#]
PP (mmHg) min	60 ± 10	52 ± 10 [*]	44 ± 5 ^{#§}	60 ± 7	53 ± 8 **	43 ± 7 ^{##§§}
max	68 ± 11	61 ± 11 **	53 ± 7 ^{##§§}	64 ± 7	57 ± 8 **	49 ± 7 ^{##§§}
delta	8 ± 4	9 ± 4	10 ± 4	4 ± 1	4 ± 2	5 ± 2
SV (ml) min	72 ± 11	55 ± 10 **	42 ± 6 ^{##§§}	70 ± 10	55 ± 9 **	43 ± 6 ^{##§§}
max	85 ± 13	67 ± 14 **	55 ± 10 ^{##§§}	79 ± 12	64 ± 11 **	50 ± 7 ^{##§§}
delta	12 ± 5	13 ± 8	14 ± 8	9 ± 4	8 ± 4	7 ± 3

Table 2. The respiratory fluctuations of hemodynamic parameters

The data for RRI, SAP, MAP, CVP, PP and SV were calculated from 1 minute recordings (so from 6 and 15 respiratory cycles), during mild and moderate central hypovolemia and at 6/min and 15/min patterned breathing and are given as mean \pm SD. *baseline vs. -20 mmHg, p &It; 0.05.

**baseline vs. -20 mmHg, p < 0.001. #baseline vs. -40 mmHg, p < 0.05. ##baseline vs. -40 mmHg, p < 0.001. §-20 vs -40 mmHg, p < 0.05.

§§-20 vs -40 mmHg, p < 0.001.

Delta RRIs were not influenced by the LBNP level, but their values were significantly greater during 6/min patterned breathing at all LBNP levels (p<0.001).

SAP and MAP respiratory fluctuations increased at both 6/min and 15/min patterned breathing in moderate central hypovolemia, but in mild central hypovolemia these changes were only significant at 6/min breathing rate. The SAP respiratory fluctuations were greater at slow patterned breathing at each LBNP level (p&It;0.001). MAP delta increased between -20 and -40 mmHg LBNP only at 6/min patterned breathing.

The CVP delta values changed significantly between baseline and mild and moderate central hypovolemia, but the fluctuations became smaller at both 6 and 15/min respiratory rate.

The assessment of SV, PP, SVV and PPV gave the most interesting results. The minimum, and maximum values of SV and PP decreased in response to all levels of hypovolemia, though the delta values did not change either at 6/min nor at 15/min breathing rate. The fluctuations of both SV and PP were more pronounced at 6/min patterned breathing, and the differences were significant in all comparisons (p<0.001 for PP; p<0.05 for SV), except at baseline in SV. However, PPV increased from baseline to -40 mmHg LBNP, especially at slow patterned breathing. PPV increased also between -20 and -40 mmHg LBNP at 15/min patterned breathing. SVV only increased at slow patterned breathing in moderate central hypovolemia, though it was greater in all 3 conditions with slow respiratory rate ().

	Baseline	−20 mmHg LBNP	−40 mmHg LBNP
PPV 6 (%)	13 ± 5	16 ± 7	<mark>20 ± 8</mark> ##
PPV 15 (%)	7 ± 3 **	8 ± 4 **	11 ± 4 ^{##§**}
<mark>SVV 6</mark> (%)	15 ± 5	20 ± 10	<mark>26 ± 12</mark> ^{##}
SVV 15 (%)	12 ± 4 [*]	13 ± 5 [*]	15 ± 6 [*]

Table 3. PPV and SVV during different levels of central hypovolemia and different respiratory rates

Data are calculated from 1 minute recordings (so from 6 and 15 respiratory cycles) at 6/min (PPV 6, SVV 6) and 15/min (PPV 15, SVV 15) patterned breathing and are given as mean ± SD.

##baseline vs. -40 mmHg, p < 0.001.

§-20 vs -40 mmHg, p < 0.05.

*6 vs 15/min patterned breathing, p < 0.05.

**6 vs 15/min patterned breathing, p &It; 0.001.

Finally we performed ROC analysis (). The AUC was the largest and the sensitivity and specificity the best for SV with a 56 ml cut-off, though PP also had acceptable sensitivity and specificity. Slow patterned breathing improved the discriminative value of SVV.

Table 4. ROC analysis of SV, PP, SVV and PPV

AUC (CI)	Cut-off	Sensitivity (%)	Specificity (%)
----------	---------	-----------------	-----------------

SV	<mark>0.97 (0.85–1.0)</mark>	56 ml	94	95
PP	<mark>0.88</mark> (0,73–0.97)	53 mmHg	88	85
SVV 6	<mark>0.83</mark> (0.67–0.94)	<mark>16%</mark>	81	75
PPV 6	0.78 (0.61–0.9)	13%	81	65
SVV 15 [*]	0.69 (0.51–0.83)	13%	69	70
PPV 15	0.83 (0.66–0.93)	8%	81	75

Data calculated from 6/min (SV, PP, SVV 6, PPV 6) and 15/min (PPV 15, SVV 15) patterned breathing during -40 mmHg LBNP.

*6 vs 15/min patterned breathing, p &It; 0.0023.

Discussion

The main findings of our study are as follows: mild and moderate central hypovolemia induced by LBNP caused significant and clinically relevant falls in PP and SV. This was accompanied by significant and relevant decreases in PcvO₂ and ScvO₂. All the investigated microcirculatory variables such as RBCV, MPCL, PRPV and MFI decreased significantly at both levels of central hypovolemia, which indicates progressive impairment of both microvessel density and perfusion. PPV increased at both 6/min and 15/min patterned breathing in moderate central hypovolemia however the magnitude of fluctuations was greater at the slow respiratory rate. SVV increased only during slow breathing, and its accentuation with low respiratory rate was less pronounced then that of PPV. To our knowledge this is the first study that investigates SVV and PPV during LBNP and at different rates of patterned breathing. ROC analysis confirmed the best predictive value for SV absolute values in this group of healthy volunteers. Slow patterned breathing improved the discriminative value of SVV, but not that of PPV.

Recently the effects of LBNP induced hypovolemia on systemic and tissue oxygenation and microcirculation have been investigated in two studies, in which sublingual side stream dark field imaging has been used in healthy volunteers. ^[35,36] In both they increased the LBNP level until hemodynamic decompensation occurred. Ward et al. found significant decrease in systemic oxygen delivery accompanied by progressive decrease in sublingual functional capillary density (FCD). The oxygen extraction ratio increased both at systemic and local (muscle) level. The fall of FCD occurred <u>earlier</u> than any other change in tissue oxygenation parameters. Bartels et al. found significant decrease not only in perfused vessel density but also in the microvascular flow index, as well as a significant increase in microvascular flow heterogeneity. Our results regarding the microcirculatory alterations were similar, confirming that even -20 mmHg LBNP causes hemodynamic changes great enough to influence microcirculation. The fall in ScvO2 with increasing levels of LBNP is consistent with decreased cardiac output and oxygen delivery, if we assume that oxygen consumption did not change during supine rest.

Since there is strong clinical need to recognize hypovolemia in patients who are not mechanically ventilated, we performed our investigations in spontaneously breathing subjects. The mechanism of respiratory fluctuations in stroke volume and blood pressure is different during spontaneous breathing and positive pressure ventilation. ^[10,11,15,37] During mechanical ventilation positive pressure inspiration impedes venous return and can increase right ventricular (RV) afterload. These result in a decrease of left ventricular (LV) preload a few heart beats later. However, inspiration also squeezes blood from the pulmonary capillaries towards the left ventricle, and at the same time decreases LV afterload by decreasing aortic transmural pressure. These effects result in an early inspiratory increase in LV SV and arterial pressure, which is called delta up. This is followed by an inspiratory decrease a few heart beats later, which is called delta down. During expiration the RV preload increases, the afterload decreases, the afterload increases, which results in lower LV SV and blood pressure. This is followed by an increase in SV and SAP a few beats later when the increased RV SV reaches the left side of the heart. In this way the delta up results mainly from the increase in LV preload and reduction of its afterload,

and delta down mainly from decreased venous return to the right heart.

During spontaneous breathing the mechanism of stroke output changes is different. During spontaneous inspiration the venous return increases without a considerable increase in RV afterload. On the left side there is no afterload reduction, so the inspiratory increase in LV SV and arterial pressure (delta up) is the result of the increased RV output. During exhalation, the venous return and RV SV decrease, which reaches the LV a few beats later. At the same time the aortic transmural pressure decreases compared to the inspiratory value, which augments LV ejection. This dampens the effect of decreased preload, so delta down is not so prominent, as during positive pressure ventilation. In summary, the mechanism of delta up and delta down is different: during mechanical ventilation it is the delta down, during spontaneous breathing it is the delta up that is the volume sensitive, preload determined variable.

There is another problem with dynamic hemodynamic parameters besides the above discussed issues. The respiratory perturbation should be great enough to cause significant preload alteration in order for clinicians to be able to assess its effects on blood pressure and left ventricular stroke volume. These effects are smaller during quiet spontaneous breathing and are clearly tidal volume dependent during mechanical ventilation. [6,16–18,20–22,24,25]

In addition, the lag between the changes of RV and LV output is determined by the pulmonary transit time. The timing of the influence of RV output changes on the LV is clearly dependent on the heart rate (HR) and respiratory rate (RR) as well as their timing to each other. This relationship is highlighted in the work of De Backer et al..^[23] They concluded that the respiratory variations in left but not in RV stroke volume and its derivates disappeared with high RR, and when the HR/RR ratio fell below 3.6. This relationship however can change from breath to breath during spontaneous respiration, or when the patient triggers the ventilator.

Based on these findings it may be assumed that there is a need to standardize conditions, especially respiratory stimuli when assessing preload responsiveness by dynamic parameters. Perel et al. tried to achieve this with a ventilatory manoeuvre called respiratory systemic variation test (RSVT). ^[38] In our study we tried to control the spontaneous breathing of our subject by patterned breathing. The 6/min respiratory rate looked more promising because it merges the effects of sympathetic activity (Mayer waves) and respiration thus increasing blood pressure fluctuations. Besides this, during slow patterned breathing the tidal volume is greater than during quiet breathing, because the subjects automatically adjust it to keep minute ventilation constant. We hypothesized that this stimulus would be regular and great enough to cause measurable changes in dynamic parameters and improve their discriminative value compared to 15/min breathing rate, which served as a control, being similar to normal quite breathing. According to our results, although the fluctuations in blood pressure and SVV and PPV increased, ROC analysis confirmed improvement in discriminative value only of SVV. Still absolute values of SV and PP performed much better in this group of healthy volunteers.

The different behaviour of SVV and PPV may be explained by the fact that the relationship between SV and PP is not linear, arterial distensibility continues to decrease with progressive reduction of central blood volume. [1,29]

Finally, we were <u>interested</u> in how the <u>respiratory fluctuations of CVP</u> were influenced, because this is easy to measure and is <u>readily available</u>. In his early study Magder tested the hypothesis that the respiratory pattern of right atrial pressure (RAP) could be used to predict the response to a fluid challenge. He concluded that it was <u>very unlikely that volume loading would increase cardiac output</u> when the <u>inspiratory fall in RAP was less 1</u> <u>mmHg</u>. ^[13] Although this <u>fall was greater than 1 mmHg</u> in our subjects even at <u>baseline</u>, the <u>amplitude</u> of <u>fluctuations decreased</u> during <u>mild and moderate hypovolemia</u>. We explain this by the <u>ceiling effect</u> due to the <u>collapse of large veins</u>.

This study has some limitations, the first one being that we investigated healthy young individuals with intact autonomic regulation therefore the results cannot be extrapolated to different patient populations. Another limitation was that our subjects were fully awake and cooperative and after a short learning period could follow the breathing commands properly. The implementation of slow patterned breathing could be more difficult and even impossible in many clinical situations where patients have dyspnea, tachypnea or altered mental state.

However we believe that not only during spontaneous but also during controlled mechanical ventilation there is a need to standardise the respiratory pattern when we want to assess functional hemodynamic parameters. For stroke volume analysis we used pulse-contour analysis with modified Modelflow method and verified manually the beat-to-beat automatic wave form analysis. Furthermore, when we investigated the effect of respiratory rate, we averaged data for 1 minute, from 6 and 15 respiratory cycles. Commercially available, built-in systems, with different softwares and time-averaged data could give different results.^[27]

Conclusions

In conclusion, slow patterned breathing may allow the assessment of hypovolemia in spontaneously breathing subjects and it is worthy for further study in different conditions and populations. In Emergency Departments and wards patients are usually not equipped with arterial lines and invasive hemodynamic monitoring is not available. In this patient group, for those, who are able to sustain a specific breathing pattern for a few minutes, the use of non-invasive blood pressure monitoring coupled with functional hemodynamic monitoring, is promising.

Sidebar

Key Messages

- Mild and moderate central hypovolemia induced by LBNP caused significant and clinically relevant falls in PP and SV. This was accompanied by significant and relevant decreases in PcvO₂ and ScvO₂ and progressive impairment of both microvessel density and perfusion.
- PPV increased during both 6/min and 15/min patterned breathing in moderate central hypovolemia however the magnitude of fluctuations was greater during the slow respiratory rate. SVV increased only during slow breathing, and its accentuation with low respiratory rate was less pronounced then that of PPV.
- ROC analysis confirmed the best predictive value for SV absolute values in this group of healthy volunteers.
- Slow patterned breathing improved the discriminative value of SVV, but not that of PPV.

References

- 1. Convertino VA, Cooke WH, Holcomb JB: Arterial pulse pressure and its association with reduced stroke volume during progressive central hypovolemia. *J Trauma* 2006, 61:629–634.
- 2. Cooke WH, Salinas J, Convertino VA, Ludwig DA, Hinds D, Duke JH, Moore FA, Holcomb JB: Heart rate variability and its association with mortality in prehospital trauma patients. *J Trauma* 2006, 60:363–370.
- Huang CC, Fu JY, Hu HC, Kao KC, Chen NH, Hsieh MJ, Tsai YH: Prediction of fluid responsiveness in acute respiratory distress syndrome patients ventilated with low tidal volume and high positive end-exspiratory pressure. *Crit Care Med* 2008, 36:2810–2816.
- Preisman S, Kogan S, Berkenstadt H, Perel A: Predicting fluid responsiveness in patients undergoing cardiac surgery: functional haemodynamic parameters including the respiratory systolic variation test and static preload indicators. *Br J Anaesth* 2005, 95:746–755.
- Michard F, Boussat S, Chemla D, Anguel N, Mercat A, Lecarpentier Y, Richard C, Pinsky MR, Toboul JL: Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure. *Am J Respir Crit Care Med* 2000, 162:134–138.
- 6. Teboul JL, Monnet X: Prediction of volume responsiveness in critically ill patients with spontaneous

breathing activity. Curr Opin Crit Care 2008, 14:334-339.

- 7. Pinsky MR: Functional hemodynamic monitoring. Intensive Care Med 2002, 28:386–388.
- Ornstein E, Eidelman LA, Drenger B, Elami A, Pizov R: Systolic pressure variation predicts the response to acute blood loss. J Clin Anesth 1998, 10:137–140.
- Pizov R, Cohen M, Weiss Y, Sgal E, Cotev S, Perel A: Positive end-exspiratory pressure-induced hemodynamic changes are reflected in the arterial pressure waveform. *Crit Care Med* 1996, 24:1381–1387.
- 10. Rooke GA: Systolic pressure variation as an indicator of hypovolemia. *Curr Opin Anaesthesiol* 1995, 8:511–515.
- Rooke GA, Schwid HA, Shapira Y: The effect of graded hemorrhage and intravascular volume replacement on systolic pressure variation in humans during mechanical and spontaneous ventilation. *Anesth Analg* 1995, 80:925–932.
- Beaussier M, Coriat P, Perel A, Lebret F, Kalfon P, Chemla D, Lienhart A, Viars P: Determinants of systolic pressure variation in patients ventilated after vascular surgery. *J Cardiothorac Vasc Anesth* 1995, 9:547–551.
- 13. Magder S, Georgiadis G, Cheong T: Respiratory variations in right atrial pressure predict the response to fluid challenge. *J Crit Care* 1992, 7:76–85.
- 14. Perel A, Pizov R, Cotev S: Systolic blood pressure variation is a sensitive indicator of hypovolemia in ventilated dogs subjected to graded hemorrhage. *Anesthesiology* 1987, 67:498–502.
- 15. Jardin F, Farcot JC, Gueret P, Prost JF, Ozier Y, Bourdarias JP: Cyclic changes in arterial pulse during respiratory support. *Circulation* 1983, 68:266–274.
- 16. Soubrier S, Daulnier F, Hubert H, Delour P, Lenci H, Onimus T, Nseir S, Durocher A: Can dynamic indicators help the prediction of fluid responsiveness in spontaneously breathing critically ill patients? *Intensive Care Med* 2007, 33:1117–1124.
- 17. Monnet X, Rienzo M, Osman D, Anguel N, Richard C, Pinsky MR, Teboul JL: Passive leg raising predicts fluid responsiveness in the critically ill. *Crit Care Med* 2006, 34:1402–1407.
- 18. Heenen S, De Backer D, Vincent JL: How can the response to volume expansion in patients with spontaneous respiratory movements be predicted? *Crit Care* 2006, 10:R102.
- 19. Kramer A, Zygun D, Hawes H, Easton P, Ferland A: Pulse pressure variation predicts fluid responsiveness following coronary artery bypass surgery. *Chest* 2004, 126:1563–1568.
- Lakhal K, Ehrmann S, Benzekri-Lefévre D, Runge I, Legras A, Dequin PF, Mercier E, Wolff M, Régnier B, Boulain T: Respiratory pulse pressure variation fails to predict fluid responsiveness in acute respiratory distress syndrome. *Crit Care* 2011, 15:R85.
- 21. Muller L, Louart G, Bousquet PJ, Candela D, Zoric L, Coussaye JE, Jaber S, Lefrant JY: The influence of the airway pressure on pulsed pressure variation as a predictor of fluid responsiveness. *Intensive Care Med* 2010, 36:496–503.
- 22. Vallée F, Richard JCM, Mari A, Gallas T, Arsac E, Verlaan PS, Chousterman B, Samii K, Genestal M, Fourcade O: Pulse pressure variations adjusted by alveolar driving pressure to assess fluid responsiveness. *Intensive Care Med* 2009, 35:1004–1010.

- 23. De Backer D, Taccone FS, Holsten R, Ibrahimi F, Vincent JL: Influence of respiratory rate on stroke volume variation in mechanically ventilated patients. *Anesthesiology* 2009, 110:1092–1097.
- 24. Charron C, Fessenmeyer C, Cosson C, Mazoit JX, Hebert JL, Benhamou D, Edouard AR: The influence of tidal volume on the dynamic variables of fluid responsiveness in critically ill patients. *Anesth Analg* 2006, 102:1511–1517.
- 25. De Backer D, Heenen S, Koch M, Vincent JL: Pulse pressure variations to predict fluid responsiveness: influence of tidal volume. *Intensive Care Med* 2005, 31:517–523.
- 26. Michard F, Chemla D, Richard C, Wysocki M, Pinsky MR, Lecarpentier Y, Teboul JL: Clinical use of respiratory changes in arterial pulse pressure to monitor the hemodynamic effect of PEEP. *Am J Respir Crit Care Med* 1999, 159:935–939.
- 27. Kim HK, Pinsky MR: Effect of tidal volume, sampling duration, and cardiac contractility on pulse pressure variation during positive-pressure ventilation. *Crit Care Med* 2008, 36:2858–2862.
- 28. Nouira S, Elatrous S, Dimassi S, Besbes L, Boukef R, Mohamed B, Abroug F: Effects of norepinephrin on static and dynamic preload indicators in experimental hemorrhagic shock. *Crit Care Med* 2005, 33:2339–2343.
- 29. Jardin F: Cyclic changes in arterial pressure during mechanical ventilation. *Intensive Care Med* 2004, 30:1047–1050.
- 30. Cooke HW, Ryan KL, Convertino VA: Lower body negative pressure as a model to study progression to acute hemorrhagic shock in humans. *J Appl Physiol* 2004, 96:1249–1261.
- 31. Trzeciak S, Rivers EP: Clinical manifestations of disordered microcirculatory perfusion in severe sepsis. *Crit Care* 2005, 9(suppl 4):S20-S26.
- Sakr Y, Dubois MJ, De Backer D, Creteur J, Vincent JL: Persistent microcirculatory alterations are associated with organ failure and death in patients with septic shock. *Crit Care Med* 2004, 32:1825–1831.
- 33. De Backer D, Creteur J, Preiser JC, Dubois MJ, Vincent JL: Microvascular blood flow is altered in patients with sepsis. *Am J Respir Crit Care Med* 2002, 166:98–104.
- 34. De Backer D, Hollenberg S, Boerma C, Goedhart P, Büchele G, Ospina-Tascon G, Dobbe I, Ince C: How to evaluate the microcirculation:report of a round table conference. *Crit Care* 2007, 11:R101.
- 35. Bartels AB, Bezemer R, Milstein DMJ, Radder M, Lima A, Cherpanath TGV, Heger M, Karemaker JM, Ince C: The microcirculatory response to compensated hypovolemia in lower body negative pressure model. *Microvasc Res* 2011, 82:374–380.
- Ward KR, Tiba MH, Ryan KL, Filho IPT, Rickards CA, Witten T, Soller BR, Ludwig DA, Convertino VA: Oxygen transport characterization of a human model of progressive hemorrhage. *Resuscitation* 2010, 81:987–993.
- 37. Magder S: Further cautions for the use of ventilatory-induced changes in arterial pressures to predict volume responsiveness. *Crit Care* 2010, 14:197.
- Perel A, Minkovich L, Preisman S, Abiad M, Segal E, Coriat P: Assessing fuid-responsiveness by a standardized ventilatory maneuver: the respiratory systolic variation test. *Anesth Analg* 2005, 100:942–945.

Abbreviations

AUC: Area under the curve; BP: Blood pressure; CVP: Central venous pressure; ECG: Electrocardiogram; FCD: Functional capillary density; GEDVI: Global end diastolic volume index; Hb: Hemoglobin; HR: Heart rate; ITBVI: Intrathoracic blood volume index; LBNP: Lower body negative pressure; LV: Left ventricle; MAP: Mean arterial pressure; MFI: Microvascular flow index; MPCL: Mean perfused capillary lenght; OPS: Orthogonal polarization spectral imaging; Paop: Pulmonary artery occlusion pressure; PcvO2: Central venous oxygen tension; PcvCO2: Central venous CO₂ tension; PEEP: Positive end exspiratory pressure; PP: Pulse pressure; PPV: Pulse pressure variation; PRPV: Proportion of perfused vessels; RAP: Right atrial pressure; RBCV: Red blood cell velocity; ROC: Receiver operating characteristic curve; RR: Respiratory rate; RRI: RR interval (the time of the cardiac cycle); RSVT: Respiratory systemic variation test; RV: Right ventricle; SaO2: Arterial oxygen saturation; SAP: Systolic arterial pressure; ScvO2: Central venous oxygen saturation; SV: Stroke volume; SVV: Stroke volume variation; VBG: Central venous blood gas analysis.

Authors' contributions

ÉZ and LR planned the study design, and contributed to hemodynamic data acquisition and analysis. The manuscript was written by ÉZ. VB, AN, PCs have been involved in coordination, hemodynamic data acquisition and analysis. The investigation and data analysis of the microcirculation was done by JK and IL. All authors were involved in the critical revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgement

The publication is supported by the European Union and co-funded by the European Social Fund. Project title: "Telemedicine-focused research activities on the field of Mathematics, Informatics and Medical sciences". Project number: TÁMOP-4.2.2.A-11/1/KONV-2012–0073.

BMC Anesthesiol. 2013;13(40) © 2013 BioMed Central, Ltd.