EDITORIAL

Open Access

Treatment of central line-associated bloodstream infections



Jérémy Guenezan¹, Bertrand Drugeon^{1,2}, Nicolas Marjanovic¹ and Olivier Mimoz^{1,2,3*}

Keywords: Central venous catheter, Bloodstream infection, Prevention, Antibiotics

Central venous catheters (CVCs) are among the most frequently used medical devices in critically ill patients [1]. Unfortunately, complications are not uncommon despite the many improvements in caregiver behaviour observed over recent decades. Central line-associated bloodstream infection (CLABSI) is probably the most feared complication [2]. It is defined as a primary laboratory-confirmed bloodstream infection in a patient with a central line at the time of, or within 24-h prior to, the onset of symptoms, in cases where the cultured organism is not related to an infection from another site. The central line should be in place for > 2 calendar days on the date of the event, with the day of device placement being day 1. Tens of thousands of patients continue to experience CLABSIs each year in the US, resulting in thousands of deaths each year and billions of dollars in added costs to the US healthcare system [3].

Of paramount importance, CLABSI prevention depends on evidence-based measures. The most effective of these are hand hygiene before catheter insertion and maintenance, full-barrier precautions during catheter insertion, 2% alcoholic chlorhexidine use for skin preparation, femoral vein avoidance whenever possible and prompt removal of useless CVCs [4–6]. The use of care protocols and the training of caregivers must also be considered. Implementation of these bundled measures has been shown in high-quality studies to reduce CLABSI incidence by 50% [7].

On the other hand, the principles of CLABSI treatment are not as well established and are mainly based on expert opinions and the results of old cohort studies [2]. Source control and intravenous administration of

* Correspondence: olivier.mimoz@chu-poitiers.fr

¹Centre Hospitalier Universitaire de Poitiers, Service des Urgences Adultes & SAMU 86-Centre 15, 1 Rue de la Milétrie, 86021 Poitiers Cedex, France ²Université de Poitiers, UFR de Médecine Pharmacie, Poitiers, France Full list of author information is available at the end of the article high doses of bactericidal agents are the most commonly cited measures.

Source control consists of prompt removal of the catheter when clinical suspicion is high, at times even before the CLABSI diagnosis is proven. Yet, catheter salvage with antibiotic lock and systemic antibiotic treatment are not recommended for critically ill patients. <u>Salvage</u> therapy is limited to cases where the CVC cannot be removed, in the absence of septic shock, and for a period of <u>at least 4 weeks</u>, with low success rate.

Empirical treatment should be initiated promptly, before confirmation of the diagnosis, after appropriate cultures have been obtained. The choice is based on host characteristics, known or suspected colonization with resistant microorganisms and local epidemiology and susceptibility patterns [8]. In general, reasonable coverage is required for common Gram-positive and Gram-negative organisms. Parenteral vancomycin (or daptomycin in patients with acute renal failure or when organisms have reduced susceptibility to vancomycin) is the cornerstone of an empirical regime in settings with a high prevalence of methicillin-resistant Staphylococcus aureus (MRSA) [9]. Otherwise, an anti-staphylococcal penicillin or a first-generation cephalosporin is enough [10]. Antibiotics active against Gram-negative bacilli include β-lactam/ β-lactamase inhibitor combinations, fourth-generation cephalosporins and carbapenems alone or in combination with an aminoglycoside [11, 12]. Agents against Pseudomonas aeruginosa are required in cases of neutropaenia, severity of disease or increased probability or known prior colonization [8]. New anti-Gram negative antibiotics are indicated for infections due to multi-drug resistant Gram-negative organisms [13].

Empiric coverage for <u>candidaemia</u> should be considered if <u>multiple sites are colonized</u> with *Candida* or for patients with bone marrow or organ transplants, <u>hematologic ma-</u> lignancy, <u>femoral catheterization</u> or when patients are



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.



receiving total parenteral nutrition or prolonged administration of broad-spectrum antibiotics [2, 8]. Parenteral echinocandins are recommended when resistance to azoles is suspected (prior exposure to azoles in the previous 3 months or high prevalence of common non-*albicans Candida* such as *Candida* glabrata or *Candida* krusei) [8, 12]. Otherwise, intravenous fluconazole is enough.

Once antimicrobial susceptibility patterns are available, de-escalation to appropriate therapy with the narrowest spectrum possible is recommended [14]. The duration of treatment depends on the organism involved and the presence of complications or not, and varies between 7 days to 6 weeks.

For **uncomplicated** infection (i.e., not associated with suppurative thrombosis, endocarditis or metastatic infection) with septic manifestations resolving within 72 h of catheter removal, intravenous antimicrobial therapy is recommended for the following duration based on the organism isolated (Fig. 1):

- Staphylococcus aureus: 14 days
- <u>Coagulase-negative staphylococci: 7 days</u>
- <u>Enterococci and Gram-negative bacilli: 10 to 14 days</u>
- <u>Candida: 14 days</u>

Recently, a retrospective cohort study of 366 uncomplicated CLBSI cases added new scientific evidence on the subject [15]. Duration of antimicrobial treatment lower than 15 days was associated with earlier mortality or recurrence, suggesting that antibiotic treatment should be <u>continued for at least 2 weeks for all microorganisms</u>. Unfortunately, the study was not powerful enough to compare outcomes amongst individual classes of causative organisms. The authors concluded that further evidence is required before amending the guidelines.

Transoesophageal echography should be performed in patients with <u>S. aureus</u>, enterococci and <u>Candida</u> infections so as to rule out complicated forms of CLBSI [2]. Similarly, an ophthalmological examination is recommended in candidaemic patients [2]. Indeed, the duration of parenteral antibiotic therapy should be longer in the case of <u>complicated CLABSIs</u>, ranging from <u>4 to 6 weeks</u>, even though there is little scientific evidence to support these recommendations [14]. Suppurative thrombophlebitis may also require anticoagulation and sometimes surgery. Infectious endocarditis and osteomyelitis may require surgical treatment.

Conclusion

The treatment of CLABSIs in 2018 is still based on expert opinions and cohort studies rather than robust scientific evidence. Such infections continue to increase complications and mortality and are quite costly. Large-scale studies are warranted to further clarify the nature and duration of treatment. In the meantime, physicians are encouraged to use algorithms such as the one proposed here (Fig. 1).

Abbreviations

CLABSI: Central line-associated bloodstream infection; CVC: Central venous catheter

Acknowledgements

Not applicable.

Funding

None.

Availability of data and materials

Not applicable.

Authors' contributions

JG wrote the first version of the manuscript. All authors have reviewed and completed the manuscript. All authors read and approved the final manuscript.

Author's information

OM is professor of Emergency Medicine and Chief of the Emergency Department at University Hospital of Poitiers in France. JG, NM and BD are physicians in the Emergency Department at University

Hospital of Poitiers in France.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable.

Competing interests

Dr. Mimoz received grant support from 3 M and Becton Dickinson and honoraria for giving lectures from 3M and Becton Dickinson.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Centre Hospitalier Universitaire de Poitiers, Service des Urgences Adultes & SAMU 86-Centre 15, 1 Rue de la Milétrie, 86021 Poitiers Cedex, France. ²Université de Poitiers, UFR de Médecine Pharmacie, Poitiers, France. ³INSERM, U1070, Pharmacologie des Agents anti-Infectieux, Poitiers, France.

Received: 17 October 2018 Accepted: 29 October 2018 Published online: 16 November 2018

References

- Frasca D, Dahyot-Fizelier C, Mimoz O. Prevention of central venous catheterrelated infection in the intensive care unit. Crit Care. 2010;14(2):212.
- Timsit JF, Rupp M, Bouza E, Chopra V, Kärpänen T, Laupland K, Lisboa T, Mermel L, Mimoz O, Parienti JJ, Poulakou G, Souweine B, Zingg W. A state of the art review on optimal practices to prevent, recognize, and manage complications associated with intravascular devices in the critically ill. Intensive Care Med. 2018; in press.
- Zimlichman E, Henderson D, Tamir O, Franz C, Song P, Yamin CK, Keohane C, Denham CR, Bates DW. Health care-associated infections: a meta-analysis of costs and financial impact on the US health care system. JAMA Intern Med. 2013;173(22):2039–46.
- 4. Mimoz O, Lucet JC, Kerforne T, Pascal J, Souweine B, Goudet V, Mercat A, Bouadma L, Lasocki S, Alfandari S, Friggeri A, Wallet F, Allou N, Ruckly S, Balayn D, Lepape A, Timsit JF, CLEAN trial investigators. Skin antisepsis with chlorhexidine–alcohol versus povidone iodine– alcohol, with and without skin scrubbing, for prevention of intravascular catheter-related infection (CLEAN): an open-label, multicentre, randomised, controlled, two-by-two factorial trial. Lancet. 2015;386(10008):2069–77.
- Mimoz O, Chopra V, Timsit JF. What's new in catheter-related infection: skin cleansing and skin antisepsis. Intensive Care Med. 2016;42(11):1784–6.
- Mimoz O, Chopra V, Widmer A. What's new in skin antisepsis for short-term intravascular catheters: new data to address old problems? Intensive Care Med. 2016;42(12):2043–5.
- Blot K, Bergs J, Vogelaers D, Blot S, Vandijck D. Prevention of central lineassociated bloodstream infections through quality improvement interventions: a systematic review and meta-analysis. Clin Infect Dis. 2014;59(1):96–105.
- Mermel LA, Allon M, Bouza E, Craven DE, Flynn P, O'Grady NP, Raad II, Rijnders BJ, Sherertz RJ, Warren DK. Clinical practice guidelines for the diagnosis and

management of intravascular catheter-related infection: 2009 update by the Infectious Diseases Society of America. Clin Infect Dis. 2009;49(1):1–45.

- Timsit JF, L'Heriteau F, Lepape A, Francais A, Ruckly S, Venier AG, Jarno P, Boussat S, Coignard B, Savey A. A multicentre analysis of catheter- related infection based on a hierarchical model. Intensive Care Med. 2012;38(10):1662–72.
- Chaves F, Garnacho-Montero J, Del Pozo JL, Bouza E, Capdevila JA, de Cueto M, Domínguez MÁ, Esteban J, Fernández-Hidalgo N, Fernández Sampedro M, Fortún J, Guembe M, Lorente L, Paño JR, Ramírez P, Salavert M, Sánchez M, Vallés J. Diagnosis and treatment of catheter-related bloodstream infection: Clinical guidelines of the Spanish Society of Infectious Diseases and Clinical Microbiology and (SEIMC) and the Spanish Society of Spanish Society of Intensive and Critical Care Medicine and Coronary Units (SEMICYUC). Med Int. 2018;42(1):5–36.
- Lorente L, Jimenez A, Santana M, Iribarren JL, Jimenez JJ, Martin MM, Mora ML. Microorganisms responsible for intravascular catheter- related bloodstream infection according to the catheter site. Crit Care Med. 2007;35(10):2424–7.
- Safdar N, Handelsman J, Maki DG. Does combination antimicrobial therapy reduce mortality in Gram-negative bacteraemia? A meta-analysis. Lancet Infect Dis. 2004;4(8):519–27.
- 13. Akova M, Daikos GL, Tzouvelekis L, Carmeli Y. Interventional strategies and current clinical experience with carbapenemase-producing gram-negative bacteria. Clin Microbiol Infect. 2012;18(5):439–48.
- 14. Rupp ME, Karnatak R. Intravascular catheter-related bloodstream infections. Infect Dis Clin N Am. 2018; in press.
- Huerta LE, Nelson GE, Stewart TG, Rice TW. Factors associated with recurrence and mortality in central line-associated bloodstream infections: a retrospective cohort study. Crit Care. 2018;22(1):266.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- · thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

