

Critical Care 2



Ethics and end-of-life care for adults in the intensive care unit

J Randall Curtis, Jean-Louis Vincent

The intensive care unit (ICU) is where patients are given some of the most technologically advanced life-sustaining treatments, and where difficult decisions are made about the usefulness of such treatments. The substantial regional variability in these ethical decisions is a result of many factors, including religious and cultural beliefs. Because most critically ill patients lack the capacity to make decisions, family and other individuals often act as the surrogate decision makers, and in many regions communication between the clinician and family is central to decision making in the ICU. Elsewhere, involvement of the family is reduced and that of the physicians is increased. End-of-life care is associated with increased burnout and distress among clinicians working in the ICU. Since many deaths in the ICU are preceded by a decision to withhold or withdraw life support, high-quality decision making and end-of-life care are essential in all regions, and can improve patient and family outcomes, and also retention of clinicians working in the ICU. To make such a decision requires adequate training, good communication between the clinician and family, and the collaboration of a well functioning interdisciplinary team.

Introduction

Critical care is an integral part of hospital care, and the intensive care unit (ICU) is the setting where patients are given the most technologically advanced life-sustaining treatments. These treatments are expensive and resource-intensive, but can sustain life despite severe and multiple organ dysfunction. The ICU is, however, also a setting where death is common and end-of-life care is frequently provided. Since the focus in ICUs is on sustaining life, the delivery of high-quality end-of-life care can be particularly challenging, and clinicians often find the dual responsibilities of saving lives and delivering end-of-life care difficult. Because of the nature of critical care, difficult decisions often need to be made about the usefulness of life-sustaining treatments, not only in terms of the probability of survival but also the quality of life associated with survival. Difficult decisions also need to be made about the fairness of expending substantial resources on one patient.¹ Furthermore, because most critically ill patients do not have the capacity to make decisions, the family frequently becomes involved in discussions about the goals of care and often represents the values and preferences of the patient.² The extent to which the family is directly involved in such decisions varies according to the countries and cultures:³ however, good communication between the clinician and family is essential for quality end-of-life care in the ICU, irrespective of the location.⁴

Data from observational studies indicate that end-of-life care in the ICU varies greatly between countries.⁵⁻⁹ The reasons for this variability have not been clearly defined, but are probably diverse, including differences in religion,¹⁰ legislation and culture,^{10,11} organisation of care in the ICU,^{7,12,13} attitudes of physicians toward end-of-life care,¹⁴ severity of illness and casemix,^{7,15} and the physician's predictions of

prognosis and future quality of life.¹⁵ Variability also exists within countries¹⁶⁻¹⁹ and between intensivists within hospitals.²⁰

Admissions and triage decisions

The availability of ICU resources vary substantially in different countries, and decisions about admission, triage, and end-of-life care vary accordingly. For example, ICU care is not available in many countries in the developing world and in rural regions of developed countries. Even where ICU care is available, the proportion of hospital beds that are ICU beds differs between centres.²¹ The availability of beds in the ICU will, by necessity, affect decisions about indications for care in the ICU; this assertion is supported by an association between the mortality rate in the ICU and availability of beds in the ICU.^{21,22} However, the availability of beds should not affect the ethical principles that guide the use of intensive care.

An integral part of the ethics of critical care is the process used for decisions about who needs treatment in the ICU, and when that treatment is no longer indicated. Treatment might be judged to be not indicated because patients are not sick enough for care in the ICU, or

Published Online

October 9, 2010

DOI:10.1016/S0140-

6736(10)60143-2

See Online/Comment

DOI:10.1016/S0140-

6736(10)61502-4

This is the second in a **Series** of three papers about critical care

Division of Pulmonary and Critical Care Medicine, Harborview Medical Center; University of Washington, Seattle, WA, USA (Prof J R Curtis MD); and Department of Intensive Care, Erasme Hospital; Université Libre de Bruxelles, Brussels, Belgium (Prof J-L Vincent MD)

Correspondence to:

Prof J Randall Curtis, Division of Pulmonary and Critical Care Medicine Harborview Medical Center, Box 359762, University of Washington, 325 Ninth Avenue, Seattle, WA 98104-2499, USA
jrc@u.washington.edu

Search strategy and selection criteria

We searched the Cochrane Library (1994–2009), Medline (1994–2009), and Embase (1994–2009) for papers published in English, using the search terms “ethics”, “end-of-life care”, or “palliative care” in combination with “critical care” or “intensive care”. We predominantly selected publications from the past 5 years, but did not exclude commonly referenced and highly regarded older publications. We also searched the reference lists of articles identified by this search strategy and selected those we judged relevant.

Panel 1: Principles of fair allocation of resources in the intensive care unit (ICU)

- 1 Every individual's life is valuable and equally so
- 2 Respect for patient autonomy, as represented by informed consent, is a central tenet for provision of health care, including care in the ICU
- 3 Enhancement of the patient's welfare, by provision of resources that meet an individual's medical needs and that the patient regards as beneficial, is the main duty of health-care providers
- 4 ICU care, when medically appropriate, is an essential component of a basic package of health-care services that should be available for all patients
- 5 The duty of health-care providers to provide benefit to a patient has limitations when doing so unfairly compromises the availability of resources needed by other patients

Reprinted with the permission of the American Thoracic Society.¹ Copyright American Thoracic Society.




Figure: Three-step approach to patient-centred and family-centred decision making

Adapted from Curtis and White⁴ with permission from the American College of Chest Physicians.

because they are too sick and such care is unlikely to provide benefit. The American Thoracic Society has outlined several important principles that should guide decision making about admission and triage.¹ The main duty of the ICU team is to ensure patient welfare and that care in the ICU, when appropriate, constitutes basic medical care (panel 1). The duty of the ICU team to provide benefit to a patient has limitations when provision of care for that patient unfairly compromises care for others.

The recommendation that every ICU should have explicit and written criteria for admission and discharge of the patient is supported by statements from critical care professional societies.^{1,23,24} Most ICUs do have such criteria, but they generally require interpretation in the application to individual patients, and intensivists report that these criteria are not explicitly used to decide admission or triage status for most patients.²¹

Availability of ICU resources in a region will have important effects on decisions about admission and triage. Importantly, these decisions should be governed by ethical principles, irrespective of the availability of beds in the ICU.

The ageing population in many countries will increase the importance of these issues in the future. In the USA, for example, the proportion of all deaths that occur in the ICU is nearly 20%, and this proportion does not decrease with increasing age until after 85 years.²⁵ The proportion of all deaths in elderly people that are preceded by cardiopulmonary resuscitation is increasing.²⁶ Society and countries need to develop approaches to address the appropriate delivery of critical care to the increasing population of elderly people, especially those with chronic life-limiting disease. These approaches will probably vary according to the country and health-care system, but the ethical principles should be similar and many regions will have similar disparities that need to be addressed.

Communication about end-of-life care in the outpatient setting between physicians and patients with life-limiting disease is not associated with patient distress, and is associated with a reduction in use of unsuccessful life-sustaining treatments, improved quality of life, and reduction in health-care costs at the end of life.^{27,28} Advance directives and advance care planning have recently been shown to be associated with patients receiving care matching their preferences, and have also been shown to be associated with less aggressive care at the end of life and with better family ratings of end-of-life care.^{29–32} Although these discussions are generally not the purview of intensivists, efforts to increase the quality and quantity of these discussions when patients are stable are likely to improve our ability to maximise the effectiveness of intensive care, and reduce the burden of end-of-life care in the ICU on patients, families, and the health-care system.

End-of-life decisions

Interdisciplinary communication

End-of-life care in most settings is delivered by an interdisciplinary team that includes nurses and physicians. Ideally, end-of-life decisions should be made after discussions between all members of the interdisciplinary team. However, interdisciplinary collaboration about end-of-life care is often poor and varies across countries.^{33,34} For example, in a prospective survey in 113 French ICUs, end-of-life decisions were made by one physician in 12% of cases, by medical staff alone in 34% of cases, and by medical and nursing staff in 54% of patients.¹⁸ In a study of an ICU in Lebanon, nurses were not involved in 26% of end-of-life decisions.³⁵ In a questionnaire study of 1961 intensivists from 21 countries, for a hypothetical patient without any family, 62% of physicians from northern and central Europe would involve nurses in end-of-life discussions compared with only 32% of physicians in southern Europe, 39% in Japan, 38% in Brazil, and

29% in the USA.⁹ Patients and families report that interdisciplinary collaboration is an essential part of good end-of-life care.³⁶ Poor interdisciplinary collaboration about end-of-life care is associated with increased symptoms of burnout, depression, and post-traumatic stress among clinicians working in the ICU.^{37–39} Additionally, conflict between clinicians in the ICU is common, is increased with delivery of end-of-life care, and is associated with increased job stress.^{40,41} Therefore, improved interdisciplinary collaboration for end-of-life care in the ICU is important for improvement of the quality of care and the work environment for clinicians.

Communication between clinician and family

Physicians caring for critically ill individuals have an obligation to disclose information about a patient's condition and prognosis to the patient and the patient's family. The families of critically ill individuals are an important source of information about the patient's values and treatment preferences. In 2005, five international critical-care societies issued a consensus statement advocating shared decision making about life-sustaining treatment in ICUs.⁴² In this statement, a shared decision is defined as one in which "responsibility for decisions is shared jointly by the treating physician and the patient's family".⁴² Guidelines for end-of-life care also emphasise the importance of involving the patient (when possible) and the family.⁴³ Nevertheless, substantial international differences exist in the amount of involvement of the patient and family in the end-of-life process. Moreover, although most families want the physician and ICU team to provide a recommendation about whether to restrict life support and then want to share in the final decision,^{44,45} some families do not wish to be involved in such decisions or want to make decisions without a recommendation from the physician.^{46–48}

Traditionally, families have been much more involved in end-of-life decision making in the USA than in Europe.⁴⁹ In the Ethicus study,⁶ done in 37 ICUs in 17 European countries, end-of-life decisions were discussed with the family more commonly in northern (84%) and central (66%) than in southern (47%) Europe. Huge variations have been reported in family involvement—from 100% in India,⁵⁰ 98% in Hong Kong,⁵¹ 79% in Lebanon,⁵² 72% in Spain,¹⁹ to just 44% in France.¹⁸ In a questionnaire study of intensivists in Italy, 19% of physicians said the close family were never involved in such decisions and 56% would never involve patients even if competent.⁵²

Physicians need to be aware of the variety and complexity of attitudes present in our increasingly multicultural society, and adapt their approach to the situation. There is a range for physician's role in decision making from parentalism in which the physician makes the decision to autonomy where the patient or family makes the decision with shared decision-making in the middle.⁵³ The figure shows a potential approach to match the clinician's role with the needs of the patient and the patient's family.

Panel 2: Proposed best practices for ethics and end-of-life care in the intensive care unit (ICU)

ICU policies

- Explicit policy for admission and transfer criteria
- Explicit policy for delivery of palliative and end-of-life care
- Training for ICU clinicians in ethics, communication, and end-of-life care

Interdisciplinary communication

- All patients included in daily interdisciplinary rounds
- Daily interdisciplinary communication about the goals of care

Communication with families

- Open visiting hours for family members
- Allow family presence during ICU rounds
- Routine interdisciplinary family conferences within 48–72 h for family of patients at high risk of death or prolonged stay

Withholding or withdrawing life support

- Explicit policy about withholding and withdrawing life support
- Protocol for withdrawing life support
- Train clinicians from all disciplines regarding ethics and communication about withholding and withdrawing life support
- Staff support programmes about end-of-life care, and withholding or withdrawing life support

Shared decision making is the default position that is modified in three steps.⁴ First, as the prognosis worsens and the certainty of the prognosis increases, so should the physician's willingness to take on the burden of making a decision. Second, the preferred decision-making role of the family is assessed. Last, the approach is adapted to the patient and family factors identified in the first and second steps. For this approach to work, communication between physicians and the family needs to be optimum. However, communication between the clinician and the family in the ICU is often inadequate; in one study, only half of families of patients in the ICU understood basic information about patients' diagnoses, prognoses, or treatments after discussion with clinicians.⁵⁴

A focus on communication with the families of all critically ill patients is important, not just those expected to die. Whether critically ill patients will survive is often not clear at the time when communication between the clinician and family should be happening. Additionally, although the patient's death is a risk factor for psychological symptoms among the family, the families of patients who survive are also at increased risk of these symptoms.⁵⁵ Families of patients who survive are less satisfied with communication from ICU clinicians than are those of patients who die.⁵⁶

Discussions between clinicians in the ICU and family about goals of care and medical decision making often take place during conferences between the inter-

Panel 3: Key stages for improvement of communication during interdisciplinary family conferences in the intensive care unit

Before

- Plan the specific details of location and setting: quiet, private place⁵⁷
- Do a preconference with the clinicians to develop consensus and ensure consistency of information provided

During

- Use active listening and provide family adequate time to speak⁵⁸
- Use empathic statements to provide support for families:⁵⁹ difficulty of having a critically ill loved one; difficulty of surrogate decision making; impending loss of a loved one
- Acknowledge and address family emotions⁶⁰
- Explore and focus on patient values and treatment preferences⁶⁰
- Affirm non-abandonment of patient and family⁶¹

Completion

- Summarise information and decisions
- Ask for questions, and allow family time to consider questions⁶²
- Reaffirm and support family with respect to their decisions⁶²

disciplinary team and family (panel 2). Features of these conferences that are associated with improved family experience or assessment of communication have been identified in several studies (panel 3). For example, improved outcomes are associated with a private place for family communication and with consistent communication by all members of the team.⁵⁷ Family are more satisfied when clinicians spend more time listening and less time talking.⁵⁸ Other features of clinician communication associated with improved family experiences include assurances that the patient will not be abandoned before death; assurances that the patient will not suffer; and explicit support for family decisions.⁶² Empathic statements by clinicians are also associated with increased family satisfaction.⁵⁹ When there is conflict between staff and families, ethics consultations have been beneficial.⁶³

When communication occurs across cultures or languages, the likelihood of miscommunication is increased.⁶⁴ Involvement of family-specific religious or community leaders and professional interpreters could be helpful.⁶⁴ Errors in communication are common even with professional medical interpreters and might affect understanding, decision making, and emotional support.^{65,66} Some simple steps can improve this communication: clinicians can meet briefly with interpreters before the conference, speak slowly allowing time for interpretation, restrict the number of simultaneous conversations, and use pictures or drawings when possible.⁶⁷ Another important part of care in the ICU

is to assess the spiritual needs of the families and then offer them spiritual care if desired. Family satisfaction with care is increased if spiritual care needs are assessed, and spiritual care is provided by a spiritual-care provider.^{68,69}

Withholding or withdrawing life support

Most patients who die in ICUs do so after a decision has been made to restrict life-sustaining treatments,^{6-8,70} but there are substantial differences in the proportion of deaths preceded by withholding or withdrawing life support internationally. In the Ethicus study,⁶ withdrawal of life-sustaining treatments was reported as more common (47% vs 18%, $p < 0.001$) in northern European countries (Denmark, Finland, Ireland, Netherlands, Sweden, and UK) than in those in southern Europe (Greece, Israel, Italy, Portugal, Spain, and Turkey). In an analysis of 14488 patients from 282 ICUs in seven different geographical regions, deaths occurring after a decision to restrict life-sustaining treatments vary from 26% in Central and South America to 48% in central and western Europe.⁷ Life support was withdrawn or withheld in 59% of patients who did not survive in Hong Kong,⁵¹ 53% in France,¹⁸ 45% in Lebanon,³⁵ 41% in Sweden,⁷¹ 35% in Spain,¹⁹ and 49% in India.⁴⁷ Differences in religious and cultural backgrounds are likely to be one of the major reasons for these international differences.

Although many ethicists and critical-care societies state that there is no ethical distinction between withholding or withdrawing life-sustaining treatments,^{43,72,73} this perspective is not universally accepted, and some ethicists and intensivists believe that an important distinction exists between withholding and withdrawing life-sustaining treatments.^{74,75} In Israel, for example, orthodox Jewish law allows life-sustaining treatments to be withheld, but withdrawal of continuous interventions is forbidden because it is regarded as an act to shorten life.⁷⁶ However, withdrawal of intermittent life-sustaining treatments is permitted because it is seen as the next treatment being withheld rather than the withdrawal of the present one.^{76,77}

Religion is an important determinant of attitudes toward dying, death, and end-of-life care, and includes the religion of patients, their families, and their clinicians. For example, in the Ethicus study,¹¹ treatment was withheld more often than it was withdrawn if the physician was Jewish (81%), Greek orthodox (78%), or Muslim (63%), whereas withdrawal occurred more often when physicians were Catholic (53%), Protestant (49%), or had no religious affiliation (47%). Religion is also an important determinant of acceptance of brain death, a state that is widely, but not universally, accepted.^{6,10}

With the large numbers of deaths now associated with a decision to withhold or withdraw life-sustaining treatments, improvement of the process by which life-sustaining treatments are withheld or withdrawn is an important aspect of improving quality of ICU care.⁴³ There are few data to guide clinicians in the practical

aspects of withdrawing life-sustaining treatments.⁷⁸ Withdrawal of these treatments is a clinical procedure that deserves the same preparation and expectation of quality as do other procedures.⁴³ These decisions can become routine for clinicians working in the ICU, and, as such, clinicians must be careful to guard against the subtle institutional pressures to withdraw life-sustaining treatments.⁷⁹ Rationale for the decision to withdraw life support should be noted in the medical record. An explicit plan for the procedure should be developed: the patient should be in the appropriate setting with irrelevant monitoring removed; the process should be carefully documented in the medical record, including the reasons why sedation or analgesia was increased; and outcomes should be assessed. The plan should also be discussed thoroughly with the patient (if possible) and family to ensure they understand the planned process, potential symptoms, and the plan for treatment of symptoms.⁸⁰

Once a decision is made to withdraw life-sustaining treatments, the time during which a treatment is withdrawn should be determined by the potential for discomfort as treatment is stopped. The only legitimate rationale why life-sustaining treatment is tapered in this setting is to allow time to treat patient's symptoms. Mechanical ventilation is one of the few life-support treatments in which abrupt termination causes discomfort (panel 4). Typically, the transition from full ventilatory support to T piece or extubation should take less than 10–20 min. Drugs, including opioids and benzodiazepines, are often used to treat the patient's distress or discomfort and there is some observational evidence to suggest that appropriate use of these drugs does not hasten death.^{81,82} Few data exist to support whether patients should be extubated after terminal discontinuation of mechanical ventilation. No significant difference was noted in patient comfort in small studies, which lacked power to detect clinically important differences.⁸³ Families rate quality of dying higher when patients are extubated, but firm conclusions cannot be drawn because of the observational nature of such studies.⁸⁴ The decision to extubate should, therefore, be made on an individual basis, depending on the anticipated time to death and family preferences about the presence of an endotracheal tube and the potential for distressing respiratory sounds.

Like many aspects of critical care, a protocol to withhold or withdraw life-sustaining treatments, if carefully developed to accommodate local standards, could provide an opportunity to improve care and reduce inappropriate variability in care. An ICU order form for withdrawal of life-support that was assessed in a before and after study included preparations before withdrawal of life support (such as discontinuation of routine laboratory tests), and protocols for analgesia and sedation in this context, and ventilator withdrawal.⁸⁵ Physicians and nurses thought that the order form was helpful, and implementation was associated with increased use of benzodiazepine and

Panel 4: An eight-step approach to terminal withdrawal of mechanical ventilation

- 1 Communicate withdrawal process with patient (if able to communicate), patient's family, and intensive care unit team, and document decision making and communication in the medical record
- 2 Explain to patient and family the likely time course, uncertainty of that time course, and possible symptoms and treatment plan for symptoms when withdrawing life-sustaining treatment
- 3 Assess patient comfort and treat discomfort as necessary, often with opiate or benzodiazepine drugs, depending on the symptoms identified
- 4 Remove positive end expiratory pressure, and turn inspired oxygen fraction to room air
- 5 Reassess patient's comfort, and treat discomfort as necessary
- 6 Reduce ventilatory support by about 50%
- 7 Repeat steps 5 and 6 until ventilator support has been discontinued
- 8 Remove mechanical ventilator, and extubate patient or attach T-piece with humidified air

opiate drugs in the hour before and the hour after ventilator withdrawal, but without an associated reduction in time from ventilator withdrawal to death, suggesting this approach can increase drug use for patient comfort without hastening death.

Conclusions

There is substantial regional and international variability in the approaches to end-of-life care. Some of this variability will diminish as we develop a global consensus about the ethics of critical care, but some variability will inevitably remain because of regional variation in religious and cultural perspectives about end-of-life care, and also variation in the availability of ICU resources. Development of global consensus about end-of-life care, to the extent possible, will require open and continued discussion of these issues in international forums. In all regions, the delivery of ethical and high-quality critical care requires training and emphasis on ethical decision making, communication and collaboration throughout the interdisciplinary team, effective communication with patients and families, and identification and resolution of conflict within the team and with patients and families.

Contributors

JRC did the literature search. JRC and J-LV did the literature review, and drafted and reviewed this report.

Conflicts of interest

We declare that we have no conflicts of interest.

Acknowledgments

Funding was provided by a grant from the National Institute of Nursing Research (R01NR-05226 to JRC). The sponsor was not involved in developing, drafting, or reviewing the report, or in the decision to publish.

References

- 1 American Thoracic Society. Fair allocation of intensive care unit resources. *Am J Respir Crit Care Med* 1997; **156**: 1282–301.
- 2 Prendergast TJ, Luce JM. Increasing incidence of withholding and withdrawal of life support from the critically ill. *Am J Respir Crit Care Med* 1997; **155**: 15–20.
- 3 Cohen S, Sprung C, Sjøkvist P, et al. Communication of end-of-life decisions in European intensive care units. *Intensive Care Med* 2005; **31**: 1215–21.

- 4 Curtis JR, White DB. Practical guidance for evidence-based ICU family conferences. *Chest* 2008; **134**: 835–43.
- 5 Vincent JL. Forgoing life support in western European intensive care units: results of an ethical questionnaire. *Crit Care Med* 1999; **16**: 1626–33.
- 6 Sprung CL, Cohen SL, Sjøkvist P, et al. End-of-life practices in European intensive care units: the Ethicus Study. *JAMA* 2003; **290**: 790–97.
- 7 Azoulay E, Metnitz B, Sprung CL, et al. End-of-life practices in 282 intensive care units: data from the SAPS 3 database. *Intensive Care Med* 2009; **35**: 623–30.
- 8 van der Heide A, Deliens L, Faisst K, et al. End-of-life decision-making in six European countries: descriptive study. *Lancet* 2003; **362**: 345–50.
- 9 Yaguchi A, Truog RD, Curtis JR, et al. International Differences in End-of-Life Attitudes in the Intensive Care Unit: Results of a Survey. *Arch Intern Med* 2005; **165**: 1970–75.
- 10 Bulow HH, Sprung CL, Reinhart K, et al. The world's major religions' points of view on end-of-life decisions in the intensive care unit. *Intensive Care Med* 2008; **34**: 423–30.
- 11 Sprung CL, Maia P, Bulow HH, et al. The importance of religious affiliation and culture on end-of-life decisions in European intensive care units. *Intensive Care Med* 2007; **33**: 1732–39.
- 12 Keenan SP, Busche KD, Chen LM, Esmail R, Inman KJ, Sibbald WJ. Withdrawal and withholding of life support in the intensive care unit: a comparison of teaching and community hospitals. *Crit Care Med* 1998; **26**: 245–51.
- 13 Kollef MH, Ward S. The influence of access to a private attending physician on the withdrawal of life-sustaining therapies in the intensive care unit. *Crit Care Med* 1999; **27**: 2125–32.
- 14 Cook DJ, Guyatt GH, Jaeschke R, et al. Determinants in Canadian health care workers of the decision to withdraw life support from the critically ill. *JAMA* 1995; **273**: 703–08.
- 15 Cook D, Rocker G, Marshall J, et al. Withdrawal of mechanical ventilation in anticipation of death in the intensive care unit. *N Engl J Med* 2003; **349**: 1123–32.
- 16 Prendergast TJ, Claessens MT, Luce JM. A national survey of end-of-life care for critically ill patients. *Am J Respir Crit Care Med* 1998; **158**: 1163–67.
- 17 Wunsch H, Harrison DA, Harvey S, Rowan K. End-of-life decisions: a cohort study of the withdrawal of all active treatment in intensive care units in the United Kingdom. *Intensive Care Med* 2005; **31**: 823–31.
- 18 Ferrand E, Robert R, Ingrand P, Lemaire F, for the French LATAREA Group. Withholding and withdrawal of life support in intensive-care units in France: a prospective study. *Lancet* 2001; **357**: 9–14.
- 19 Esteban A, Gordo F, Solsona JF, et al. Withdrawing and withholding life support in the intensive care unit: a Spanish prospective multicenter observational study. *Intensive Care Med* 2001; **27**: 1744–49.
- 20 Garland A, Connors AF. Physicians' influence over decisions to forego life support. *J Palliat Med* 2007; **10**: 1298–305.
- 21 Wunsch H, Angus DC, Harrison DA, et al. Variation in critical care services across North America and Western Europe. *Crit Care Med* 2008; **36**: 2787–93.
- 22 Vincent JL, Suter P, Bihari D, Bruining H. Organization of intensive care units in Europe: lessons from the EPIC study. *Intensive Care Med* 1997; **23**: 1181–84.
- 23 Guidelines for intensive care unit admission, discharge, and triage. Task Force of the American College of Critical Care Medicine, Society of Critical Care Medicine. *Crit Care Med* 1999; **27**: 633–38.
- 24 Department of Health National Executive. Guidelines on admission to and discharge from intensive care and high dependency units. London: Department of Health, 1996.
- 25 Angus DC, Barnato AE, Linde-Zwirble WT, et al on behalf of the Robert Wood Johnson Foundation ICU End-of-Life Peer Group. Use of intensive care at the end of life in the United States: an epidemiologic study. *Crit Care Med* 2004; **32**: 638–43.
- 26 Ehlenbach WJ, Barnato AE, Curtis JR, et al. Epidemiologic study of in-hospital cardiopulmonary resuscitation in the elderly. *N Engl J Med* 2009; **361**: 22–31.
- 27 Wright AA, Zhang B, Ray A, et al. Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA* 2008; **300**: 1665–73.
- 28 Zhang B, Wright AA, Huskamp HA, et al. Health care costs in the last week of life: associations with end-of-life conversations. *Arch Intern Med* 2009; **169**: 480–88.
- 29 Silveira MJ, Kim SY, Langa KM. Advance directives and outcomes of surrogate decision making before death. *N Engl J Med* 2010; **362**: 1211–18.
- 30 Teno JM, Gruneir A, Schwartz Z, Nanda A, Wetle T. Association between advance directives and quality of end-of-life care: a national study. *J Am Geriatr Soc* 2007; **55**: 189–94.
- 31 Norris K, Merriman MP, Curtis JR, Asp C, Tuholske L, Byock IR. Next of kin perspectives on the experience of end-of-life care in a community setting. *J Palliat Med* 2007; **10**: 1101–15.
- 32 Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ* 2010; **340**: c1345.
- 33 Ferrand E, Lemaire F, Regnier B, et al. Discrepancies between perceptions by physicians and nursing staff of intensive care unit end-of-life decisions. *Am J Respir Crit Care Med* 2003; **167**: 1310–15.
- 34 Benbenishty J, Ganz FD, Lippert A, et al. Nurse involvement in end-of-life decision making: the ETHICUS Study. *Intensive Care Med* 2006; **32**: 129–32.
- 35 Yazigi A, Riachi M, Dabbar G. Withholding and withdrawal of life-sustaining treatment in a Lebanese intensive care unit: a prospective observational study. *Intensive Care Med* 2005; **31**: 562–67.
- 36 Carline JD, Curtis JR, Wenrich MD, Shannon SE, Ambrozio DM, Ramsey PG. Physicians' interactions with health care teams and systems in the care of dying patients: perspectives of dying patients, family members, and health care professionals. *J Pain Symptom Manage* 2003; **25**: 19–28.
- 37 Poncet MC, Toullic P, Papazian L, et al. Burnout syndrome in critical care nursing staff. *Am J Respir Crit Care Med* 2007; **175**: 698–704.
- 38 Embriaco N, Azoulay E, Barrau K, et al. High level of burnout in intensivists: prevalence and associated factors. *Am J Respir Crit Care Med* 2007; **175**: 686–92.
- 39 Mealer ML, Shelton A, Berg B, Rothbaum B, Moss M. Increased prevalence of post traumatic stress disorder symptoms in critical care nurses. *Am J Respir Crit Care Med* 2007; **175**: 693–97.
- 40 Azoulay E, Timsit JF, Sprung CL, et al. Prevalence and factors of intensive care unit conflicts: the conflictus study. *Am J Respir Crit Care Med* 2009; **180**: 853–60.
- 41 Studdert DM, Mello MM, Burns JP, et al. Conflict in the care of patients with prolonged stay in the ICU: types, sources, and predictors. *Intensive Care Med* 2003; **29**: 1489–97.
- 42 Carlet J, Thijs LG, Antonelli M, et al. Challenges in end-of-life care in the ICU. Statement of the 5th International Consensus Conference in Critical Care: Brussels, Belgium, April 2003. *Intensive Care Med* 2004; **30**: 770–84.
- 43 Truog RD, Campbell ML, Curtis JR, et al. Recommendations for end-of-life care in the intensive care unit: A consensus statement by the American Academy of Critical Care Medicine. *Crit Care Med* 2008; **36**: 953–63.
- 44 Heyland DK, Rocker GM, O'Callaghan CJ, Dodek PM, Cook DJ. Dying in the ICU: perspectives of family members. *Chest* 2003; **124**: 392–97.
- 45 Evans LR, Boyd EA, Malvar G, et al. Surrogate decision-makers' perspectives on discussing prognosis in the face of uncertainty. *Am J Respir Crit Care Med* 2009; **179**: 48–53.
- 46 Heyland DK, Tranmer J, O'Callaghan CJ, Gafni A. The seriously ill hospitalized patient: preferred role in end-of-life decision making? *J Crit Care* 2003; **18**: 3–10.
- 47 Azoulay E, Pochard F, Chevret S, et al. Half the family members of intensive care unit patients do not want to share in the decision-making process: a study in 78 French intensive care units. *Crit Care Med* 2004; **32**: 1832–38.
- 48 White DB, Evans LR, Bautista CA, Luce JM, Lo B. Are physicians' recommendations to limit life support beneficial or burdensome? Bringing empirical data to the debate. *Am J Respir Crit Care Med* 2009; **180**: 320–25.
- 49 Moselli NM, Debernardi F, Piovano F. Forgoing life sustaining treatments: differences and similarities between North America and Europe. *Acta Anaesthesiol Scand* 2006; **50**: 1177–86.
- 50 Mani RK, Mandal AK, Bal S, et al. End-of-life decisions in an Indian intensive care unit. *Intensive Care Med* 2009; **35**: 1713–19.

- 51 Buckley TA, Joynt GM, Tan PY, Cheng CA, Yap FH. Limitation of life support: frequency and practice in a Hong Kong intensive care unit. *Crit Care Med* 2004; **32**: 415–20.
- 52 Giannini A, Pessina A, Tacchi EM. End-of-life decisions in intensive care units: attitudes of physicians in an Italian urban setting. *Intensive Care Med* 2003; **29**: 1902–10.
- 53 White DB, Malvar G, Karr J, Lo B, Curtis JR. Expanding the paradigm of the physician's role in surrogate decision-making: an empirically derived framework. *Crit Care Med* 2010; **38**: 743–50.
- 54 Azoulay E, Chevret S, Leleu G, et al. Half the families of intensive care unit patients experience inadequate communication with physicians. *Crit Care Med* 2000; **28**: 3044–49.
- 55 Azoulay E, Pochard F, Kentish-Barnes N, et al. Risk of post-traumatic stress symptoms in family members of intensive care unit patients. *Am J Respir Crit Care Med* 2005; **171**: 987–94.
- 56 Wall RJ, Curtis JR, Cooke CR, Engelberg RA. Family satisfaction in the ICU: differences between families of survivors and nonsurvivors. *Chest* 2007; **132**: 1425–33.
- 57 Pochard F, Azoulay E, Chevret S, et al, for the French FAMIREA group. Symptoms of anxiety and depression in family members of intensive care unit patients: Ethical hypothesis regarding decision-making capacity. *Crit Care Med* 2001; **29**: 1893–97.
- 58 McDonagh JR, Elliott TB, Engelberg RA, et al. Family satisfaction with family conferences about end-of-life care in the intensive care unit: increased proportion of family speech is associated with increased satisfaction. *Crit Care Med* 2004; **32**: 1484–88.
- 59 Selph RB, Shiang J, Engelberg R, Curtis JR, White DB. Empathy and life support decisions in intensive care units. *J Gen Intern Med* 2008; **23**: 1311–17.
- 60 Curtis JR, Engelberg RA, Wenrich MD, Shannon SE, Treece PD, Rubenfeld GD. Missed opportunities during family conferences about end-of-life care in the intensive care unit. *Am J Respir Crit Care Med* 2005; **171**: 844–49.
- 61 West HF, Engelberg RA, Wenrich MD, Curtis JR. Expressions of nonabandonment during the intensive care unit family conference. *J Palliat Med* 2005; **8**: 797–807.
- 62 Stapleton RD, Engelberg RA, Wenrich MD, Goss CH, Curtis JR. Clinician statements and family satisfaction with family conferences in the intensive care unit. *Crit Care Med* 2006; **34**: 1679–85.
- 63 Schneiderman LJ, Gilmer T, Teetzel HD, et al. Effect of ethics consultations on nonbeneficial life-sustaining treatments in the intensive care setting: a randomized controlled trial. *JAMA* 2003; **290**: 1166–72.
- 64 Kagawa-Singer M, Blackhall LJ. Negotiating cross-cultural issues at the end of life: "You got to go where he lives". *JAMA* 2001; **286**: 2993–3001.
- 65 Pham K, Thornton JD, Engelberg RA, Jackson JC, Curtis JR. Alterations during medical interpretation of ICU family conferences that interfere with or enhance communication. *Chest* 2008; **134**: 109–16.
- 66 Thornton JD, Pham K, Engelberg RA, Jackson JC, Curtis JR. Families with limited English proficiency receive less information and support in interpreted intensive care unit family conferences. *Crit Care Med* 2009; **37**: 89–95.
- 67 Norris WM, Wenrich MD, Nielsen EL, Treece PD, Jackson JC, Curtis JR. Communication about end-of-life care between language-discordant patients and clinicians: Insights from medical interpreters. *J Palliat Med* 2005; **8**: 1016–24.
- 68 Wall RJ, Engelberg RA, Gries CJ, Glavan B, Curtis JR. Spiritual care of families in the intensive care unit. *Crit Care Med* 2007; **35**: 1084–90.
- 69 Gries CJ, Curtis JR, Wall RJ, Engelberg RA. Family member satisfaction with end-of-life decision making in the ICU. *Chest* 2008; **133**: 704–12.
- 70 Gajewska K, Schroeder M, De Marre F, Vincent JL. Analysis of terminal events in 109 successive deaths in a Belgian intensive care unit. *Intensive Care Med* 2004; **30**: 1224–27.
- 71 Nolin T, Andersson R. Withdrawal of medical treatment in the ICU. A cohort study of 318 cases during 1994–2000. *Acta Anaesthesiol Scand* 2003; **47**: 501–07.
- 72 Luce JM, Alpers A. Legal aspects of withholding and withdrawing life support from critically ill patients in the United States and providing palliative care to them. *Am J Respir Crit Care Med* 2000; **162**: 2029–32.
- 73 American Thoracic Society. Withholding and withdrawing life-sustaining therapy. *Ann Intern Med* 1991; **115**: 478–85.
- 74 Levin PD, Sprung CL. Withdrawing and withholding life-sustaining therapies are not the same. *Crit Care* 2005; **9**: 230–02.
- 75 Melltorp G, Nilstun T. The difference between withholding and withdrawing life-sustaining treatment. *Intensive Care Med* 1997; **23**: 1264–67.
- 76 Steinberg A, Sprung CL. The dying patient: new Israeli legislation. *Intensive Care Med* 2006; **32**: 1234–37.
- 77 Ravitsky V. Timers on ventilators. *BMJ* 2005; **330**: 415–17.
- 78 Lanken PN, Terry PB, Delisser HM, et al. An official American Thoracic Society clinical policy statement: palliative care for patients with respiratory diseases and critical illnesses. *Am J Respir Crit Care Med* 2008; **177**: 912–27.
- 79 Luce JM, White DB. The pressure to withhold or withdraw life-sustaining therapy from critically ill patients in the United States. *Am J Respir Crit Care Med* 2007; **175**: 1104–08.
- 80 Kompanje EJ, van der Hoven B, Bakker J. Anticipation of distress after discontinuation of mechanical ventilation in the ICU at the end of life. *Intensive Care Med* 2008; **34**: 1593–99.
- 81 Chan JD, Treece PD, Engelberg RA, et al. Association between narcotic and benzodiazepine use after withdrawal of life support and time to death. *Chest* 2004; **126**: 286–93.
- 82 Bakker J, Jansen TC, Lima A, Kompanje EJ. Why opioids and sedatives may prolong life rather than hasten death after ventilator withdrawal in critically ill patients. *Am J Hosp Palliat Care* 2008; **25**: 152–54.
- 83 Campbell ML, Bizek KS, Thill M. Patient responses during rapid terminal weaning from mechanical ventilation: a prospective study. *Crit Care Med* 1999; **27**: 73–77.
- 84 Gerstel E, Engelberg RA, Koepsell T, Curtis JR. Duration of withdrawal of life support in the intensive care unit and association with family satisfaction. *Am J Respir Crit Care Med* 2008; **178**: 798–804.
- 85 Treece PD, Engelberg RA, Crowley L, et al. Evaluation of a standardized order form for the withdrawal of life support in the intensive care unit. *Crit Care Med* 2004; **32**: 1141–48.