

This Provisional PDF corresponds to the article as it appeared upon acceptance. Copyedited and fully formatted PDF and full text (HTML) versions will be made available soon.

Death after percutaneous dilatational tracheostomy: a systematic review and analysis of risk factors

Critical Care 2013, **17**:R258 doi:10.1186/cc13085

Marcel Simon (m.simon@uke.de)
Maria Metschke (m.metschke@uke.de)
Stephan A Braune (sbraune@uke.de)
Klaus Püschel (pueschel@uke.de)
Stefan Kluge (s.kluge@uke.de)

ISSN 1364-8535

Article type Research

Submission date 8 May 2013

Acceptance date 7 October 2013

Publication date 29 October 2013

Article URL <http://ccforum.com/content/17/5/R258>

This peer-reviewed article can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Critical Care* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Critical Care* go to

<http://ccforum.com/authors/instructions/>

Death after percutaneous dilatational tracheostomy: a systematic review and analysis of risk factors

Marcel Simon^{1,†}
Email: m.simon@uke.de

Maria Metschke^{1,†}
Email: m.metschke@uke.de

Stephan A Braune¹
Email: sbraune@uke.de

Klaus Püschel²
Email: pueschel@uke.de

Stefan Kluge^{1*}
* Corresponding author
Email: s.kluge@uke.de

¹ Department of Intensive Care Medicine, University Medical Center Hamburg-Eppendorf, Martinistr 52, 20246 Hamburg, Germany

² Department of Forensic Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[†] Equal contributors.

Abstract

Introduction

Since the introduction and widespread acceptance of percutaneous techniques in the intensive care unit (ICU) setting, the number of critically ill patients undergoing tracheostomy has steadily increased. However, this procedure can be associated with major complications, including death. The purpose of this study is to estimate the incidence and analyze the causes of lethal complications due to percutaneous dilatational tracheostomy (PDT).

Methods

We analyzed cases of lethal outcome due to complications from PDT including cases published between 1985 and April 2013. A systematic literature search was performed and unpublished cases from our own departmental records were retrospectively analyzed.

Results

71 cases of lethal outcome following PDT were identified including 68 published cases and 3 of our own patients. The incidence of lethal complications was calculated to be 0.17%. Of the

fatal complications, 31.0% occurred during the procedure and 49.3% within seven days of the procedure. The main causes of death were: hemorrhage (38.0%), airway complications (29.6%), tracheal perforation (15.5%), and pneumothorax (5.6%). We found specific risk factors for complications in 73.2% of patients, 25.4% of patients had more than one risk factor. Bronchoscopic guidance was used in only 46.5% of cases.

Conclusions

According to this analysis, PDT-related death occurs in 1 out of 600 patients receiving a PDT. Careful patient selection, bronchoscopic guidance, and securing the tracheal cannula with sutures are likely to reduce complication rates.

Introduction

Since the introduction and widespread acceptance of percutaneous techniques in the intensive care unit (ICU) setting, the number of critically ill patients undergoing tracheostomy has increased in recent years [1,2]. Given a predicted increase in numbers of mechanically ventilated patients, a further increase in the number of tracheostomy procedures in the ICU is to be expected [1-4].

The ideal timing of tracheostomy is still a subject of debate as there is no clear evidence that early tracheostomy improves relevant endpoints, such as duration of mechanical ventilation, length of ICU stay, and mortality [5]. Nevertheless, tracheostomy is being undertaken significantly earlier during ICU stay, as the intervention appears to be beneficial in terms of patient comfort, mobility, and reducing the requirement for sedation [1,6]. Percutaneous dilatational tracheostomy (PDT) has gained wide acceptance and has become the procedure of choice for tracheostomy in critically ill patients worldwide [7]. However this procedure, just like surgical tracheostomy, is associated with major complications, including death. It is estimated that each year approximately 500 patients in the United States die or are permanently disabled because of a tracheostomy [8].

Three fatalities due to PDT in our department in recent years prompted us to perform this study. We aimed to analyze cases from the literature and from our own database to determine the incidence of lethal complications due to percutaneous tracheostomy, to reveal the causes of death, to identify risk factors and possible mechanisms for lethal complications, and finally to develop recommendations to further minimize the risks of complication.

Materials and methods

The study was conducted and reported according to PRISMA guidelines [9].

Search strategy

A systematic search for articles was performed in PubMed, Embase and the Cochrane Library without restrictions for language searching for case reports, case series, observational studies and randomized trials describing or including fatalities associated with PDT. We focused on studies published from 1985 onwards as the percutaneous approach gained popularity after its description by Ciaglia et al. [10] in that specific year.

An extensive and sensitive search strategy was chosen using the keywords “percutaneous tracheostomy” and “percutaneous tracheotomy” to ensure comprehensive retrieval of articles. The search was last updated on 30th April 2013.

Study selection

Studies were selected independently by two authors (MS and SK). Disagreements between the two authors were resolved by discussion. The titles and abstracts of all publications retrieved by the search strategy were screened for eligibility. Studies on PDT were selected for further evaluation. Full-text review of these studies was performed. Studies were included if one or more PDT-related deaths were described. In addition, the reference sections of all studies on PDT were handsearched for additional relevant publications.

Data extraction, assessment, completion and synthesis

Data was independently extracted with standardized forms and interpreted by three authors (MS, SB, SK). The results of data extraction were then compared and disagreement resolved by discussion. Included publications were reviewed manually for relevant data. Information concerning patient characteristics (age, gender, main diagnosis, and duration of mechanical ventilation before tracheostomy), tracheostomy procedure (technique, bronchoscopic guidance, pre-interventional ultrasound, performing physician or team, procedure-related difficulties, special circumstances, and risk factors for complications), cause and time of death as well as departmental characteristics (type of ICU, year of introduction of PDT in the department, number of PDTs performed annually and the number of fatalities related to PDT since its introduction in the department) was extracted. If the published data set was incomplete, we contacted the corresponding author via email or post. If the corresponding author did not respond within 4 weeks, we sent a reminder.

Retrieval of PDT-related fatalities from our own department

To identify tracheostomy-related fatalities in our own department, all patients who had undergone PDT between 1st January 2005 and 31st December 2012 were identified from the departmental electronic patient database. Cases retrieved from this search were further evaluated for procedure-related deaths by manual analysis of their medical records.

Since 1st January 2005, the Ciaglia blue rhino® technique assisted by bronchoscopic guidance has been the standard technique for PDT in our department. The PDTs were performed at the bedside in one of the 10 departmental ICUs according to a standardized operating procedure [11]. After completion, a protocol documenting key aspects of the procedure was generated and saved in the electronic medical record.

The ethics committee of the Hamburg chamber of physicians and the institutional data protection official approved the collection, analysis, and publication of the retrospectively obtained and anonymized data for this non-interventional study.

Data analysis

Results are presented as medians and ranges or as absolute numbers with percentages. The software used for descriptive analyses was Microsoft Excel 2011.

Results

Study and case selection

The search strategy yielded a total of 1963 articles. From these publications 45 studies describing 65 PDT-related fatalities were included in the study. 3 more cases could be added from personal communication with the corresponding authors. 3 additional cases were included from our own departmental records. Overall, 71 cases of PDT-related death were included and analyzed in this study. The process of case selection is summarized in Figure 1.

Figure 1 Process of case selection.* [12-14]. [15,16].**

By contacting the corresponding authors, case specific data could be added in 20 cases and departmental data concerning the total numbers of PDT procedures and PDT-related fatalities could be obtained from 17 departments.

Study characteristics

We found 16 case reports or case series, 7 retrospective studies, 20 prospective observational studies and 2 randomized trials. Study characteristics are summarized in Table 1.

Table 1 Characteristics of studies included

Study					Departmental data obtained by personal communication with the corresponding author				
Author	Year	Design	Population	Number of PDTs	Number of deaths	Mortality rate	Number of PDTs	Number of deaths	Mortality rate
Byhahn	[16]	CR			2				
Cobean	[15]	CR			1				
Hutchinson	[17]	1991 CR			1				
Cokis	[18]	2000 CR			1				
Hürter	[19]	2000 CR	Surgical		1		252	2	0.79
Drage	[20]	2002 CR			1				
Maeda	[21]	2002 CR	Mixed		1				
Soubirou	[22]	2002 CR	Surgical		1				
Ryan	[23]	2003 CR			1				
Shlugman	[24]	2003 CR	Neurological		1		700	1	0.14
McCormick	[25]	2005 CR	Mixed		3		2100	4	0.19
Grant	[26]	2006 CR	Mixed		3				
Rosolski	[27]	2006 CR	Mixed		1		420	1	0.24
Ayoub	[28]	2007 CR			1				
Zehlicke	[29]	2007 CR			1				
Wang	[30]	2009 CR			1				
Hoiting	[31]	2010 CR			2				
Gilbey	[32]	2012 CR	Mixed		1		420	1	0.24
Ivatury	[33]	1992 RS		61	1	1.64			
Cobean	[34]	1996 RS	Mixed	65	1	1.54	1080	2	0.19
van Heurn	[35]	1996 RS	Mixed	150	1	0.67	880	1	0.11
Thompson	[36]	2001 RS	Medical	300	1	0.33			
Pandit	[37]	2006 RS	Mixed	501	1	0.20	1265	2	0.16
Klein	[38]	2007 RS	Mixed	207	1	0.48			
Dennis	[39]	2013 RS		3162	5	0.16	3162	5	0.16
Toye	[40]	1986 POS		94	1	1.06			
Marelli	[41]	1990 POS		61	1	1.64			
Wang	[42]	1992 POS	Surgical	7	1	14.29			
Friedman	[43]	1993 POS		100	1	1.00			
Cole	[44]	1994 POS		55	1	1.82			
Barba	[45]	1995 POS	Surgical	27	1	3.70			
Muhl	[46]	1995 POS	Surgical	14	1	7.14	720	1	0.14
Joosten	[47]	1996 POS	Surgical	53	1	1.89	420	1	0.24
Marx	[48]	1996 POS		254	1	0.39			
Berrouschot	[49]	1997 POS	Neurological	76	2	2.63	900	2	0.22
Walz	[50]	1998 POS		337	2	0.59			
Suh	[51]	1999 POS		95	2	2.11	1040	2	0.19
Escarment	[52]	2000 POS		162	2	1.23			
Kearney	[53]	2000 POS	Mixed	827	5	0.60	5400	9	0.17
Lim	[54]	2000 POS		261	2	0.77			
Norwood	[55]	2000 POS		422	1	0.24			
Tan	[56]	2004 POS		352	1	0.28			
Byhahn	[57]	2005 POS	Surgical	474	2	0.42	2775	4	0.14
Chiu	[58]	2005 POS	Medical	107	1	0.93	900	1	0.11
Páez	[59]	2005 POS		38	2	5.26			
Porter	[60]	1999 RT	Surgical	12	1	8.33			
Massick	[61]	2001 RT	Medical	50	1	2.00			
Own cases			Mixed		3		1873	3	0.16
Summary				8324	71	2.18*	24307	42	0.17

Abbreviations: *CR* case report, *RS* retrospective study, *POS* prospective observational study, *RT* randomized trial

*Calculated mean mortality rate including data from retrospective studies, prospective observational studies and randomized trials.

Summary and analysis of all cases

The median age was 66 years (range 4 to 95). For further details concerning patient characteristics see Table 2. Most tracheostomies were performed using the Ciaglia (45.1%) and the Ciaglia blue rhino (26.8%) technique. Bronchoscopic guidance was used in 46.5% of cases, but no cases involved pre- or peri-interventional ultrasound.

Table 2 Patient characteristics

Characteristic	Values	
Number of patients	71	
Median age [years]	66	(range 4–95)
Gender		
female	33	(46.5%)
male	16	(22.5%)
not specified	22	(31.0%)
Main diagnosis		
pulmonary disease	15	(21.1%)
neurologic disease	18	(25.4%)
trauma	9	(12.7%)
cardiac disease	5	(7.0%)
surgical procedure	4	(5.6%)
vascular disease	3	(4.2%)
not specified	17	(23.9%)
Type of intensive care unit		
interdisciplinary	13	(18.3%)
surgical	11	(15.5%)
medical	7	(9.9%)
neurological	6	(8.5%)
cardio-thoracic	1	(1.4%)
trauma	3	(4.2%)
not specified	30	(42.3%)
Median duration of intubation before tracheostomy [days]	11	(range: 0–33)

The major causes of death were tracheostomy-related hemorrhage in 27 patients (38.0%) and airway complications in 21 patients (29.6%). In 31.0% of cases, fatal complications occurred during the procedure and in 49.3% of cases within seven days of the procedure. 73.2% of patients had specific risk factors and 25.4% of patients had more than one risk factor. For further details about the causes of death and time of complications see Table 3.

Table 3 Causes and time of death after PDT

Cause of death	Total number		Time of death			
			intra-procedural		post-procedural	
Total number	71		22	(31.0%)	49	(69.0%)
Hemorrhage	27	(38.0%)	3	(11.1%)	24	(88.9%)
- innominate artery	11	(40.7%)			11	(100.0%)
- aortic arch	2	(7.4%)			2	(100.0%)
- subclavian artery	1	(3.7%)	1	(100.0%)		
- thyroid artery	1	(3.7%)			1	(100.0%)
- other artery	1	(3.7%)	1	(100.0%)		
- venous	5	(18.5%)	1	(20.0%)	4	(80.0%)
- diffuse/unknown	6	(22.2%)			6	(100.0%)
Airway complications	21	(29.6%)	7	(33.3%)	14	(66.7%)
- dislocation of the tracheal cannula	11	(52.4%)	1	(9.1%)	10	(90.9%)
- lost airway during the procedure	4	(19.0%)	3	(75.0%)	1	(25.0%)
- paratracheal misplacement of the tracheal cannula	3	(14.3%)	3	(100.0%)		
- obstruction of tracheal cannula	2	(9.5%)			2	(100.0%)
- hypoxemia during cannula replacement	1	(4.8%)			1	(100.0%)
Tracheal perforation	11	(15.5%)	1	(9.1%)	10	(90.9%)
Pneumothorax	4	(5.6%)	4	(100.0%)		
Bronchospasm	3	(4.2%)	3	(100.0%)		
Cardiac arrest / arrhythmia	3	(4.2%)	3	(100.0%)		
Sepsis	1	(1.4%)			1	(100.0%)
Unknown	1	(1.4%)	1	(100.0%)		

Hemorrhage

We found 27 deaths (38.0%) due to hemorrhage [18,19,22-26,28-31,35,37,39,43,46,49,54,57-59]. The source of bleeding was arterial in 16 cases (59.3%). Most often arterial bleeding originated from tracheo-vascular fistula formation involving the innominate artery (11 cases). The majority of bleeding incidents (75.0%) occurred between 1 day and 1 month after the tracheostomy procedure, with a median of 5 days. The following known risk factors were retrospectively attributed as potentially relevant in these fatalities secondary to hemorrhagic complications: not using bronchoscopic guidance (8 cases), low tracheostomy site (5 cases), coagulopathy (5 cases), previous surgery to the neck (4 cases), previous radiotherapy (1 case), obesity (1 case), anatomical abnormality (1 case), paratracheal misplacement of the tracheal cannula (1 case), malpositioned cannula tip (1 case), and high cuff pressure (1 case). 21 patients (77.8%) had at least one of the described risk factors.

Airway complications

21 deaths (29.6%) due to airway complications were reported [32,34,38-40,42,44,48,50-53,56,60,61]. Main causes of death in this group were dislocation of the tracheal cannula (52.4%), lost airway during the procedure (19.0%) and paratracheal misplacement of the tracheal cannula (14.3%). Performing the procedure without bronchoscopic guidance (6 cases), by a team relatively inexperienced with the procedure (5 cases), in obese patients (8 cases), in patients with a difficult airway (2 cases), not securing the tracheal cannula with sutures (3 cases), early cannula replacement (1 case), and post-procedural care by an inexperienced team (1 case) were determined as risk factors for these fatalities attributed to airway complications. 19 patients (90.5%) had at least one of the described risk factors.

Tracheal perforation

Including our own cases, a total of 11 deaths (15.5%) were due to tracheal perforation [17,20,47,49,53,57,59]. Performing the procedure without using bronchoscopy (5 cases), by an inexperienced team (4 cases), in an obese patient (1 case), in a child (1 case), noticing a kinking of the guidewire post-procedurally (2 cases), and anatomic abnormalities of the spine (1 case) were determined as risk factors for this kind of complication. 9 patients (81.8%) had at least one of the described risk factors.

Pneumothorax

Four deaths (5.6%) resulting from a pneumothorax were reported [15,27,36,55]. Two deaths were due to tension pneumothoraces and two were due to bilateral pneumothorax. Primary symptoms were subcutaneous emphysema and/or hypoxemia. Chronic obstructive pulmonary disease (3 cases) was stated as underlying condition putting patients at risk for this type of complication.

Bronchospasm

Three deaths (4.4%) due to bronchospasm were reported [33,45,53]. Severe acute respiratory distress syndrome (ARDS) (2 cases) and chronic obstructive pulmonary disease (1 case) were stated as underlying conditions putting patients at risk for this type of complication.

Cardiac arrest/arrhythmia

Three intraoperative deaths (4.4%) due to cardiac arrest were reported [41,50,57]. Cardiac disease and surgery to the heart were recognized as underlying conditions putting patients at risk for this type of complication.

Sepsis

One death (1.5%) was due to sepsis [21]. Sepsis was due to mediastinitis originating from the tracheostomy site. Previous sternotomy performed for the treatment of an aneurysm of the thoracic aorta was identified as a risk factor.

Mortality rate

The mean mortality rate calculated from retrospective studies, prospective observational studies and randomized trials included in this study and overlooking 8,324 PDT procedures was 2.18%.

Departmental data could be obtained from 17 departments responsible for a total number of 24,307 PDT procedures and 42 fatalities. The incidence of death attributable to PDT calculated from these numbers was 0.17%.

In our own department 1873 patients underwent PDT between 1st January 2005 and 31st December 2012 and 3 procedure-related fatalities occurred resulting in an incidence of PDT-related lethal complications of 0.16%.

Discussion

We analyzed 71 cases of death due to PDT. The incidence of death related to PDT calculated from departmental data provided by corresponding authors was 0.17%, which is in line with the lethal complication rate in our own institution. The main reasons for the catastrophic events resulting in death were vascular injuries and airway complications. We found specific risk factors in 73.2% of patients, and 25.4% of patients had more than one risk factor. To our knowledge this is the first systematic analysis of lethal complications due to PDT.

Gilbey et al. recently published a case series of fatal cases due to PDT and concluded that this event usually results from vascular injury [32]. However, their report included only 7 cases, without providing further information about demographics, PDT technique, use of bronchoscopy or ultrasound.

Even after surgical tracheostomy, fatal complications can occur. In a survey of members of the American Academy of Otolaryngology, Head and Neck Surgery two thirds of (mainly surgical) tracheostomy-related catastrophic events were reported to be mainly due to loss of airway or bleeding [8].

Tracheostomy-related hemorrhage was the most common cause of death in our study. Massive hemorrhage is a rare but devastating complication after any form of tracheostomy and usually originates from tracheo-arterial fistula formation. The majority of cases occur within 3 days to 6 weeks of tracheostomy, and risk factors include pressure necrosis from high cuff pressure, mucosal trauma, malpositioned cannula tip, low tracheal incision, excessive neck movement, radiotherapy, or prolonged intubation [26]. We found such risk factors in 77.8% of these patients. In 29.6% of studied patients with fatal bleeding, performing the procedure without bronchoscopy was determined as a risk factor. In this group, placement of tracheostomy was too low in 5 patients, of which 2 had an aortic arch laceration. Coagulation dysfunction or platelet dysfunction were present in 5 patients. These were not deemed to be the primary cause of bleeding in 2 cases as these patients died from acute and sudden bleeding from specific blood vessels and without diffuse bleeding. Indeed, it has been shown previously, that even in severe thrombocytopenia, PDT can be safely performed after preprocedural correction [62].

A major risk factor seems to be a low tracheal incision, as was the case in 5 of 27 patients with bleeding complications in our study. The site of puncture should ideally be selected between first and second or second and third tracheal rings [63]. In one study, the site of tracheal puncture was changed in 24% of patients as a result of prior ultrasound [64]. Furthermore preprocedural ultrasound and clinical examination have been used to detect abnormal pretracheal vascular anatomy. Therefore several authors recommend ultrasound to improve the safety of PDT [64-67].

The second most frequent cause of death was airway complications. The tracheal cannula was placed outside the tracheal lumen in 3 cases. Of note, in all of these cases bronchoscopic guidance was not used. Despite the lack of randomized controlled trials many authors think that the use of bronchoscopic guidance significantly increases the safety of PDT as it can guide correct placement of the introducer needle, the guidewire, and the cannula during the procedure [68,69].

In a further 11 patients, the tracheal cannula accidentally dislocated postprocedurally. While the surgical approach allows easy re-insertion of the tracheal cannula, airway complications such as accidental decannulation or tube obstruction are well described problems of the percutaneous technique [70]. Some researchers have proposed that fixing the tracheal cannula to the skin with sutures for the first postprocedural week may decrease cannula-related complications such as accidental decannulation and postoperative bleeding [71,72].

In 2 of the 11 deaths (18.2%) due to posterior tracheal wall perforation, kinking of the guidewire was noticed after its removal during the procedure. Other researchers have postulated guidewire kinking as a possible mechanism for perforation of the posterior tracheal wall [73,74]. Thus, kinking of the guidewire must be avoided and its occurrence should raise suspicion of potential injury to the posterior tracheal wall prompting further investigation.

To avoid complications, PDT should only be considered in selected patients without contraindications. Contraindications to PDT include anatomic distortion of the neck, the presence of a difficult airway, severe ARDS, uncorrectable coagulopathy, and the presence of an unstable cervical spine [68]. However, most of these contraindications are relative and also dependent on the skill of the operator. Of the 10 cases (14.1%) in our study, where the performing team was deemed to be relatively inexperienced in the procedure, contraindications were also present in 2 patients (20.0%).

Because of the growing numbers of patients requiring ventilatory support, the frequency of tracheostomy in the ICU has increased over the last decades as has the number of PDT-related publications (Additional file 1). In addition, it is conceivable that after the introduction and widespread acceptance of percutaneous techniques, enthusiasm about the ease of performing this procedure at the bedside may have resulted in relative overuse by intensivists.

A general limitation of this systematic review is the possibility of under- or over-estimating the true incidence of PDT-related death. Accurate risk assessment in clinical medicine is most difficult when an event is rare and available evidence is based on self-reported data [75]. Additionally, concerning the topic of this study, there is sometimes considerable difficulty in differentiating whether an adverse event is due to being critically ill with a tracheostomy or whether it is a complication of the PDT procedure itself. However, in all cases included in this study, the authors of the original publications judged death to be most likely related to the PDT procedure itself. Calculated from departmental data provided by the corresponding authors - where we are confident of both the completeness of reporting and the denominator - the mortality rate from PDT was 0.17%. This is very much in line with the mortality rate of 0.16% reported in the largest study published to date [39] and the mortality rate of 0.16% in our own institution.

In our opinion, this first systemic analysis of the incidence and causes of PDT-related lethal complications adds important evidence to the literature and serves to remind clinicians about potentially life-threatening complications and to help them minimize risk factors by choosing suitable patients and safe procedural strategies.

Conclusions

In conclusion, PDTs, which are frequently performed in ICUs worldwide, are associated with an average mortality of one in every 600 procedures. Major risk factors are present in a substantial proportion of these patients. To prevent severe complications, the results of this systematic review, several authors and professional guidelines suggest that the following measures may improve the safety of PDT: strict consideration of contraindications, bronchoscopic guidance during the entire procedure, performance by an experienced team, avoidance of a low tracheostomy puncture site and avoidance of guidewire kinking as well as the use of outer flange tracheal cannula sutures.

Key messages

- PDT-related death occurs in 1 out of 600 procedures.
- Careful patient selection, bronchoscopic guidance, and securing the tracheal cannula with sutures are likely to reduce complication rates.

Abbreviations

ARDS, Acute respiratory distress syndrome; ICU, Intensive care unit; PDT, Percutaneous dilatational tracheostomy

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MS, MM and SK have made substantial contributions to conception and design of the study as well as to the acquisition, analysis and interpretation of data. SB and KP have made substantial contributions to analysis and interpretation of data. MS, MM and SK have drafted the submitted article. SB and KP have revised it critically for important intellectual content. All authors read and approved the final manuscript.

References

1. Cox CE, Carson SS, Holmes GM, Howard A, Carey TS: **Increase in tracheostomy for prolonged mechanical ventilation in North Carolina, 1993–2002.** *Crit Care Med* 2004, **32**:2219–2226.
2. Simpson TP, Day CJ, Jewkes CF, Manara AR: **The impact of percutaneous tracheostomy on intensive care unit practice and training.** *Anaesthesia* 1999, **54**:186–189.
3. Needham DM, Bronskill SE, Calinawan JR, Sibbald WJ, Pronovost PJ, Laupacis A: **Projected incidence of mechanical ventilation in Ontario to 2026: Preparing for the aging baby boomers.** *Crit Care Med* 2005, **33**:574–579.

4. Zilberberg MD, de Wit M, Pirone JR, Shorr AF: **Growth in adult prolonged acute mechanical ventilation: implications for healthcare delivery.** *Crit Care Med* 2008, **36**:1451–1455.
5. Wang F, Wu Y, Bo L, Lou J, Zhu J, Chen F, Li J, Deng X: **The timing of tracheotomy in critically ill patients undergoing mechanical ventilation: a systematic review and meta-analysis of randomized controlled trials.** *Chest* 2011, **140**:1456–1465.
6. Kluge S, Baumann HJ, Maier C, Klose H, Meyer A, Nierhaus A, Kreymann G: **Tracheostomy in the intensive care unit: a nationwide survey.** *Anesth Analg* 2008, **107**:1639–1643.
7. Freeman BD, Morris PE: **Tracheostomy practice in adults with acute respiratory failure.** *Crit Care Med* 2012, **40**:2890–2896.
8. Das P, Zhu H, Shah RK, Roberson DW, Berry J, Skinner ML: **Tracheotomy-related catastrophic events: results of a national survey.** *Laryngoscope* 2012, **122**:30–37.
9. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, Clarke M, Devereaux PJ, Kleijnen J, Moher D: **The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration.** *In PLoS Med* 2009, **6**:e1000100.
10. Ciaglia P, Firsching R, Syniec C: **Elective percutaneous dilatational tracheostomy. A new simple bedside procedure; preliminary report.** *Chest* 1985, **87**:715–719.
11. Braune S, Kluge S: **Die perkutane Dilatationstracheotomie.** *Dtsch Med Wochenschr* 2011, **136**:1265–1269.
12. Dempsey GA, Grant CA, Jones TM: **Percutaneous tracheostomy: a 6 year prospective evaluation of the single tapered dilator technique.** *Br J Anaesth* 2010, **105**:782–788.
13. Hill BB, Zweng TN, Maley RH, Charash WE, Toursarkissian B, Kearney PA: **Percutaneous dilational tracheostomy: report of 356 cases.** *J Trauma* 1996, **41**:238–243.
14. Toursarkissian B, Zweng TN, Kearney PA, Pofahl WE, Johnson SB, Barker DE: **Percutaneous dilational tracheostomy: report of 141 cases.** *Ann Thorac Surg* 1994, **57**:862–867.
15. Cobean R: *Personal communication*
16. Byhahn C: *Personal communication*
17. Hutchinson RC, Mitchell RD: **Life-threatening complications from percutaneous dilational tracheostomy.** *Crit Care Med* 1991, **19**:118–120.
18. Cokis C, Towler S: **Tracheo-innominate fistula after initial percutaneous tracheostomy.** *Anaesth Intensive Care* 2000, **28**:566–569.

19. Hürter H, Post-Stanke A: **Tödliche Gefäßarrosion nach Dilatationstracheotomie.** *Anesthesiol Intensivmed Notfallmed Schmerzther* 2000, **35**:658–660.
20. Drage SM, Pac Soo C, Dexter T: **Delayed presentation of tracheo-oesophageal fistula following percutaneous dilatational tracheostomy.** *Anaesthesia* 2002, **57**:932–933.
21. Maeda K, Ninomiya M, Moyairi T, Morota T, Kitamura R, Takamoto S: **Mediastinitis after percutaneous dilatational tracheostomy.** *Thorac Cardiovasc Surg* 2002, **50**:123–124.
22. Soubirou JL, Puidupin A, Augeul G, Leclerc T, Combourieu E, Patrigeon RG, Escarment J: **Complication sévère après une trachéotomie translaryngée.** *Ann Fr Anesth Reanim* 2002, **21**:728–730.
23. Ryan DW, Kilner AJ: **Another death after percutaneous dilational tracheostomy.** *Br J Anaesth* 2003, **91**:925–926.
24. Shlugman D, Satya-Krishna R, Loh L: **Acute fatal haemorrhage during percutaneous dilatational tracheostomy.** *Br J Anaesth* 2003, **90**:517–520.
25. McCormick B, Manara AR: **Mortality from percutaneous dilatational tracheostomy. A report of three cases.** *Anaesthesia* 2005, **60**:490–495.
26. Grant CA, Dempsey G, Harrison J, Jones T: **Tracheo-innominate artery fistula after percutaneous tracheostomy: three case reports and a clinical review.** *Br J Anaesth* 2006, **96**:127–131.
27. Rosolski T: **Spannungspneumothorax Nach Punktionstracheotomie.** *J Anaesth Intensivbehandlung* 2006, **3**:36.
28. Ayoub OM, Griffiths MV: **Aortic arch laceration: A lethal complication after percutaneous tracheostomy.** *Laryngoscope* 2007, **117**:176–178.
29. Zehlicke T, Dommerich S, Rummel J: **Letale Blutung aus dem Truncus brachiocephalicus - Spätkomplikation nach Fehleinschätzung eines Dilatationstracheostomas bei rudimentärer Halsrippe.** *Laryngorhinootologie* 2007, **86**:655–659.
30. Wang P-K, Yen P-S, Shyr M-H, Chen T-Y, Chen A, Liu H-T: **Endovascular repair of tracheo-innominate artery fistula.** *Acta Anaesthesiol Taiwan* 2009, **47**:36–39.
31. Hoiting O, van den Brule J, van Zwam PH: **Late fatal bleeding after percutaneous dilatational tracheostomy.** *Neth J Crit Care* 2010, **14**:335–337.
32. Gilbey P: **Fatal complications of percutaneous dilatational tracheostomy.** *Am J Otolaryngol* 2012, **32**:770–773.
33. Ivatury R, Siegel JH, Stahl WM, Simon R, Scorpio R, Gens DR: **Percutaneous tracheostomy after trauma and critical illness.** *J Trauma* 1992, **32**:133–140.

34. Cobean R, Beals M, Moss C: **Percutaneous dilatational tracheostomy: a safe, cost-effective bedside procedure.** *Arch Surg* 1996, **131**:265–271.
35. van Heurn LW, van Geffen GJ, Brink PR: **Clinical experience with percutaneous dilatational tracheostomy: report of 150 cases.** *Eur J Surg* 1996, **162**:531–535.
36. Thompson EC, Fernandez LG, Norwood S, Wilkins H, Vallina VL: **Percutaneous dilatational tracheostomy in a community hospital setting.** *South Med J* 2001, **94**:208–211.
37. Pandit RA, Jacques TC: **Audit of over 500 percutaneous dilational tracheostomies.** *Crit Care Resusc* 2006, **8**:146–150.
38. Klein M, Agassi R, Shapira A-R, Kaplan DM, Koiffman L, Weksler N: **Can intensive care physicians safely perform percutaneous dilational tracheostomy? An analysis of 207 cases.** *Isr Med Assoc J* 2007, **9**:717–719.
39. Dennis BM, Eckert MJ, Gunter OL, Morris JA, May AK: **Safety of bedside percutaneous tracheostomy in the critically ill: evaluation of more than 3,000 procedures.** *J Am Coll Surg* 2013, **216**:858–867.
40. Toye FJ, Weinstein JD: **Clinical experience with percutaneous tracheostomy and cricothyroidotomy in 100 patients.** *J Trauma* 1986, **26**:1034–1040.
41. Marelli D, Paul A, Manolidis S, Walsh G, Odum JN, Burdon TA, Shennib H, Vestweber KH, Fleischer DM, Mulder DS: **Endoscopic guided percutaneous tracheostomy: early results of a consecutive trial.** *J Trauma* 1990, **30**:433–435.
42. Wang MB, Berke GS, Ward PH, Calcaterra TC, Watts D: **Early experience with percutaneous tracheotomy.** *Laryngoscope* 1992, **102**:157–162.
43. Friedman Y, Mayer AD: **Bedside percutaneous tracheostomy in critically ill patients.** *Chest* 1993, **104**:532–535.
44. Cole IE: **Elective percutaneous (Rapitrac) tracheotomy: results of a prospective trial.** *Laryngoscope* 1994, **104**:1271–1275.
45. Barba C, Angood P, Kauder D, Latenser B: **Bronchoscopic guidance makes percutaneous tracheostomy a safe, cost-effective, and easy-to-teach procedure.** *Surgery* 1995, **118**:879–883.
46. Muhl E, Franke C, BRUCH HP: **Verbesserte Technik der Dilatationstracheostomie und erste Ergebnisse.** *Anesthesiol Intensivmed Notfallmed Schmerzther* 1995, **30**:497–500.
47. Joosten U, Sturbeck K, Hohlbach G: **Die Punktionstracheotomie beim langzeitbeatmeten Intensivpatienten.** *Langenbecks Arch Chir Suppl Kongressbd* 1996, **113**:356–359.

48. Marx WH, Ciaglia P, Graniero KD: **Some important details in the technique of percutaneous dilatational tracheostomy via the modified Seldinger technique.** *Chest* 1996, **110**:762–766.
49. Berrouschot J, Oeken J, Schneider D: **Perioperative complications of percutaneous dilatational tracheostomy.** *Laryngoscope* 1997, **107**:1538–1544.
50. Walz MK, Peitgen K, Thürauf N, Trost HA, Wolfhard U, Sander A, Ahmadi C, Eigler FW: **Percutaneous dilatational tracheostomy—early results and long-term outcome of 326 critically ill patients.** *Intensive Care Med* 1998, **24**:685–690.
51. Suh RH, Margulies DR, Hopp ML, Ault M, Shabot MM: **Percutaneous dilatational tracheostomy: still a surgical procedure.** *Am Surg* 1999, **65**:982–986.
52. Escarment J, Suppini A, Sallaberry M, Kaiser E, Cantais E, Palmier B, Quinot JF: **Percutaneous tracheostomy by forceps dilation: report of 162 cases.** *Anaesthesia* 2000, **55**:125–130.
53. Kearney PA, Griffen MM, Ochoa JB, Boulanger BR, Tseui BJ, Mentzer RM: **A single-center 8-year experience with percutaneous dilatational tracheostomy.** *Ann Surg* 2000, **231**:701–709.
54. Lim JW, Friedman M, Tanyeri H, Lazar A, Caldarelli DD: **Experience with percutaneous dilatational tracheostomy.** *Ann Otol Rhinol Laryngol* 2000, **109**:791–796.
55. Norwood S, Vallina VL, Short K, Saigusa M, Fernandez LG, McLarty JW: **Incidence of tracheal stenosis and other late complications after percutaneous tracheostomy.** *Ann Surg* 2000, **232**:233–241.
56. Tan C, Lee H: **Percutaneous dilatational tracheostomy - a 3 year experience in a general hospital in Malaysia.** *Med J Malaysia* 2004, **59**:591–597.
57. Byhahn C, Lischke V, Meininger D, Halbig S, Westphal K: **Peri-operative complications during percutaneous tracheostomy in obese patients.** *Anaesthesia* 2005, **60**:12–15.
58. Chiu C-T, Chung Y-H, Lu H-I, Lin M-C: **Weaning of long-term mechanically-ventilated patients following video bronchoscopy-guided percutaneous dilatational tracheostomy.** *Chang Gung Med J* 2005, **28**:829–836.
59. Páez M, Buisán F, Almaraz A, Martínez-Martínez A, Muñoz F: **Percutaneous tracheotomy with the Ciaglia Blue Rhino technique: a critical analysis after 1 year.** *Rev Esp Anesthesiol Reanim* 2005, **52**:466–473.
60. Porter JM, Ivatury RR: **Preferred route of tracheostomy—percutaneous versus open at the bedside: a randomized, prospective study in the surgical intensive care unit.** *Am Surg* 1999, **65**:142–146.
61. Massick DD, Yao S, Powell DM, Griesen D, Hobgood T, Allen JN, Schuller DE: **Bedside tracheostomy in the intensive care unit: a prospective randomized trial**

comparing open surgical tracheostomy with endoscopically guided percutaneous dilational tracheotomy. *Laryngoscope* 2001, **111**:494–500.

62. Kluge S, Meyer A, Kühnelt P, Baumann HJ, Kreymann G: **Percutaneous tracheostomy is safe in patients with severe thrombocytopenia.** *Chest* 2004, **126**:547–551.

63. ANZICS: *Percutaneous Dilatational Tracheostomy Consensus Statement.* ; 2010.

64. Kollig E, Heydenreich U, Roetman B, Hopf F, Muhr G: **Ultrasound and bronchoscopic controlled percutaneous tracheostomy on trauma ICU.** *Injury* 2000, **31**:663–668.

65. Rajajee V, Fletcher JJ, Rochlen LR, Jacobs TL: **Real-time ultrasound-guided percutaneous dilational tracheostomy: a feasibility study.** *Crit Care* 2011, **15**:R67.

66. Otchwemah R, Defosse J, Wappler F, Sakka SG: **Percutaneous dilatation tracheostomy in the critically ill: use of ultrasound to detect an aberrant course of the brachiocephalic trunk.** *J Cardiothorac Vasc Anesth* 2012, **26**:e72–e73.

67. Rudas M, Seppelt I: **Safety and efficacy of ultrasonography before and during percutaneous dilational tracheostomy in adult patients: a systematic review.** *Crit Care Resusc* 2012, **14**:297–301.

68. Braune S, Kluge S: **Update Tracheotomie.** *Med Klin Intensivmed Notfmed* 2012, **107**:543–547.

69. De Leyn P, Bedert L, Delcroix M, Depuydt P, Lauwers G, Sokolov Y, Van Meerhaeghe A, Van Schil P, Belgian Association of Pneumology and Belgian Association of Cardiothoracic Surgery: **Tracheotomy: clinical review and guidelines.** *Eur J Cardiothorac Surg* 2007:412–421.

70. Higgins KM, Punthakee X: **Meta-analysis comparison of open versus percutaneous tracheostomy.** *Laryngoscope* 2007, **117**:447–454.

71. Halum SL, Ting JY, Plowman EK, Belafsky PC, Harbarger CF, Postma GN, Pitman MJ, LaMonica D, Moscatello A, Khosla S, Cauley CE, Maronian NC, Melki S, Wick C, Sinacori JT, White Z, Younes A, Ekbom DC, Sardesai MG, Merati AL: **A multi-institutional analysis of tracheotomy complications.** *Laryngoscope* 2012, **122**:38–45.

72. Beiderlinden M, Karl Walz M, Sander A, Groeben H, Peters J: **Complications of bronchoscopically guided percutaneous dilational tracheostomy: beyond the learning curve.** *Intensive Care Med* 2002, **28**:59–62.

73. Byhahn C, Wilke HJ, Halbig S, Lischke V, Westphal K: **Percutaneous tracheostomy: ciaglia blue rhino versus the basic ciaglia technique of percutaneous dilational tracheostomy.** *Anesth Analg* 2000, **91**:882–886.

74. Thant M, Samuel T: **Posterior tracheal wall tear with PercuTwist.** *Anaesthesia* 2002, **57**:507–508.

75. Eibling DE, Roberson DW: **Managing tracheotomy risk: time to look beyond hospital discharge.** *Laryngoscope* 2012, **122**:23–24.

Additional file

Additional_file_1 as TIFF

Additional file 1 Number of publications reporting PDT-related fatalities and total number of publications on PDT procedures.




Figure 1

Additional files provided with this submission:

Additional file 1: 9119462499884424_add1.tiff, 6077K
<http://ccforum.com/imedia/1246765791103206/supp1.tiff>