

Revisiting, Reframing, and Casting a New Light on Liberation From Mechanical Ventilation

Timothy D. Girard, MD, MSCI; Karen E. A. Burns, MD, FRCPC, MSc

Every year, more than 1 million patients throughout the world receive mechanical ventilation for acute respiratory failure. One of the most important decisions clinicians make in managing these critically ill patients is how to liberate them from invasive ventilation. Ventilator liberation poses an important dilemma for clinicians because premature or failed attempts at extubation, which typically necessitate reintubation, increase rates of ventilator-associated pneumonia, mortality, and other adverse outcomes.¹ Conversely, delaying extubation also increases a patient's risk of being oversedated and developing delirium or ventilator-associated events.²

The current standard of care for managing patients with invasive ventilation involves daily screenings to identify patients who are ready to undergo a spontaneous breathing trial (SBT) to ascertain those ready to be separated from ventilator support.^{3,4} Additionally, clinicians determine whether patients can be liberated from an endotracheal tube by assessing for extubation failure risk factors (eg, weak cough, heavy secretions, and depressed level of consciousness) before deciding to extubate patients who have had a successful SBT. Before the SBT became the standard assessment of ventilator liberation readiness, a series of investigations searched for the ideal predictor, but the “weaning parameters” examined were limited in their ability to predict successful extubation. Consequently, observing a still-intubated patient during a trial of unassisted breathing (ie, the SBT) remained the preferred approach. In 1995, Esteban et al⁵ published a randomized trial that found that using daily SBTs hastened successful liberation compared with gradually titrating mandatory ventilation rate or inspiratory pressure. Shortly thereafter, Ely et al⁶ showed that a respiratory therapy-driven SBT protocol resulted in earlier extubation than a physician-directed approach.

When conducting an SBT, a number of different techniques can be used. During a T-piece SBT, which was used by Esteban et al,⁵ a patient receives supplemental oxygen but no ventilatory assistance. Alternatively, a patient may receive pressure support ventilation (PSV), during which a small amount of positive pressure (eg, 5-8 cm H₂O) assists inspiration. Additionally, the safety criteria used to determine when an SBT should be started can be either conservative or liberal, and SBTs can be conducted for variable durations. Thus, more than 2 decades after Esteban et al⁵ and Ely et al⁶ published trials ushering SBTs into routine use, new evidence to guide decision making about liberation

from invasive ventilation in current practice is still warranted and needed.

In this issue of the *JAMA*, Subirà and colleagues⁷ present the results of the largest randomized clinical trial ever conducted comparing alternative SBT techniques. The authors randomized 1153 mechanically ventilated adults in 18 Spanish intensive care units (ICUs) to 2 different SBT techniques: (1) a “highly demanding” 2-hour T-piece SBT or (2) a “less demanding” 30-minute PSV SBT with 8-cm H₂O inspiratory pressure and 0-cm H₂O positive end-expiratory pressure. By study design, participants in both groups who successfully completed an SBT were extubated because the trial eligibility criteria ensured that enrolled participants were good candidates for extubation. In this trial, participants in whom the initial SBT was unsuccessful were ventilated at the discretion of the ICU team, and subsequent SBTs were not directed by the trial protocol.

Subirà et al⁷ found that participants in the 30-minute PSV SBT group were more likely than those in the 2-hour T-piece SBT group to have a successful initial SBT and therefore to be extubated. Thus, successful extubation (the primary outcome) occurred in 82.3% of patients in the PSV group and in 74.0% of participants in the T-piece group, an absolute increase of 8.2% (95% CI, 3.4%-13.0%). Additionally, although more participants in the less demanding SBT group had a successful initial SBT and were extubated, they did not experience a higher reintubation rate in the 72 hours after extubation. Moreover, those in the PSV SBT group were also significantly less likely to die in the hospital or during the 90 days after randomization—results that are difficult to explain given that the SBT techniques did not appear to affect other secondary outcomes. Nevertheless, these results are important because they provide evidence that, at least during an initial SBT, most patients can be tested for 30 minutes using 8-cm H₂O PSV.

Less demanding SBT techniques were previously studied in 2 multicenter randomized trials conducted in the late 1990s.^{8,9} Similar to the trial by Subirà et al, these older investigations found no significant differences in reintubation rates when comparing 30-minute SBTs with 120-minute SBTs⁹ and comparing SBTs conducted with 7-cm H₂O PSV vs T-piece SBTs.⁸ Given these results, it is worth considering why many clinicians continue to rely on 120-minute SBTs and many still prefer to conduct T-piece SBTs.⁴ Clinicians' tendency to be risk averse and delays in knowledge transfer may explain some of the variation in ventilator liberation practices, but limitations in the evidence are also

likely a contributing factor. No equivalence trials have been conducted comparing different SBT techniques, and questions remain regarding the quality and generalizability of the evidence. These new results reported by Subirà and colleagues⁷ are therefore an important and welcome addition to the literature. At the same time, their trial raises a number of important questions.

First, when should SBTs be started during a patient's recovery from acute respiratory failure? Similar to previous trials,⁵ a high percentage of participants in the trial by Subirà et al had a successful initial SBT, which leads to the question of whether some participants were ready to undergo an SBT and to be considered for extubation at an earlier time point. That only half of the patients who experience unplanned extubation require reintubation¹⁰ suggests that recognizing the earliest time when patients are ready for extubation is a challenge for ICU clinicians.

Second, should a 30-minute PSV SBT be used regardless of a patient's characteristics and circumstances, or should patient-related factors influence how clinicians conduct SBTs? Even though Subirà et al enrolled a diverse group of participants in multiple centers, their protocol applied only to each participant's initial SBT. Consequently, the results of this trial cannot be generalized to patients in whom an initial SBT is unsuccessful, a group that may be more likely to have extubation failure after a successful 30-minute SBT.¹¹ Future trials of ventilator liberation techniques should continue to apply study interventions until patients achieve successful extubation (or another trial end point).

Third, does the use of related interventions—eg, computerized ventilator weaning¹² or extubation to noninvasive ventilation or high-flow nasal cannula oxygen therapy after a successful (or even a failed) SBT¹³—modify the effects or the implementation of different SBT techniques? The availability of noninvasive modes of support, for example, may encourage clinicians to extubate high-risk patients after a less demanding SBT. It is notable that in the trial by Subirà and colleagues, the use of prophylactic noninvasive ventilation or

high-flow nasal cannula oxygen therapy after extubation was more common in the 30-minute PSV SBT group than in the 2-hour T-piece SBT group (25% vs 19%; $P = .01$).

Fourth, does the SBT technique affect long-term outcomes? If so, how? The effects of interventions on short-term outcomes are important and generally simple to ascertain in clinical trials. Long-term outcomes, however, including long-term survival and functional status, are equally or perhaps even more important to patients.¹⁴ The trial by Subirà and colleagues⁷ suggests that 90-day mortality may be reduced by using 30-minute PSV SBTs rather than more demanding SBTs. If this finding is true, identifying the mechanism(s) underlying this putative benefit will not only promote knowledge transfer but may also help in the identification of other interventions that favorably influence long-term outcomes.

Fifth, how should outcomes be defined and reported in ventilator liberation trials? Although successful extubation is an outcome valued by clinicians, patients, and patients' families,¹⁴ there is no consensus regarding how to define, measure, and report this outcome. Are patients who are extubated to noninvasive ventilation, for example, successfully extubated? At what time point should investigators report successful extubation? Which study participants should be included in the denominator (all participants randomized to an intervention or only those who are extubated)? Which is more important, the proportion of participants successfully extubated or the time to successful extubation?

In summary, mechanical ventilation research in the ICU is challenging. With their publication in this issue of *JAMA*, Subirà and colleagues have addressed an important knowledge gap, advancing the science of ventilator liberation and revitalizing an important area of critical care research. Their results support the use of 30 minutes of PSV during an initial SBT for most patients. Yet much work remains to be done to improve the outcomes of mechanically ventilated ICU patients, a population that both is highly vulnerable and has great potential to respond to evidence-based care.¹⁵

ARTICLE INFORMATION

Author Affiliations: Clinical Research, Investigation, and Systems Modeling of Acute Illness (CRISMA) Center, Department of Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (Girard); Interdepartmental Division of Critical Care, University of Toronto, Toronto, Ontario, Canada (Burns); Division of Critical Care Medicine, Department of Medicine, St Michael's Hospital, Toronto, Ontario, Canada (Burns); Li Ka Shing Knowledge Institute, St Michael's Hospital, Toronto, Ontario, Canada (Burns).

Corresponding Author: Timothy D. Girard, MD, MSCI, University of Pittsburgh School of Medicine, Critical Care Medicine, 3520 Fifth Ave, Ste 100, Pittsburgh, PA 15213 (timothy.girard@pitt.edu).

Conflict of Interest Disclosures: None reported.

REFERENCES

1. Thille AW, Harrois A, Schortgen F, Brun-Buisson C, Brochard L. Outcomes of extubation failure in

medical intensive care unit patients. *Crit Care Med*. 2011;39(12):2612-2618. doi:10.1097/CCM.0b013e3182282a5a

2. Cook DJ, Walter SD, Cook RJ, et al. Incidence of and risk factors for ventilator-associated pneumonia in critically ill patients. *Ann Intern Med*. 1998;129(6):433-440. doi:10.7326/0003-4819-129-6-199809150-00002

3. Schmidt GA, Girard TD, Kress JP, et al; ATS/CHEST Ad Hoc Committee on Liberation From Mechanical Ventilation in Adults. Official executive summary of an American Thoracic Society/American College of Chest Physicians clinical practice guideline: liberation from mechanical ventilation in critically ill adults. *Am J Respir Crit Care Med*. 2017;195(1):115-119. doi:10.1164/rccm.201610-2076ST

4. Burns KEA, Raptis S, Nisenbaum R, et al. International practice variation in weaning critically ill adults from invasive mechanical ventilation. *Ann Am Thorac Soc*. 2018;15(4):494-502. doi:10.1513/AnnalsATS.201705-4100C

5. Esteban A, Frutos F, Tobin MJ, et al; Spanish Lung Failure Collaborative Group. A comparison of four methods of weaning patients from mechanical ventilation. *N Engl J Med*. 1995;332(6):345-350. doi:10.1056/NEJM199502093320601

6. Ely EW, Baker AM, Dunagan DP, et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med*. 1996;335(25):1864-1869. doi:10.1056/NEJM199612193352502

7. Subirà C, Hernández G, Vázquez A, et al. Effect of pressure support vs T-piece ventilation strategies during spontaneous breathing trials on successful extubation among patients receiving mechanical ventilation: a randomized clinical trial [published June 11, 2019]. *JAMA*. doi:10.1001/jama.2019.7234

8. Esteban A, Alía I, Gordo F, et al; Spanish Lung Failure Collaborative Group. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. *Am J Respir Crit Care Med*. 1997;156(2, pt 1):459-465. doi:10.1164/ajrccm.156.2.9610109

9. Esteban A, Alía I, Tobin MJ, et al; Spanish Lung Failure Collaborative Group. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. *Am J Respir Crit Care Med*. 1999;159(2):512-518. doi:10.1164/ajrccm.159.2.9803106
10. Krinsley JS, Barone JE. The drive to survive: unplanned extubation in the ICU. *Chest*. 2005;128(2):560-566. doi:10.1378/chest.128.2.560
11. Vallverdú I, Calaf N, Subirana M, Net A, Benito S, Mancebo J. Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med*. 1998;158(6):1855-1862. doi:10.1164/ajrccm.158.6.9712135
12. Lellouche F, Mancebo J, Jolliet P, et al. A multicenter randomized trial of computer-driven protocolized weaning from mechanical ventilation. *Am J Respir Crit Care Med*. 2006;174(8):894-900. doi:10.1164/rccm.200511-1780OC
13. Perkins GD, Mistry D, Gates S, et al; Breathe Collaborators. Effect of protocolized weaning with early extubation to noninvasive ventilation vs invasive weaning on time to liberation from mechanical ventilation among patients with respiratory failure: the Breathe randomized clinical trial. *JAMA*. 2018;320(18):1881-1888. doi:10.1001/jama.2018.13763
14. Burns KE, Jacob SK, Aguirre V, Gomes J, Mehta S, Rizvi L. Stakeholder engagement in trial design: survey of visitors to critically ill patients regarding preferences for outcomes and treatment options during weaning from mechanical ventilation. *Ann Am Thorac Soc*. 2016;13(11):1962-1968. doi:10.1513/AnnalsATS.201606-445OC
15. Pun BT, Balas MC, Barnes-Daly MA, et al. Caring for critically ill patients with the ABCDEF bundle: results of the ICU Liberation Collaborative in over 15,000 adults. *Crit Care Med*. 2019;47(1):3-14. doi:10.1097/CCM.0000000000003482

Effect of Pressure Support vs T-Piece Ventilation Strategies During Spontaneous Breathing Trials on Successful Extubation Among Patients Receiving Mechanical Ventilation: A Randomized Clinical Trial

Carles Subirà, MD; Gonzalo Hernández, MD, PhD; Antònia Vázquez, MD, PhD; Raquel Rodríguez-García, MD; Alejandro González-Castro, MD; Carolina García, MD; Olga Rubio, MD, PhD; Lara Ventura, MD; Alexandra López, MD; María-Carmen de la Torre, MD, PhD; Elena Keough, MD; Vanesa Arauzo, MD; Cecilia Hermosa, MD; Carmen Sánchez, MD; Ana Tizón, MD; Eva Tenza, MD, PhD; César Laborda, MD; Sara Cabañes, MD; Victoria Lacueva, MD; María del Mar Fernández, MD, PhD; Anna Arnau, MSc, PhD; Rafael Fernández, RMD, PhD

IMPORTANCE Daily spontaneous breathing trials (SBTs) are the best approach to determine whether patients are ready for disconnection from mechanical ventilation, but mode and duration of SBT remain controversial.

OBJECTIVE To evaluate the effect of an SBT consisting of 30 minutes of pressure support ventilation (an approach that is less demanding for patients) vs an SBT consisting of 2 hours of T-piece ventilation (an approach that is more demanding for patients) on rates of successful extubation.

DESIGN, SETTING, AND PARTICIPANTS Randomized clinical trial conducted from January 2016 to April 2017 among 1153 adults deemed ready for weaning after at least 24 hours of mechanical ventilation at 18 intensive care units in Spain. Follow-up ended in July 2017.

INTERVENTIONS Patients were randomized to undergo a 2-hour T-piece SBT (n = 578) or a 30-minute SBT with 8-cm H₂O pressure support ventilation (n = 557).

MAIN OUTCOME AND MEASURES The primary outcome was successful extubation (remaining free of mechanical ventilation 72 hours after first SBT). Secondary outcomes were reintubation among patients extubated after SBT; intensive care unit and hospital lengths of stay; and hospital and 90-day mortality.

RESULTS Among 1153 patients who were randomized (mean age, 62.2 [SD, 15.7] years; 428 [37.1%] women), 1018 (88.3%) completed the trial. Successful extubation occurred in 473 patients (82.3%) in the pressure support ventilation group and 428 patients (74.0%) in the T-piece group (difference, 8.2%; 95% CI, 3.4%-13.0%; P = .001). Among secondary outcomes, for the pressure support ventilation group vs the T-piece group, respectively, reintubation was 11.1% vs 11.9% (difference, -0.8%; 95% CI, -4.8% to 3.1%; P = .63), median intensive care unit length of stay was 9 days vs 10 days (mean difference, -0.3 days; 95% CI, -1.7 to 1.1 days; P = .69), median hospital length of stay was 24 days vs 24 days (mean difference, 1.3 days; 95% CI, -2.2 to 4.9 days; P = .45), hospital mortality was 10.4% vs 14.9% (difference, -4.4%; 95% CI, -8.3% to -0.6%; P = .02), and 90-day mortality was 13.2% vs 17.3% (difference, -4.1% [95% CI, -8.2% to 0.01%; P = .04]; hazard ratio, 0.74 [95% CI, 0.55-0.99]).

CONCLUSIONS AND RELEVANCE Among patients receiving mechanical ventilation, a spontaneous breathing trial consisting of 30 minutes of pressure support ventilation, compared with 2 hours of T-piece ventilation, led to significantly higher rates of successful extubation. These findings support the use of a shorter, less demanding ventilation strategy for spontaneous breathing trials.

TRIAL REGISTRATION ClinicalTrials.gov Identifier: [NCT02620358](https://clinicaltrials.gov/ct2/show/study/NCT02620358)

JAMA. 2019;321(22):2175-2182. doi:10.1001/jama.2019.7234

[+ Visual Abstract](#)

[← Editorial page 2167](#)

[+ Supplemental content](#)

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Carles Subirà, MD, Althaia Xarxa Assistencial Universitària de Manresa, Dr Soler 1-3, 08243 Manresa, Spain (csubira@althaia.cat).

Section Editor: Derek C. Angus, MD, MPH, Associate Editor, JAMA (angusdc@upmc.edu).

Among patients receiving mechanical ventilation, readiness for extubation and liberation from ventilatory support is evaluated with a spontaneous breathing trial (SBT).¹ Daily screening of respiratory function by SBT is associated with a shorter duration of mechanical ventilation.² After a successful SBT and extubation, 10% to 25% of patients require reintubation, and reintubation is associated with higher mortality.^{3,4}

The most common modes of SBT are T-piece ventilation and pressure support ventilation (PSV), lasting between 30 minutes and 2 hours.⁵⁻⁷ There are no differences in the rate of successful extubation between 2-hour PSV and 2-hour T-piece ventilation,⁸ between T-piece ventilation for 30 minutes vs 2 hours,⁹ or between PSV for 30 minutes vs 2 hours.¹⁰ Although shorter SBTs are better tolerated, there is no evidence that they result in higher successful extubation rates.^{9,10} Some patients in whom a T-piece SBT failed might have been successfully extubated after a PSV SBT.¹¹

A recent meta-analysis suggested that T-piece SBTs are the optimal method for evaluating weaning readiness.¹² Nevertheless, another meta-analysis found that PSV resulted in higher rates of successful extubation than T-piece SBTs.¹³ Moreover, the latest American Thoracic Society guidelines for weaning recommend PSV SBTs with moderate-quality evidence.¹⁴ Thus, further investigation is needed to determine the best approach for SBTs.

This study hypothesized that less demanding SBTs could result in a higher rate of successful extubation without increasing the reintubation rate. To test this hypothesis, 2 weaning strategies were compared: an approach that is more demanding for patients (T-piece SBT for 2 hours) vs an approach that is less demanding for patients (8-cm H₂O PSV for 30 minutes).

Methods

From January 2016 through April 2017, a multicenter randomized clinical trial was conducted in 18 Spanish intensive care units. The ethics committee of each hospital approved the study, and all patients or their relatives provided written informed consent. The study protocol is available in [Supplement 1](#).

Patients aged 18 years or older undergoing mechanical ventilation for at least 24 hours who fulfilled the weaning criteria were eligible. The weaning criteria were (1) the resolution or improvement of the condition leading to intubation; (2) hemodynamic stability, defined as systolic blood pressure between 90 and 160 mm Hg and heart rate less than 140/min without vasopressors or with low doses of vasopressors; (3) Glasgow Coma Scale score of 13 or greater; (4) respiratory stability (oxygen saturation >90% with fraction of inspired oxygen [FiO₂] ≤0.4, respiratory rate <35/min, spontaneous tidal volume >5 mL/kg, ratio of respiratory rate to tidal volume <100/min per liter, and maximal inspiratory pressure >15 cm H₂O); and (5) noncopious secretions (<3 aspirations in the last 8 hours). Patients with tracheostomies or do-not-reintubate orders were excluded.

Randomization

Patients were randomized in a 1:1 ratio to one of the two weaning strategies by means of tables of computer-generated ran-

Key Points

Question What is the effect of a less demanding (30 minutes of pressure support ventilation) vs a more demanding (2 hours of T-piece ventilation) spontaneous breathing trial on rates of successful extubation?

Findings In this randomized clinical trial that included 1153 adults receiving mechanical ventilation, the proportion of patients successfully extubated was 82.3% among those who received 30 minutes of pressure support ventilation compared with 74% among those who received 2 hours of T-piece ventilation, a difference that was statistically significant.

Meaning These findings support the use of a shorter, less demanding strategy of 30 minutes of pressure support ventilation for spontaneous breathing trials.

dom numbers in blinded blocks of 4 patients for each center. A central administrator who was not involved in the analyses used an opaque envelope to allocate patients to receive one of the two treatments. The intervention was not blinded for the investigators or attending physicians.

Interventions

Patients randomized to undergo a highly demanding SBT underwent a 2-hour T-piece SBT; patients randomized to undergo a less demanding SBT underwent a 30-minute SBT with 8-cm H₂O PSV and zero positive end-expiratory pressure; FiO₂ remained unchanged from the mechanical ventilation period leading up to the SBT.

Before randomization, attending physicians had to decide on the extubation strategy (whether to reconnect the patient to the ventilator for 1 hour before extubation and whether to administer noninvasive ventilation or high-flow nasal cannula after extubation).

Patients who successfully completed the SBT were extubated. Arterial blood gas analysis was not required, but when it was done, the results were recorded. Physicians were also recommended to record dyspnea using the Borg Dyspnea Scale (score range, 0-10; 0 indicates no dyspnea and 10 indicates maximal dyspnea) at the beginning and at the end of SBTs and to ask patients about their confidence in their ability to sustain breathing without a ventilator.

Patients who did not tolerate the SBT were reconnected to a ventilator. Criteria for failure to tolerate the SBT were agitation, anxiety, low level of consciousness (Glasgow Coma Scale score <13), respiratory rate higher than 35/min and/or use of accessory muscles, oxygen saturation by pulse oximetry less than 90% with FiO₂ higher than 0.5, heart rate higher than 140/min or greater than a 20% increase from baseline, systolic blood pressure lower than 90 mm Hg, or development of arrhythmia. Additional SBTs were not protocolized, and mode and duration were left to the discretion of attending teams.

Respiratory failure within 72 hours of extubation was defined as the occurrence of at least 1 of the following: respiratory acidosis with pH lower than 7.32 and PaCO₂ higher than 45 mm Hg, oxygen saturation less than 90% with FiO₂ higher than 0.5, respiratory rate higher than 35/min, low level of consciousness (Glasgow Coma Scale score <13), severe agitation,

or clinical signs of respiratory fatigue. Treatment of postextubation respiratory failure was not protocolized. When noninvasive ventilation was used, duration, maximum inspiratory and expiratory pressures, and maximum FiO_2 were recorded. When respiratory failure was treated with a high-flow nasal cannula, duration, maximum flow, and maximum FiO_2 were recorded.

Patients needing reintubation within 72 hours were not randomized again for weaning, but the need for tracheostomy and the date of final liberation from mechanical ventilation were registered.

Outcomes

The primary outcome was successful extubation, defined as remaining free of invasive mechanical ventilation 72 hours after the first SBT.

Secondary outcomes were rate of reintubation among patients who were extubated after the SBT; intensive care unit and hospital lengths of stay; and hospital and 90-day mortality.

Exploratory outcomes were time to reintubation and reasons for reintubation, incidence of tracheostomy, and use of noninvasive ventilation and high-flow nasal cannula as prophylaxis against postextubation respiratory failure and to treat it.

Post hoc outcomes were intensive care unit mortality, Borg Dyspnea Scale score at the end of the SBT, patients' confidence in their ability to breathe without the ventilator, and arterial blood analysis after successful SBT.

Statistical Analysis

Based on previous studies,^{8,9} a successful extubation rate of 75% and an absolute increase in successful extubation of 7% were expected. Thus, the required sample for an $\alpha=.05$ and a power of 80% was estimated to be 540 patients in each group.

A prespecified interim analysis was performed when 500 patients were enrolled. The results showed a nonsignificant difference in primary outcome between groups. For this reason, the investigators decided to complete the estimated sample enrollment.

All patients were analyzed in the group to which they were randomized using the intention-to-treat principle, with no exclusion after randomization. Patients extubated outside of protocol were analyzed as having a failed SBT. No participants were excluded from main or secondary analyses because of missing or incomplete data. Reintubation was recorded only among patients who completed the trial.

Categorical variables are presented as absolute and relative frequencies. Continuous variables are summarized as medians and interquartile ranges (IQRs) for nonnormal distributions. The Mann-Whitney U was used for nonparametric continuous variables. To compare categorical variables, the χ^2 test was used, except when expected frequencies in contingency tables were less than 5, in which case the Fisher exact test or the Monte Carlo method was used.

Time-to-event outcomes were analyzed with Kaplan-Meier curves and compared by log-rank test. For the time-to-event outcome of 72-hour successful extubation, deaths occurring before 72 hours were introduced in the survival analysis as censored data. Event or censored times for all patients were

calculated from the time of randomization. Crude hazard ratios and 95% confidence intervals were calculated using a univariable Cox proportional regression model to estimate the effect size of randomization group. Proportionality of hazards was verified by examining Schoenfeld residual plots.

A post hoc random-effects multilevel logistic regression model was used to determine variables associated with 72-hour successful extubation, taking into account the effect of hospital. Patient characteristics that were associated with 72-hour successful extubation in the bivariable analysis were introduced in the random-effects multilevel logistic regression model as first-level variables and hospital as a second-level variable (random effect). Odds ratios (ORs) and median ORs with 95% confidence intervals were used to measure the association between each covariate and 72-hour successful extubation. The median OR is a measure of the variation between rates of 72-hour successful extubation at different hospitals that is unexplained by the modeled risk factors; it is defined as the median of the set of ORs that could be obtained by comparing 2 patients with identical patient-level characteristics from 2 randomly chosen hospitals. Covariates were introduced in the random-effects multilevel logistic regression model using a researcher-controlled backward exclusion strategy.

Post hoc analyses were performed for primary, secondary, exploratory, and post hoc outcomes among the following populations: patients extubated after the first SBT, patients extubated outside of protocol, patients treated per protocol, and several subgroups defined by baseline demographic characteristics. Effect sizes were evaluated by computing absolute risk differences with 95% confidence intervals for binary outcomes and differences in means with 95% confidence intervals for continuous outcomes. Figures were plotted for unadjusted risk ratios and 95% confidence intervals in the subgroup analysis by age; days of mechanical ventilation; Acute Physiology and Chronic Health Evaluation (APACHE) II score; chronic obstructive pulmonary disease (COPD); and medical, surgical, or trauma admission. No tests for interaction were conducted for the subgroup analyses.


A 2-sided $\alpha=.05$ was considered statistically significant. Data were analyzed using SPSS version 22 (IBM Corp) and Stata version 14 (StataCorp). Subgroup analysis graphs were generated using R version 3.5.2 (R Foundation for Statistical Computing). There was no adjustment for multiple comparisons. Therefore, the results of the subgroup analyses and the analyses for secondary and exploratory outcomes should be interpreted as exploratory.

Results

Study Participants

Figure 1 shows participant flow through the trial. During the study period, 2649 patients received mechanical ventilation for at least 24 hours in the participating intensive care units; 1501 of these fulfilled the inclusion criteria, and 1153 were included in the study; 578 patients were randomized to undergo a 2-hour T-piece SBT and 575 patients were randomized to undergo a 30-minute SBT with 8-cm H_2O PSV). The

Figure 1. Flow of Participants in a Trial of Pressure Support vs T-Piece Ventilation Strategies for SBTs



2 groups were similar in age, sex, APACHE II score on admission, reason for intensive care unit admission, and length of mechanical ventilation before the SBT (Table 1). No patients were lost to follow-up.

Primary Outcome

Successful extubation, defined as remaining free of mechanical ventilation 72 hours after the SBT, occurred in 473 patients (82.3%) in the PSV group and 428 patients (74%) in the T-piece group (difference, 8.2%; 95% CI, 3.4%-13%) (Table 2).

The Kaplan-Meier curves show a significant difference, with a higher successful extubation rate in the PSV group (hazard ratio, 1.54; 95% CI, 1.19-1.97; $P < .001$) (Figure 2).

Secondary Outcomes

After the first SBT, 486 patients (92.5%) undergoing the 30-minute PSV SBT and 532 patients (84.1%) undergoing the 2-hour T-piece SBT were extubated (difference, 8.4%; 95% CI, 4.7%-12.1%; $P < .001$). Reintubation within 72 hours occurred in 59 patients (11.1%) in the PSV group and in 58 patients (11.9%) in the T-piece group (difference, -0.8% ; 95% CI, -4.8% to -3.1% ; $P = .63$) (Table 2). The median intensive care unit length of stay was 9 days (IQR, 5-17) in the PSV group and 10 days (IQR, 5-17) in the T-piece group (mean difference, -0.3 days; 95% CI, -1.7 to 1.1 days; $P = .69$). The median hospital length of stay was 24 days (IQR, 15-40) in the PSV group and 24 days (IQR, 15-39) in the T-piece group (mean difference, 1.3 days; 95% CI, -2.2 to 4.9 days; $P = .45$). Hospital mortality rates were 10.4% ($n = 60$) in the PSV group and 14.9% ($n = 86$) in the T-piece group (difference, -4.4% ; 95% CI, -8.3% to -0.6% ; $P = .02$) (Table 2).

Mortality at 90 days was significantly lower in the PSV group (13.2%) compared with the T-piece group (17.3%)

(difference, -4.1% [95% CI, -8.2% to 0.01% ; $P = .04$]; hazard ratio, 0.74 [95% CI, 0.55-0.99]) (eFigure 1 in Supplement 2).

Exploratory Outcomes

In the T-piece group, 58 patients required reintubation, and in the PSV group, 59 patients required reintubation. The median time to reintubation was 23 hours (IQR, 9-45 hours) in the PSV group and 24.5 hours (IQR, 9.8-44 hours) in the T-piece group (mean difference, 0.53 hours; 95% CI, -7.2 to 8.3 hours). Reasons for reintubation were not significantly different in the 2 groups; excessive work of breathing was the most common in both groups, followed by inability to clear secretions and hypoxemia (Table 3). Four patients (3 in the T-piece group and 1 in the PSV group) had cardiac arrest within 72 hours after extubation.

Among reintubated patients, tracheostomy was performed in 41 patients (7.1%) in the PSV group and in 50 patients (8.7%) in the T-piece group (difference, -1.5% ; 95% CI, -4.6% to 1.6%) (Table 2).

Before randomization, physicians had to decide on an extubation strategy (standard oxygen, reconnection to the ventilator for a 1-hour rest after the SBT, and/or prophylactic non-invasive ventilation or high-flow nasal cannula after extubation). The use of each treatment was not significantly different in both groups (Table 1).

Postextubation respiratory failure occurred in 110 patients (20.7%) in the PSV group and in 103 patients (21.2%) in the T-piece group (difference, -0.5% ; 95% CI, -5.5% to 4.5%). Among these 213 patients, 117 (11.4%) were reintubated. Respiratory failure was treated by noninvasive ventilation in 91 (42.7%) patients, and 36 (39.6%) of these patients were reintubated. Respiratory failure was treated by high-flow nasal cannula in 47 (22.1%) patients, and 20 (42.6%) of these patients were reintubated. The remaining 75 patients

(35.2%) received standard oxygen, and 61 (81.3%) of these patients were reintubated.

Post Hoc Analysis

In patients extubated after the first SBT, the 72-hour successful extubation rate was not significantly different between groups (eTable 1 in Supplement 2).

The post hoc analysis showed that 29 patients (5%) in the PSV group and 38 patients (6.6%) in the T-piece group died in the intensive care unit (difference, -1.5%; 95% CI, -4.5% to 1.1%).

Multilevel logistic regression found a hospital-level random effect on successful extubation (median OR, 1.56; $P < .001$). After adjustment for this random effect, the effect of the PSV persisted (adjusted OR, 1.64; 95% CI, 1.23-2.20; $P = .001$). Other patient characteristics independently associated with 72-hour successful extubation were length of mechanical ventilation before SBT (adjusted OR, 0.96; 95% CI, 0.94-0.98; $P < .001$) and COPD (adjusted OR, 0.62; 95% CI, 0.44-0.87; $P = .006$). Multilevel logistic regression did not find an association between hospital and risk of reintubation (median OR, 1.19; $P = .30$). After adjustment for this random effect, reintubation in the PSV group was not significantly different than in the T-piece group (adjusted OR, 0.92; 95% CI, 0.62-1.35; $P = .67$). The only variable independently associated with reintubation was length of mechanical ventilation before SBT (adjusted OR, 1.04; 95% CI, 1.01-1.07; $P = .03$).

A total of 36 patients (3.1%) in whom the SBT failed were extubated, either because of physician decision or self-extubation during the SBT (eTable 2 in Supplement 2). The results of the per-protocol analysis were similar to those of the intention-to-treat analysis (eAppendix and eTable 3 in Supplement 2).

Subgroup analyses were generally consistent with the overall study findings (Figure 3; eFigures 2 and 3 in Supplement 2). eTables 4, 5, and 6 in Supplement 2 report post hoc analyses of Borg Dyspnea Scale scores at the end of the SBT, patients' confidence in breathing without a ventilator, and blood gas analyses.

During the study, there were no severe adverse events attributable to the randomization group. The adverse events that occurred after extubation, such as difficulty managing secretions or excessive work of breathing, are inherent to critically ill patients.

Discussion

In this randomized trial of patients receiving mechanical ventilation, a 30-minute PSV-SBT resulted in a significantly higher rate of successful extubation than a 2-hour T-piece SBT without significantly increasing reintubation. The higher rate was related to more patients being extubated after the PSV-SBT, suggesting that a less demanding SBT better allows critically ill patients to demonstrate their ability to sustain breathing.

A recent meta-analysis concluded that breathing through a T piece requires the same amount of work as breathing after extubation, and the authors recommended that SBTs should be performed with T pieces because this approach better reflects the physiologic conditions after extubation.¹² Sup-

Table 1. Baseline Participant Characteristics^a

| Characteristics | 30-min PSV SBT (n = 575) | 2-h T-Piece SBT (n = 578) |
|--|--------------------------|---------------------------|
| Age, median (IQR), y | 65 (52-75) | 63 (53-73) |
| Sex | | |
| Men | 352 (61.2) | 373 (64.5) |
| Women | 223 (38.8) | 205 (35.5) |
| APACHE II score, median (IQR) ^b | 16 (11-22) | 16 (11-22) |
| Comorbidity | | |
| Cardiovascular disease | 146 (25.4) | 162 (28.0) |
| Diabetes mellitus ^c | 123 (22.0) | 147 (25.8) |
| Chronic obstructive pulmonary disease | 110 (19.1) | 118 (20.4) |
| Neurological disease | 107 (18.6) | 99 (17.1) |
| Cancer | 87 (15.1) | 94 (16.3) |
| Renal disease | 76 (13.2) | 68 (11.8) |
| Liver disease | 64 (11.1) | 63 (10.9) |
| Reason for admission | | |
| Medical, nonrespiratory | 215 (37.4) | 206 (35.6) |
| Medical, respiratory | 189 (32.9) | 190 (32.9) |
| Emergency surgery | 105 (18.3) | 113 (19.6) |
| Planned surgery | 35 (6.1) | 29 (5.0) |
| Trauma | 31 (5) | 40 (6.9) |
| Length of mechanical ventilation before SBT, median (IQR), d | 4 (2-8) | 4 (2-8) |
| Reconnection to ventilator before extubation ^d | 145 (25.2) | 158 (27.3) |
| Prophylactic noninvasive ventilation after extubation ^d | 51 (8.9) | 34 (5.9) |
| Prophylactic high-flow nasal cannula after extubation ^d | 91 (15.8) | 74 (12.8) |

Abbreviations: APACHE, Acute Physiology and Chronic Health Evaluation; IQR, interquartile range; PSV, pressure support ventilation; SBT, spontaneous breathing trial.

^a Data are expressed as No. (%) of participants unless otherwise indicated.

^b Data were missing for 8 patients in the 2-hour T-piece SBT group and for 15 patients in the 30-minute PSV SBT group.

^c The APACHE II score ranges from 0 to 71, with higher scores indicating higher mortality risk. A patient with a score of 16 has an estimated mortality risk of 25%.

^d Attending physicians decided on the extubation strategy (reconnection to ventilator and prophylactic noninvasive ventilation or high-flow nasal cannula) prior to randomization.

ported by anecdotal reports, physicians may be concerned that some patients who breathe comfortably with low levels of PSV and/or positive end-expiratory pressure could develop respiratory failure immediately after extubation, which might even be followed by cardiac arrest.¹⁵ The results of this randomized trial designed to study extubation outcomes of opposing SBT strategies suggest that this concern may not be warranted. The current study found that the T-piece SBT was less well tolerated than the PSV SBT, although the work of breathing with the T piece may have been similar to breathing spontaneously. In patients who successfully completed the SBT, the reintubation rate was not significantly different in the 2 groups, and no imminent respiratory failure was observed after extubation from PSV. Moreover, the time to reintubation was about 24 hours in both groups, and the incidence of cardiac arrest was very low and even nominally higher in the T-piece group than in the PSV group.

Table 2. Primary, Secondary, Exploratory, and Post Hoc Outcomes of Patients Who Underwent a 2-Hour T-Piece SBT or a 30-Minute PSV SBT^a

| Outcomes | 30-min PSV SBT (n = 575) | 2-h T-Piece SBT (n = 578) | Difference, PSV SBT Minus T-Piece SBT (95% CI) ^b | P Value |
|--|--------------------------|---------------------------|---|---------|
| Primary Outcome | | | | |
| Successful extubation, No. (%) ^c | 473 (82.3) | 428 (74.0) | 8.2 (3.4 to 13.0) | .001 |
| Secondary Outcomes | | | | |
| Extubation after first SBT, No. (%) | 532 (92.5) | 486 (84.1) | 8.4 (4.7 to 12.1) | <.001 |
| Reintubation within 72 h, No. (%) ^d | 59 (11.1) | 58 (11.9) | -0.8 (-4.8 to 3.1) | .63 |
| ICU length of stay, median (IQR), d | 9 (5-17) | 10 (5-17) | -0.3 (-1.7 to 1.1) | .69 |
| Hospital length of stay, median (IQR), d | 24 (15-40) | 24 (15-39) | 1.3 (-2.2 to 4.9) | .45 |
| Hospital mortality, No. (%) | 60 (10.4) | 86 (14.9) | -4.4 (-8.3 to -0.6) | .02 |
| 90-Day mortality, No. (%) | 76 (13.2) | 100 (17.3) | -4.1 (-8.2 to 0.01) | .04 |
| Exploratory Outcome | | | | |
| Tracheostomy, No. (%) | 41 (7.1) | 50 (8.7) | -1.5 (-4.6 to 1.6) | .38 |
| Post Hoc Outcome | | | | |
| ICU mortality, No. (%) | 29 (5.0) | 38 (6.6) | -1.5 (-4.2 to 1.1) | .26 |

Abbreviations: ICU, intensive care unit; IQR, interquartile range; PSV, pressure support ventilation; SBT, spontaneous breathing trial.


^a No patients were lost to follow-up.

^b Differences were calculated as absolute risk differences for successful extubation, extubated after first SBT, reintubation within 72 hours, mortality, and tracheostomy and as difference in means for length of stay.

^c Defined as remaining free of mechanical ventilation for 72 hours after first SBT.

^d Among patients extubated after first SBT.

Figure 2. Probability of Successful Extubation After First SBT in Each Group



PSV indicates pressure support ventilation; SBT, spontaneous breathing trial. Successful extubation was defined as remaining free of mechanical ventilation for 72 hours after first SBT.

Vallverdú et al¹⁶ showed that among patients in whom a 2-hour T-piece SBT failed, 64% of failures occurred in the first 30 minutes, 12% between 30 and 60 minutes, and 24% between 60 minutes and 2 hours. In a recent observational study including 352 patients who underwent an SBT with PSV, Liang et al¹⁷ sought to identify the characteristics of the 41 patients (11.6%) in whom a 120-minute SBT failed after successful completion of the first 30 minutes. Patients with SBT failures after 30 minutes were older, had more cardiopulmonary disease, had spent more time receiving mechanical ventilation before the SBT, and had undergone more previous SBTs. The authors suggested that patients with these characteristics might need a longer SBT to ensure that their ability to breathe is not overestimated. Nevertheless, it is unknown what the outcomes of these patients would have been if the SBT had been limited to 30 minutes.

In the present study, the 30-minute PSV SBT was enough to check patients' ability to breathe without increasing the rates of postextubation respiratory failure and reintubation.

Another finding related to tolerance of the 2 SBT approaches is that self-extubation during the SBT was more common in the T-piece group. Tolerance to SBTs in this trial may be compared with the studies in the late 1990s by Esteban et al^{8,9}: patients' tolerance to T-piece SBTs in the present study was better than in the first trial by Esteban et al (84% vs 78%) and similar to their second trial (84.6% vs 84.1%). Moreover, tolerance to the 30-minute T-piece SBT in the second trial by Esteban et al was worse than tolerance to the 30-minute PSV in the present study (87.7% vs 92.5%). However, patients in the studies by Esteban et al received longer mechanical ventilation before the SBT, and this could contribute to worse tolerance and a higher reintubation rate.

In a single-center study comparing a 2-hour T-piece SBT and a 2-hour PSV SBT, Matic et al¹⁸ found a higher rate of successful extubation with PSV than with a T piece (80% vs 73%), similar to the difference found in the present study despite a longer duration of the PSV SBT. This suggests that tolerance is not only about duration but also about the mode of SBT. Along the same lines, Ezingear et al¹¹ found that some patients who did not tolerate a T-piece SBT went on to tolerate a PSV SBT and had a reintubation rate similar to patients who underwent a PSV SBT without having attempted a T-piece SBT. Taken together with these studies, the results of the present study suggest that a T-piece SBT is not the best way to check a patient's ability to breathe.

In this study, the reintubation rate was not significantly different between the 2 groups (about 11%), which is lower than the 17% in the first study by Esteban et al⁸ and similar to the 13% in their second study.⁹ Conversely, the reintubation rate was higher than in a study by Perren et al¹⁰ (9% for short SBTs and 4% for long SBTs), but that study's single-center design and small sample size preclude direct comparison.

Table 3. Exploratory Outcomes: Reasons for Reintubation


| Outcomes | 30-min PSV SBT (n = 59) | 2-h T-Piece SBT (n = 58) | Difference, PSV SBT Minus T-Piece SBT (95% CI) ^a |
|--|-------------------------|--------------------------|---|
| Time to reintubation, median (IQR), h | 23.0 (9.0-45.0) | 24.5 (9.8-44.0) | 0.53 (-7.2 to 8.3) |
| Reasons for reintubation, No. (%) ^b | | | |
| Excessive work of breathing | 23 (39.0) | 24 (41.4) | -2.4 (-20.2 to 15.4) |
| Difficulty managing secretions | 19 (32.2) | 18 (31.0) | 1.2 (-15.7 to 18.0) |
| Refractory hypoxemia | 14 (23.7) | 11 (19.0) | 4.8 (-10.1 to 19.6) |
| Level of consciousness | 6 (10.2) | 11 (19.0) | -8.7 (-21.5 to 3.9) |
| Airway obstruction | 6 (10.2) | 8 (13.8) | -3.6 (-15.4 to 8.1) |
| Surgery | 4 (6.8) | 4 (6.9) | -0.1 (-9.2 to 9.0) |
| Cardiac arrest | 1 (1.7) | 3 (5.1) | -3.4 (-10.0 to 3.1) |
| Agitation | 4 (6.8) | 3 (5.2) | 1.6 (-7.0 to 10.2) |
| Aspiration | 1 (1.7) | 3 (5.2) | -3.5 (-10.1 to 3.1) |
| Bradycardia (heart rate <50/min) | 0 | 1 (1.7) | -1.7 (-5.1 to 1.6) |
| Hemodynamic shock | 1 (1.7) | 1 (1.7) | -0.0 (-4.7 to 4.7) |

Abbreviations: IQR, interquartile range; PSV, pressure support ventilation; SBT, spontaneous breathing trial.

^a Differences were calculated as absolute risk differences for reasons for reintubation and as difference in means for time to reintubation.

^b Patients could have more than 1 reason for reintubation.

Figure 3. Unadjusted Risk Ratios for Successful Extubation After First SBT in Predefined Subgroups



PSV indicates pressure support ventilation; SBT, spontaneous breathing trial; APACHE, Acute Physiology and Chronic Health Evaluation; COPD, chronic obstructive pulmonary disease. Successful extubation was defined as remaining free of mechanical ventilation for 72 hours after first SBT.

Logistic regression analysis showed that the 30-minute PSV SBT was associated with successful extubation, whereas longer duration of mechanical ventilation before the SBT as well as COPD were associated with extubation failure. However, only length of mechanical ventilation was significantly associated with reintubation. This result lends additional support to the idea that the concern that PSV SBTs increase risk of respiratory failure and reintubation may be exaggerated.

Hospital mortality and 90-day mortality were significantly higher in the T-piece group. This finding cannot be explained by the reintubation rate, days of mechanical venti-

lation after a failed SBT, APACHE II score at admission, or hospital length of stay, which were not significantly different between the 2 groups.

Limitations

This study has several limitations. First, prophylactic use of noninvasive ventilation and high-flow nasal cannula after extubation was not protocolized. In some cases, these approaches were routinely used based on recent studies, but in others, they were used only in patients with more comorbidities, such as heart failure or COPD, or more risk factors for

extubation failure. For this reason, it is impossible to draw conclusions about the use of noninvasive ventilation and high-flow nasal cannula for postextubation respiratory failure.

Second, patients extubated outside of protocol, although few, could be expected to influence the main results, but the sensitivity analysis ruled out such bias (eTable 2 in Supplement 2).

Third, investigators and attending physicians were not blinded to treatment randomization group.

Conclusions

Among mechanically ventilated patients, an SBT consisting of 30 minutes of PSV, compared with 2 hours of T-piece ventilation, led to significantly higher rates of successful extubation. These findings support the use of a shorter, less demanding ventilation strategy for SBTs.

ARTICLE INFORMATION

Accepted for Publication: May 9, 2019.

Author Affiliations: Althaia Xarxa Assistencial Universitària de Manresa, Manresa, Spain (Subirà, Rubio, Ventura, Arnau, Fernández); Hospital Virgen de la Salud, Toledo, Spain (Hernández); Hospital del Mar, Barcelona, Spain (Vázquez); Hospital U, Central de Asturias, Oviedo, Spain (Rodríguez-García); Hospital Marqués de Valdecilla, Santander, Spain (González-Castro); Hospital Universitario de Canarias, Santa Cruz de Tenerife, Spain (García); Universitat Internacional de Catalunya, Barcelona, Spain (Rubio, Fernández); Hospital General de Granollers, Granollers, Spain (López); Hospital de Mataró, Mataró, Spain (de la Torre); Hospital Gregorio Marañón, Madrid, Spain (Keough); Consorci Sanitari de Terrassa, Terrassa, Spain (Arauzo); Hospital del Henares, Coslada, Spain (Hermosa); Complejo Hospitalario Universitario de Cáceres, Cáceres, Spain (Sánchez); Complejo Hospitalario Universitario de Ourense, Ourense, Spain (Tizón); Hospital d'Elx, Elche, Alicante, Spain (Tenza); Hospital Vall d'Hebron, Barcelona, Spain (Laborda); Hospital Universitario de Araba, Araba, Spain (Cabañes); Hospital de Sagunt, Sagunt, Spain (Lacueva); Hospital Mútua Terrassa, Terrassa, Spain (del Mar Fernández); CIBER Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain (Fernández).

Author Contributions: Dr Subirà had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Subirà, Keough, Fernández. **Acquisition, analysis, or interpretation of data:** Subirà, Hernández, Vázquez, Rodríguez-García, González-Castro, García, López, de la Torre, Rubio, Ventura, Keough, Arauzo, Hermosa, Sánchez, Tizón, Tenza, Cabañes, Laborda, Lacueva, del Mar Fernández, Arnau.

Drafting of the manuscript: Subirà, López, Ventura, Keough, Sánchez, Cabañes, Arnau, Fernández. **Critical revision of the manuscript for important intellectual content:** Hernández, Vázquez, Rodríguez-García, González-Castro, García, de la Torre, Rubio, Keough, Arauzo, Hermosa, Tizón, Tenza, Laborda, Lacueva, del Mar Fernández, Fernández.

Statistical analysis: Keough, Arnau, Fernández. **Obtained funding:** Subirà, García, Ventura. **Administrative, technical, or material support:** García, Ventura, Keough, Laborda, del Mar Fernández.

Supervision: Rubio, Keough, Tenza, Fernández.

Conflict of Interest Disclosures: Dr Hernández reported receipt of personal fees from Fisher & Paykel. Dr Keough reported consulting for Vapotherm Inc. No other disclosures were reported.

Funding/Support: This study was supported by an unrestricted grant from Societat Catalana de Medicina Intensiva i Crítica (SOCMIC).

Role of the Funder/Sponsor: SOCMIC had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Additional Contributions: We thank the patients and medical and nursing staff for their cooperation. John Giba, BSc, received financial compensation for services as a professional freelance translator.

Data Sharing Statement: See Supplement 3.

REFERENCES

- Esteban A, Frutos F, Tobin MJ, et al; Spanish Lung Failure Collaborative Group. A comparison of four methods of weaning patients from mechanical ventilation. *N Engl J Med*. 1995;332(6):345-350. doi:10.1056/NEJM199502093320601
- Ely EW, Baker AM, Dunagan DP, et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med*. 1996;335(25):1864-1869. doi:10.1056/NEJM199612193352502
- Thille AW, Cortés-Puch I, Esteban A. Weaning from the ventilator and extubation in ICU. *Curr Opin Crit Care*. 2013;19(1):57-64. doi:10.1097/MCC.0b013e32835c5095
- Jaber S, Quintard H, Cinotti R, et al. Risk factors and outcomes for airway failure versus non-airway failure in the intensive care unit: a multicenter observational study of 1514 extubation procedures. *Crit Care*. 2018;22(1):236. doi:10.1186/s13054-018-2150-6
- Esteban A, Anzueto A, Frutos F, et al; Mechanical Ventilation International Study Group. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA*. 2002;287(3):345-355. doi:10.1001/jama.287.3.345
- Peñuelas O, Frutos-Vivar F, Fernández C, et al; VENTILA Group. Characteristics and outcomes of ventilated patients according to time to liberation from mechanical ventilation. *Am J Respir Crit Care Med*. 2011;184(4):430-437. doi:10.1164/rccm.201011-1887OC
- Esteban A, Ferguson ND, Meade MO, et al; VENTILA Group. Evolution of mechanical ventilation in response to clinical research. *Am J Respir Crit Care Med*. 2008;177(2):170-177. doi:10.1164/rccm.200706-893OC
- Esteban A, Alía I, Gordo F, et al; Spanish Lung Failure Collaborative Group. Extubation outcome after spontaneous breathing trials with T-tube or

pressure support ventilation. *Am J Respir Crit Care Med*. 1997;156(2, pt 1):459-465. doi:10.1164/ajrccm.156.2.9610109

9. Esteban A, Alía I, Tobin MJ, et al; Spanish Lung Failure Collaborative Group. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. *Am J Respir Crit Care Med*. 1999;159(2):512-518. doi:10.1164/ajrccm.159.2.9803106

10. Perren A, Domenighetti G, Mauri S, Genini F, Vizzardi N. Protocol-directed weaning from mechanical ventilation: clinical outcome in patients randomized for a 30-min or 120-min trial with pressure support ventilation. *Intensive Care Med*. 2002;28(8):1058-1063. doi:10.1007/s00134-002-1353-z

11. Ezingard E, Diconne E, Guyomarc'h S, et al. Weaning from mechanical ventilation with pressure support in patients failing a T-tube trial of spontaneous breathing. *Intensive Care Med*. 2006; 32(1):165-169. doi:10.1007/s00134-005-2852-5

12. Sklar MC, Burns K, Rittayamai N, et al. Effort to breathe with various spontaneous breathing trial techniques: a physiologic meta-analysis. *Am J Respir Crit Care Med*. 2017;195(11):1477-1485. doi:10.1164/rccm.201607-1338OC

13. Burns KEA, Soliman I, Adhikari NKJ, et al. Trials directly comparing alternative spontaneous breathing trial techniques: a systematic review and meta-analysis. *Crit Care*. 2017;21(1):127. doi:10.1186/s13054-017-1698-x

14. Schmidt GA, Girard TD, Kress JP, et al. Liberation from mechanical ventilation in critically ill adults: executive summary of an official American College of Chest Physicians/American Thoracic Society clinical practice guideline. *Chest*. 2017;151(1):160-165. doi:10.1016/j.chest.2016.10.037

15. Tobin MJ. Extubation and the myth of "minimal ventilator settings". *Am J Respir Crit Care Med*. 2012;185(4):349-350. doi:10.1164/rccm.201201-0050ED

16. Vallverdú I, Calaf N, Subirana M, Net A, Benito S, Mancebo J. Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med*. 1998;158(6):1855-1862. doi:10.1164/ajrccm.158.6.9712135

17. Liang G, Liu T, Zeng Y, et al. Characteristics of subjects who failed a 120-minute spontaneous breathing trial. *Respir Care*. 2018;63(4):388-394. doi:10.4187/respcare.05820

18. Matic I, Majeric-Kogler V. Comparison of pressure support and T-tube weaning from mechanical ventilation: randomized prospective study. *Croat Med J*. 2004;45(2):162-166.