Weaning from Mechanical Ventilation

Have We, so Far, Missed a Crucial Point?

Giacomo Grasselli, M.D., Giacomo Bellani, M.D., Ph.D.

UNCTIONAL impairment of respiratory muscles is considered one of the determinants of difficult weaning from the ventilator,¹ which results in prolonged mechanical ventilation and is associated with worse patient outcome.² Patients requiring invasive ventilation are exquisitely prone to the development of respiratory muscle dysfunction (including ventilator-induced diaphragmatic dysfunction), which is worsened by a number of concomitant factors like sepsis, electrolyte imbalance, malnutrition, and administration of aminoglicosides, steroids, and neuromuscular blockers. Most of the studies tackling this issue were focused on the dysfunction of inspiratory muscles, mostly the dia-



"...[E]xpiratory muscle recruitment significantly contributes to the total respiratory effort of the patient during weaning from mechanical ventilation."

phragm. At variance, very few studies aimed at assessing the role of expiratory muscles.

In healthy subjects at rest, expiration is a passive process: when the inspiratory muscles relax at the end of inhalation, the elastic recoil of the lung tissue (*i.e.*, the "release" of the elastic energy stored during lung inflation) generates a positive pressure gradient between the alveoli and the atmosphere, causing the outflow of air. Although expiration is usually passive, it can become an active process involving the contraction of muscles with expiratory activity, in particular abdominal muscles (which push the diaphragm upward, reducing the volume of the chest cavity) and internal intercostal muscles (which pull the ribs downward, compressing the rib cage). This happens during forced breathing and in some pathologic conditions characterized by airways obstruction (like asthma or chronic obstructive pulmonary disease). On one hand, recruitment of expiratory muscles may facilitate the subsequent inspiration and limit hyperinflation; on the other hand, it may result in a significant increase in the oxygen consumption of respiratory muscles (i.e., in the work of breathing). Another mechanism acting to limit hyperinflation in the presence of airway obstruction is the persistence of diaphragm contraction during exhalation: diaphragmatic "tonic this activity" has been described in different experimental models and might contribute to stabilize peripheral airways and prevent expiratory lung collapse.

In the current issue of ANES-THESIOLOGY, Doorduin *et al.*³ report the results of an interesting study exploring the role

of expiratory muscle recruitment and electrical diaphragmatic postinspiratory activity in patients undergoing a weaning trial after at least 3 days of mechanical ventilation. The spontaneous breathing trial was performed with a T-tube (no positive end-expiratory pressure) for at least 60 min. Patients were instrumented with a modified nasogastric tube equipped with an esophageal and a gastric balloon and with microelectrodes for diaphragm electromyography recording, underlining the potential for electromyography as a monitoring tool,^{4,5} irrespective of the application of neurally adjusted ventilatory assist. Expiratory muscle recruitment was computed as the pressure-time product of gastric pressure, and tonic activity of the diaphragm was expressed as a percentage of the peak inspiratory diaphragm electromyography activity during the last quartile of neural expiration. As expected, patients who failed weaning (*i.e.*, those who failed the spontaneous breathing trial

Copyright © 2018, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2018; 129:00-00

Image: J. P. Rathmell.

Accepted for publication May 29, 2018. From the Department of Anesthesiology, Intensive Care and Emergency, Foundation IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy (G.G.); Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy (G.G.); Department of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy (G.B.); and Department of Emergency and Intensive Care, San Gerardo Hospital, Monza, Italy (G.B.).

or required reintubation within 48 h from extubation) had a higher rapid shallow breathing index, a higher inspiratory electrical activity of the diaphragm, and a significantly lower neuromechanical efficiency of diaphragm contraction (indicative of an impaired force-generating capacity of the diaphragm per unit of electrical activity). Weaning-failure patients also presented a significant increase in expiratory gastric pressure and the contribution of expiratory muscles to the total pressure-time product increased from 13% in the weaning-success group to 24% in the weaning-failure group.

These findings indicate that recruitment of expiratory muscles is increased in patients failing weaning from mechanical ventilation and account for a significant proportion of the total energy expenditure of respiratory muscles. On the contrary, tonic diaphragmatic activity was not different between the two patient subgroups. However, this finding is not unexpected because tonic contraction of the diaphragm during expiration seems to play a meaningful role in specific patient subgroups (*e.g.*, neonates) or in particular experimental settings (*e.g.*, as a means to prevent derecruitment⁶), whereas its relevance in adult subjects with acute or chronic respiratory failure remains to be demonstrated.

The study by Doorduin *et al.*³ highlights the concept that expiratory muscle recruitment significantly contributes to the total respiratory effort of the patient during weaning from mechanical ventilation. From the findings of the present article, it is unknown whether the presence of expiratory activity is only "associated" with weaning failure, for example in patients with worst respiratory system mechanics, or whether it is a factor also contributing to weaning failure, for example by increasing oxygen consumption.⁷ In addition, this study underlines the importance of monitoring the work of breathing and respiratory mechanics in patients failing a weaning attempt to identify the mechanisms contributing to the failure and to drive the therapeutic approach.⁸

The experimental setup implemented in the study patients is relatively complex, requiring the simultaneous acquisition of esophageal and gastric pressure signals and of diaphragm electromyography. Hence, an adequate knowledge and level of experience are necessary to obtain reliable and reproducible data, to integrate the information, and to apply them at the bedside. The development of noninvasive techniques based on surface electromyography⁹ of both inspiratory and expiratory muscles may represent a significant advance in this respect, although signal interference and electrical crosstalks might arise as relevant technical limitations. Doorduin *et al.*³ should be congratulated for performing such a complex, elegant study and for dragging our attention to an issue that should receive more attention in daily clinical practice and become the object of further scrutiny in research.

Competing Interests

The authors are not supported by, nor maintain any financial interest in, any commercial activity that may be associated with the topic of this article.

Correspondence

Address correspondence to Dr. Bellani: giacomo.bellani1@ unimib.it

References

- Perren A, Brochard L: Managing the apparent and hidden difficulties of weaning from mechanical ventilation. Intensive Care Med 2013; 39:1885–95
- Goligher EC, Dres M, Fan E, Rubenfeld GD, Scales DC, Herridge MS, Vorona S, Sklar MC, Rittayamai N, Lanys A, Murray A, Brace D, Urrea C, Reid WD, Tomlinson G, Slutsky AS, Kavanagh BP, Brochard LJ, Ferguson ND: Mechanical ventilation-induced diaphragm atrophy strongly impacts clinical outcomes. Am J Respir Crit Care Med 2018; 197:204–13
- 3. Doorduin J, Roesthuis LH, Jansen D, van der Hoeven JG, van Hees HWH, Heunks LMA: Respiratory muscle effort during expiration in successful and failed weaning from mechanical ventilation. ANESTHESIOLOGY 2018 May 16 [Epub ahead of print]
- Bellani G, Mauri T, Coppadoro A, Grasselli G, Patroniti N, Spadaro S, Sala V, Foti G, Pesenti A: Estimation of patient's inspiratory effort from the electrical activity of the diaphragm. Crit Care Med 2013; 41:1483–91
- Bellani G, Coppadoro A, Patroniti N, Turella M, Arrigoni Marocco S, Grasselli G, Mauri T, Pesenti A: Clinical assessment of auto-positive end-expiratory pressure by diaphragmatic electrical activity during pressure support and neurally adjusted ventilatory assist. ANESTHESIOLOGY 2014; 121:563–71
- Pellegrini M, Hedenstierna G, Roneus A, Segelsjö M, Larsson A, Perchiazzi G: The diaphragm acts as a brake during expiration to prevent lung collapse. Am J Respir Crit Care Med 2017; 195:1608–16
- Bellani G, Foti G, Spagnolli E, Milan M, Zanella A, Greco M, Patroniti N, Pesenti A: Increase of oxygen consumption during a progressive decrease of ventilatory support is lower in patients failing the trial in comparison with those who succeed. ANESTHESIOLOGY 2010; 113:378–85
- 8. Silva S, Ait Aissa D, Cocquet P, Hoarau L, Ruiz J, Ferre F, Rousset D, Mora M, Mari A, Fourcade O, Riu B, Jaber S, Bataille B: Combined thoracic ultrasound assessment during a successful weaning trial predicts postextubation distress. ANESTHESIOLOGY 2017; 127:666–74
- Schmidt M, Kindler F, Gottfried SB, Raux M, Hug F, Similowski T, Demoule A: Dyspnea and surface inspiratory electromyograms in mechanically ventilated patients. Intensive Care Med 2013; 39:1368–76

2

Respiratory Muscle Effort during Expiration in Successful and Failed Weaning from Mechanical Ventilation

Jonne Doorduin, Ph.D., Lisanne H. Roesthuis, M.Sc., Diana Jansen, M.D., Johannes G. van der Hoeven, M.D., Ph.D., Hieronymus W. H. van Hees, Ph.D., Leo M. A. Heunks, M.D., Ph.D.

ABSTRACT

Background: Respiratory muscle weakness in critically ill patients is associated with difficulty in weaning from mechanical ventilation. Previous studies have mainly focused on inspiratory muscle activity during weaning; expiratory muscle activity is less well understood. The current study describes expiratory muscle activity during weaning, including tonic diaphragm activity. The authors hypothesized that expiratory muscle effort is greater in patients who fail to wean compared to those who wean successfully.

Methods: Twenty adult patients receiving mechanical ventilation (more than 72 h) performed a spontaneous breathing trial. Tidal volume, transdiaphragmatic pressure, diaphragm electrical activity, and diaphragm neuromechanical efficiency were calculated on a breath-by-breath basis. Inspiratory (and expiratory) muscle efforts were calculated as the inspiratory esophageal (and expiratory gastric) pressure–time products, respectively.

Results: Nine patients failed weaning. The contribution of the expiratory muscles to total respiratory muscle effort increased in the "failure" group from $13\pm9\%$ at onset to $24\pm10\%$ at the end of the breathing trial (P = 0.047); there was no increase in the "success" group. Diaphragm electrical activity (expressed as the percentage of inspiratory peak) was low at end expiration (failure, $3\pm2\%$; success, $4\pm6\%$) and equal between groups during the entire expiratory phase (P = 0.407). Diaphragm neuromechanical efficiency was lower in the failure *versus* success groups ($0.38\pm0.16 vs$, $0.71\pm0.36 \text{ cm H}_2\text{O}/\mu\text{V}$; P = 0.054). **Conclusions:** Weaning failure (*vs.* success) is associated with increased effort of the expiratory muscles and impaired neuromechanical efficiency of the diaphragm but no difference in tonic activity of the diaphragm. **(ANESTHESIOLOGY 2018; 129:490-501)**

PROLONGED weaning from mechanical ventilation develops in 6 to 15% of mechanically ventilated patients and is associated with increased morbidity and mortality.^{1–4} A major determinant of weaning failure is respiratory muscle dysfunction.^{5–7} The respiratory muscles are profoundly affected by critical illness and mechanical ventilation.^{8–15} The respiratory muscle pump is made up of a number of muscles, but research in weaning failure has mainly focused on the inspiratory muscles, in particular the diaphragm.^{16–20} The role of the expiratory muscles including the internal intercostals and abdominal wall muscles during weaning failure is less well understood.

Recruitment of the expiratory muscles has been demonstrated in patients with chronic obstructive pulmonary disease (COPD) during a failed weaning trial.^{21,22} Expiratory muscle recruitment may help inspiration because the active reduction in end-expiratory lung volume stores

What We Already Know about This Topic

 Inspiratory muscle weakness delays weaning from mechanical ventilation, but the contributions of expiratory muscles activity are less well understood.

What This Article Tells Us That Is New

• Twenty adult patients receiving mechanical ventilation (more than 72 h) performed a spontaneous breathing trial. Nine patients failed weaning (11 succeeded), and these patients exhibited increased effort of the expiratory muscles and impaired neuromechanical efficiency of the diaphragm but no difference in tonic activity of the diaphragm.

elastic energy, facilitating subsequent inspiration. In addition, active reduction in end-expiratory lung volume may help to limit hyperinflation.²³ However, recruitment of the expiratory muscles increases the energy expenditure of

Copyright © 2018, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2018; 129:490-501

Corresponding article on page 394. Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are available in both the HTML and PDF versions of this article. Links to the digital files are provided in the HTML text of this article on the Journal's Web site (www.anesthesiology.org).

Submitted for publication March 22, 2017. Accepted for publication April 11, 2018. From the Departments of Critical Care Medicine (J.D., L.H.R., J.G.v.d.H., L.M.A.H.), Anesthesiology (D.J.), and Pulmonary Diseases (H.W.H.v.H.) and the Donders Institute for Brain, Cognition and Behaviour, Department of Neurology (J.D.), Radboud University Medical Center, Nijmegen, The Netherlands; and the Department of Intensive Care Medicine, VU University Medical Center, Amsterdam, The Netherlands (L.M.A.H.).

breathing, although the relative contribution of expiratory muscle effort to total breathing effort in patients weaning the ventilator is unknown. Expiratory muscles effort can be quantified as the pressure-time product of gastric pressure during expiration.^{24,25} Several studies have demonstrated that the diaphragm exhibits tonic activity during expiration.^{26–31} Recently, Pellegrini *et al.*³² have demonstrated in pigs with mild acute respiratory distress syndrome that the diaphragm acts as a brake during expiration to prevent lung collapse. In weaning-failure patients, tonic activity of the diaphragm during expiration might prevent airway closure and consequently limit intrinsic positive end-expiratory pressure (PEEP).

Accordingly, the aim of the current physiologic study was to quantify the expiratory pressure-time product and the expiratory activity of the diaphragm in weaning-success and weaning-failure patients. In addition, we studied respiratory muscle activity effort during inspiration. We hypothesized that expiratory muscle effort is higher in weaning-failure patients compared to successfully weaned patients.

Materials and Methods

Study Design and Population

This cross-sectional physiologic study was conducted in the intensive care unit of the Radboud University Medical Center, Nijmegen, The Netherlands. We recruited 20 adult patients invasively ventilated for at least 3 days and considered ready for a spontaneous breathing trial. *A priori* sample size calculation was not performed, but the number of required subjects was based on previous physiologic studies in mechanically ventilated patients.^{33–36} The decision to extubate or resume mechanical ventilation was made solely by the clinical team, blinded to the study data. Exclusion criteria were a past medical history of neuromuscular disorders, upper airway or esophageal pathology (*e.g.*, recent surgery, esophageal varices, diaphragmatic hernia), and recent (less than 1 month) nasal bleeding.

The protocol was approved by the local ethics review committee (approval number 2010-058) and conducted in accordance with the Declaration of Helsinki and its later amendments. Written informed consent was obtained in patients that were not already instrumented with the dedicated nasogastric catheter.

Study Protocol

All patients were ventilated with the SERVO-i ventilator (Maquet Critical Care, Sweden). Per clinical protocol, sedatives were discontinued before the spontaneous breathing trial, and patients were verified to be responsive and adequate. If not already *in situ*, a nasogastric catheter with multiple electrodes and two balloons (NeuroVent Research, Inc., Canada) was inserted nasally. Catheter characteristics and positioning techniques have been described previously.^{37–39} Baseline data were acquired while patients were ventilated in pressure support mode with an inspiratory support of 8 cm H₂O and a PEEP level of 5 cm H₂O for 10 min. Subsequently, a spontaneous breathing trial with T-tube and supplemental oxygen was performed for up to 60 min, according to our clinical protocol. The criteria for spontaneous breathing trial failure were tachypnea (more than 35 breaths/min), low peripheral capillary oxygen saturation (less than 90%), tachycardia (more than 140 beats/ min), systolic hypertension (more than 180 mmHg), systolic hypotension (less than 90 mmHg), agitation, diaphoresis, and anxiety during the spontaneous breathing trial. Patients were extubated after a successful spontaneous breathing trial. Weaning failure was defined as a failed spontaneous breathing trial or reintubation within 48 h after extubation. Arterial blood samples were collected before disconnection from the ventilator and at the end of the spontaneous breathing trial before reinstitution of ventilator support.

Data Acquisition

Flow was measured with a Fleisch pneumotachograph (Hans Rudolph, USA) placed at the endotracheal tube or cannula. The pneumotachograph was connected to a differential pressure transducer (range \pm 50 kPa; Freescale, USA).

The air-filled esophageal balloon and gastric balloon of the nasogastric catheter were connected to differential pressure transducers (range \pm 50 kPa; Freescale). Pressure and flow signals were digitized (Porti 16, 22 bits, 1.4 μ V/least significant bit; TMSi, The Netherlands) at a sampling frequency of 2 kHz. Transdiaphragmatic pressure was calculated as esophageal pressure subtracted from gastric pressure.

Electrical activity of the diaphragm recorded from the electrodes of the nasogastric catheter were amplified and digitized (Porti 16, 22 bits, 71.5 nV/least significant bit; TMSi) at a sampling frequency of 2kHz. Signal processing of the electrical activity of the diaphragm was performed according to the method of Sinderby *et al.*^{40–42} Details are provided in the Supplemental Digital Content (http://links.lww.com/ALN/B717).

Flow, esophageal pressure, gastric pressure, and diaphragm electrical activity were acquired synchronously (maximal 0.5-ms delay between mechanics and diaphragm electrical activity) using dedicated software (NeuroVent Research, Inc.) and stored on a hard disk for offline analysis in a software routine developed for Matlab (R2014b; Mathworks, USA).

Data Analysis

The data were analyzed on a breath-by-breath basis. For the period of mechanical ventilation, a 2-min epoch at the end was selected for analysis. The duration of the spontaneous breathing trial varied from patient to patient. Therefore, data were analyzed at seven points in time: the first and last minute and five epochs of at least 1 min taken at equal time intervals in between. The epochs were visually inspected for artefacts, such as movement and esophageal and gastric contractions. In the presence of artefacts, an adjacent



Fig. 1. Example of calculation of the different variables from the recorded signals from a weaning-failure patient during the end of the spontaneous breathing trial. The dotted vertical lines represent the start and end of neural inspiration. and the solid vertical lines represent the start and end of inspiratory flow. The neural expiratory phase was divided into four quartiles (Q1-Q4, dotted and dashed lines) to calculate the mean signal per quartile for diaphragm electrical activity (EAdi), transdiaphragmatic pressure (Pdi), and gastric pressure (Pga). Different areas under the curves are shown. The gray area in the flow signal represents tidal volume. The area with three shades of gray in the esophageal pressure (Pes) signal represents the different components of the esophageal pressure-time product during inspiration (PTPesinger): intrinsic positive end-expiratory pressure (PEEPi: light gray), elastic (medium gray), and resistive (dark gray). The components of PTPes_{insp} are defined by the recoil pressures of the chest wall and the lung. The gray area in the Pga signal represents the gastric pressure time product during expiration (PTPesexp).

artefact-free epoch was selected. In figure 1, the acquired signals (flow, diaphragm electrical activity, esophageal pressure, gastric pressure, and transdiaphragmatic pressure) are shown with the calculated variables.

Mechanical inspiratory time, expiratory time, total breath cycle time, and respiratory rate were derived from the flow signal. Tidal volume (V_T) was calculated as the integral of inspiratory flow. Neural inspiration was defined as the period between the onset of diaphragm electrical activity and the peak of diaphragm electrical activity, and neural expiration was defined as the period between peak of diaphragm electrical activity and the onset of the next diaphragm electrical activity.^{26,32}

The rise in expiratory gastric pressure (ΔPga_{exp}) was calculated as the difference in gastric pressure between endinspiration and the start of decrease in esophageal pressure (*i.e.*, start of inspiration). As described previously,^{26,32} expiratory diaphragm electrical activity ($\Delta EAdi_{exp}$) was calculated as the mean signal during ongoing neural expiration divided into four equally sized quartiles (Q1EAdi_{exp}, Q2EAdi_{exp}, Q3EAdi_{exp}, Q4EAdi_{exp}). The mean Δ EAdi_{exp} of each quartile was expressed as the percentage of the peak diaphragm electrical activity of that breath during inspiration after the subtraction of the noise level. Quartile 4 was considered the index closest to the tonic diaphragmatic activity. Details on the noise level calculation are provided in the Supplemental Digital Content (http://links.lww.com/ALN/B717). Analogous analysis was performed on the transdiaphragmatic pressure and gastric pressure signal, resulting in Q1Pdieve, Q2Pdi_{exp}, Q3Pdi_{exp}, Q4Pdi_{exp}, Q1Pga_{exp}, Q2Pga_{exp}, Q3Pga_{exp}, Q3Pga_{exp}, and Q4Pga_{exp}.

During inspiration, the drop in esophageal pressure $(\Delta \text{Pes}_{\text{insp}})$ was calculated as the difference between the start of decrease in esophageal pressure and the negative peak value of esophageal pressure. ΔPga_{insp} was calculated as the difference between the start of increase in gastric pressure and the positive (or negative) peak value of gastric pressure during inspiration. ΔPdi_{insp} was calculated as the difference between the start of increase in transdiaphragmatic pressure and the positive peak value of transdiaphragmatic pressure during inspiration. ΔPes_{insp} and ΔPga_{insp} were corrected for ΔPga_{exp} in the preceding breath. The physiologic base for these corrections have been described in detail.^{22,43,44} In short, expiratory muscle activity can increase the end-expiratory alveolar pressure, leading to an overestimation of the decrease in esophageal pressure during inspiration. In the case of expiratory muscle recruitment, part of the decrease in esophageal pressure preceding inspiration is actually due to relaxation of the expiratory muscles rather than contraction of the inspiratory muscles. Therefore, the amount of pressure due to expiratory muscle activity should be subtracted from the decrease in esophageal pressure. Intrinsic PEEP was calculated as the rise in transdiaphragmatic pressure until the start of inspiratory flow and thus corrected for a drop in gastric pressure at the start of inspiration. $\Delta EAdi_{insp}$, an estimation of neural respiratory drive, was calculated as the peak root mean square per 50 samples of the diaphragm electrical activity during inspiration. Neuromechanical efficiency of diaphragm during inspiration (NMEdi_{insp}) was computed as $\Delta Pdi_{insp}/\Delta EAdi_{insp}$.

The effort of the inspiratory muscles was quantified by calculating the esophageal pressure-time product (PTPes_{insp}).

492

PTPes_{insp} was calculated as the time integral of the difference between esophageal pressure and the recoil pressure of the chest wall, as described previously.^{45,46} PTPes_{insp} was partitioned in resistive, elastic, and intrinsic PEEP components. A calculated theoretical value was used for the recoil pressure of the chest wall, as described previously.⁴⁷ Effort of the expiratory muscles was quantified by calculating the gastric pressure–time product (PTPga_{exp}) during expiration. PTPga_{exp} was calculated as the time integral of the rise in gastric pressure–time product for the respiratory muscles was calculated as PTPes_{insp} + PTPga_{exp}.

lated as PTPes_{insp} + PTPga_{exp}. Dynamic compliance (C_{dyn}) and airway resistance of the lung and airways (R_{aw}) were calculated as described previously.⁴⁸ In short, C_{dyn} was measured as the ratio of the V_T to the change in transpulmonary pressure (difference in pressure between the esophagus and the mouthpiece) between instants of zero air flow. For calculation of R_{aw} , points were selected during the inspiratory phase and expiratory phase when lung volumes were identical and flow rates were about maximal. The ratio of the change in transpulmonary pressure between these points to the corresponding change in flow between these points represents an "average" flow resistance for inspiration and expiration.

Statistical Analysis

Statistical analysis was performed using IBM SPSS Statistics version 22 (IBM Corp., USA), and the data were visualized using GraphPad Prism version 5 (GraphPad Software, Inc., USA). In response to peer review, we adapted the statistical analysis of our data by replacing a linear mixed model design with a two-way repeated measures ANOVA. Assumption of normality was tested using the Shapiro-Wilk normality test. Note that because of the limited sample size per group, this test may lack power to detect deviation from normality. First, to analyze the effects of removing ventilator assist, a paired Student's t test (pressure support ventilation vs. first minute of the spontaneous breathing trial) or its nonparametric equivalent, the Wilcoxon signed-rank test, was performed for the different variables per group (failure and success). Second, to analyze the effect of time and group on the different variables during the spontaneous breathing trial, a two-way repeated measures ANOVA with time as withinsubjects factor (seven time points from minute one to the last minute) and group as between-subjects factor (success or failure) was performed. Third, to analyze respiratory muscle activity during the expiratory phase a two-way repeated measures ANOVA was performed with time as withinsubjects factor (peak and quartile 1 to quartile) and group as between-subjects factor (success or failure). A fourth-order robust polynomial fit was calculated to visualize the trend of respiratory muscle activity during the expiratory phase. Mauchly's test of sphericity was used to test the homogeneity of variance for the mixed ANOVAs. Where Mauchly's test of sphericity was significant ($P \le 0.05$), Greenhouse–Geisser

corrections were applied. In the presence of a significant interaction or between-subject factor, *post hoc* pairwise comparisons between groups at each time point were performed by paired Student's *t* tests with Bonferroni correction. The correlation between the peak expiratory flow and ΔPga_{exp} for the average values per time point during the spontaneous breathing trial was calculated using a Pearson correlation for both weaning groups. The resulting Pearson correlation coefficients were transformed by performing the Fisher's r to z transformation and subsequently compared by determining the observed z test statistic. For all tests, a two-tailed $P \leq 0.05$ was considered significant. The data are described as means \pm SD.

Results

Nine patients met the criteria for weaning failure, five patients failed the spontaneous breathing trial, and four were reintubated within 48 h after extubation. Eleven patients completed the 60-min spontaneous breathing trial and remained extubated for at least 48 h. Patient characteristics and ventilator settings at study inclusion are presented in table 1. Diaphragm electrical activity and transdiaphragmatic pressure signals of one patient in the weaning-success group were excluded from analysis due to dislocation of the catheter during the spontaneous breathing trial. From another patient in the failure group, diaphragm electrical activity was excluded from analysis due to electrode artefacts detected during offline signal analysis.

Ventilation, Respiratory Timing, and Lung Mechanics

Variables of ventilation, respiratory time, and lung mechanics are listed in table 2. Immediately after the transition from pressure support ventilation to spontaneous breathing trial, V_T decreased in both groups but thereafter remained stable for the rest of the spontaneous breathing trial. In both groups, respiratory frequency and the ratio between respiratory frequency and V_T (*i.e.*, an index of rapid shallow breathing) increased after the transition to the spontaneous breathing trial. Only in the weaning-failure group, inspiratory time significantly decreased, and rapid shallow breathing index increased (borderline significant) during the spontaneous breathing trial. V_E did not change significantly during the spontaneous breathing trial and was not different between groups.

Respiratory Muscle Activity during Expiration

In figure 2 (A–F), activity of the diaphragm (Δ EAdi_{exp} and Δ Pdi_{exp}) during the expiration (divided into four equally sized quartiles) together with peak activity during inspiration is shown during pressure support ventilation and during the first and last minute of the spontaneous breathing trial. Changes in expiratory gastric pressure are shown in figure 2 (G–I), together with peak activity of gastric pressure during expiration. At the end of the spontaneous breathing trial, expiratory gastric pressure increased in the failure group

493

Copyright © 2018, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. Unauthorized reproduction of this article is prohibited.

Patient	No. and Diagnosis	Age (yr)	BMI (kg/m²)	Sex	Days on Ventilator	PSV Level (cm H ₂ O)	NAVA Level (cm $H_2O/\mu V$)	PEEP Level (cm H ₂ O)	FI02	Pao ₂ /Fio ₂ (mmHg)	Time to Failure (min)
Failure	aroup (n = 9)										
-	Cardiac arrest	88	24	ш	24	ı	2.3	9	0.40	268	60 [*]
0	Pneumonia	64	22	ш	49	10		9	0.30	328	52
Ю	Postoperative, CABG	72	27	ш	83	10		ø	0.40	194	60 [*]
4	Multitrauma	45	28	Σ	4	10		5	0.30	318	24
ß	Exacerbation COPD	58	25	Σ	14	80		10	0.30	405	60 [*]
9	Postoperative, CABG	67	29	ш	œ	0		ø	0.40	214	16
7	Postoperative, AAA	63	27	Σ	7	4		8	0.40	199	60 [*]
80	Postoperative, papillary muscle rupture	80	24	Σ	œ	0		ø	0.45	132	13
6	Cardiogenic shock	76	23	Σ	21	ı	0.4	10	0.30	228	23
Succes	ss group (n = 11)										
-	Tracheal stenosis, COPD	69	25	Σ	7	14		က	0.45	190	
0	Postoperative, aortoenteric fistula, COPD	71	26	Σ	80	5		9	0.40	÷	
ო	Sepsis	58	26	ш	7	0		9	0.35	330	
4	Postoperative, CABG	73	29	Σ	14	10		10	ı	÷	
5	Cardiac arrest	76	29	ш	17	9		9	0.45	248	
9	Postoperative, AAA	82	28	Σ	5	0		5	0.45	258	
7	Sepsis	73	29	Σ	24	80		9	0.25	306	
ø	Multitrauma	46	25	Σ	6	0		5	0.30	÷	
o	Sepsis	63	24	ш	7	0		8	0.30	+	
10	Pneumonia	60	21	Σ	12	9		9	0.35	291	
:	Postoperative, CABG	78	30	Σ	က	4		10	0.40	358	
*Reintuk	ation within 48h. †Patient without arterial line.										
AAA = a	bdominal aortic aneurvsm: BMI = bodv mass index: (CABG = cc	pronary artery	bvpass ar	aft: COPD = chr	onic obstructive i	oulmonary disease:	F = female: FIO ₂ =	fractional	inspired oxvaen	tension: M = male:
NAVA =	neurally adjusted ventilatory assist; PEEP = positive (end-expira	tory pressure;	PSV = pre	ssure support v	entilation.		2	2		

Table 1. Patient Characteristics

Anesthesiology 2018; 129:490-501

494 Copyright © 2018, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. Unauthorized reproduction of this article is prohibited.

Respiratory Muscle Effort during Weaning

Table 2.Variables of Ventilation, Respiratory Timing, and Mechanics during PSV and the SBT for the Failure Group and SuccessGroup

						P Val	ue
					Main	Effect	to to see at the se
	Group	PSV	SBT Start	SBT End	Time	Group	(Time × Group)
V _T (ml)	F S	535±153 [*] 486±233 [*]	384±183 414±205	401±214 413±171	0.901	0.830	0.670
Peak inspiratory flow (ml/s)	F S	766±232 [*] 714±246 [*]	647±213 637±270	801±356 650±257	0.017	0.399	0.055
Peak expiratory flow (ml/s)	F S	-618±132 -586±193 [*]	-556±151 -525±216	-651±244 -547±194	0.079	0.362	0.417
Frequency (breaths/min)	F S	22.1±8.8 [*] 24.9±8.2 [*]	26.9±8.2 27.6±9.1	30.2 ± 8.4 28.0 ± 9.4	0.128	0.743	0.171
Minute ventilation (I/min)	F S	11.3±4.0 11.5±4.8	10.0±4.5 11.1±5.2	11.6±6.4 11.1±4.6	0.247	0.985	0.385
Inspiratory time (s)	F S	$1.01 \pm 0.33^{*}$ 0.95 ± 0.22	0.87 ± 0.26 0.96 ± 0.28	0.72 ± 0.20 0.96 ± 0.32	0.024	0.126	0.033
Expiratory time (s)	F S	2.20±1.16 [*] 1.76±0.81 [*]	1.70 ± 0.87 1.48 ± 0.64	1.52±0.60 1.53±0.16	0.414	0.747	0.375
Inspiratory time/breath cycle time	F S	0.29 ± 0.12 $0.37 \pm 0.06^{*}$	0.35 ± 0.08 0.41 ± 0.07	0.33 ± 0.06 0.40 ± 0.06	0.202	0.048	0.466
Frequency/V _T (breaths \cdot min ⁻¹ \cdot ⁻¹)	F S	46±28 [*] 61±29 [*]	83±44 78±34	98±55 78±33	0.182	0.437	0.057
PEEPi (cm H ₂ O)	F S	2.7±2.7 [*] 2.0±1.5	4.9 ± 3.6 2.8 ± 2.6	5.3 ± 4.5 3.4 ± 3.0	0.621	0.160	0.649
C _{dyn} (ml/cm H ₂ O)	F	$39.5 \pm 19.4^{*}$ 35.1 ± 12.6	31.9 ± 16.8 40.8 ± 28.6	29.4 ± 24.2 35.9 ± 23.5	0.464	0.529	0.695
$R_{aw} (cm H_2O \cdot I^{-1} \cdot s^{-1})$	F S	9.1±7.4 [*] 7.3±3.6 [*]	16.6±8.6 14.4±7.9	17.2 ± 8.3 15.9 ± 10.3	0.833	0.682	0.501

Note that only data at start and end of the SBT are shown and not the time intervals in between. The P values in table reflect results during the SBT (mixed ANOVA).

* $P \le 0.05 \text{ PSV}$ versus SBT start.

 C_{dyn} = dynamic compliance; F = weaning-failure group; PEEPi = intrinsic positive end-expiratory pressure; PSV = pressure support ventilation; R_{aw} = airway resistance; S = weaning-success group; SBT = spontaneous breathing trial; V_T = tidal volume.

compared to the success group. A significant correlation was found between peak expiratory flow and ΔPga_{exp} during the spontaneous breathing trial in the weaning-failure group (R² = 0.90, *P* = 0.001) but not in the weaning-success group (R² = 0.10, *P* = 0.486). The correlation coefficients were significantly different (*P* = 0.039).

Respiratory Muscle Activity during Inspiration

Figure 3 shows respiratory muscle activity during inspiration under pressure support and during the spontaneous breathing trial. Removal of ventilator assist caused immediate increases in ΔPdi_{insp} , ΔPes_{insp} and $\Delta EAdi_{insp}$ in both groups, whereas NMEdi_{insp} did not change in the first minute after the removal of inspiratory support.

During the course of the spontaneous breathing trial, no significant interactions were found between time and group for ΔPdi_{insp} , ΔPes_{insp} , $\Delta EAdi_{insp}$, and $NMEdi_{insp}$. However, $\Delta EAdi_{insp}$ was significantly higher in the failure group compared to the success group during the entire spontaneous breathing trial. Consequently, NMEdi_{insp} was lower in the weaning-failure group compared to the success group (fig. 3E). In the failure group, intrinsic PEEP increased after removal of

ventilator assist, but during the spontaneous breathing trial, there were no differences between both groups (table 2).

Pressure–Time Product of the Respiratory Muscles

Figure 4 shows the pressure-time products of the expiratory muscle and inspiratory muscles during pressure support ventilation and the first and last minute of the spontaneous breathing trial. The transition from pressure support ventilation to spontaneous breathing did not affect PTPga_{exp} but increased in PTPes_{insp} in both groups. During the spontaneous breathing trial, there were no differences in PTPes,insp between weaning-failure and weaning-success patients. In addition, there were no differences between the intrinsic PEEP and the elastic and resistive components of $PTPes_{insp}$ between groups. The apparent increase in PTPgaern in the failure group did not reach statistical significance (P = 0.099). However, the contribution of the expiratory muscles to the total pressure-time product significantly increased from $13\pm9\%$ at the start of the spontaneous breathing trial to $24 \pm 10\%$ at the end of the spontaneous breathing trial in the weaning-failure group but did not change in the weaning-success group (fig. 5).



Fig. 2. Diaphragm electrical activity (EAdi), transdiaphragmatic pressure (Pdi), and gastric pressure (Pga) during expiration. Mean expiratory diaphragm electrical activity (Δ EAdi_{exp}) values were normalized to the individual EAdi peak during inspiration (insp). Expiratory (exp) peak values of Pga were plotted between quartile (Q) 3 and Q4, the most common position of the peak during expiration (see fig. 1). There is no expiratory activity of the diaphragm in Q4 during pressure support ventilation (PSV) or during the spontaneous breathing trial (SBT). There is a significant increase in expiratory gastric pressure (Δ Pga_{exp}) at the end of the SBT in the failure group compared to the success group. The *circles* in the figure represent the calculated mean values per quartile and the peak values (*white circles* indicate success, *black circles* indicate failure), whereas the *dashed curves* show the fourth order polynomial fit. SD ranged between 1.9 and 11.7%, between 0.6 and 8.8 cm H₂O and between 0.4 and 3.3 cm H₂O for % Δ EAdi_{exp}, Δ Pdi_{exp}, and Δ Pga_{exp}, respectively. The reported *P* values in each panel represent the interaction term (time × group) of the two-way repeated measures ANOVA.

Arterial Blood Gas Measurements and Hemodynamics

In table 3, the blood gas values and hemodynamic variables are presented. There were significant effects of time and group on $Paco_2$, such that $Paco_2$ increased over time during the spontaneous breathing trial and was higher in the failure group. However, the interaction between the effects of time and group were not significant, indicating that the increase in $Paco_2$ was not different between both groups.

Discussion

The present study provides new insights in the recruitment and effort of the respiratory muscles during expiration in a heterogeneous group of patients weaning from mechanical ventilation. First, we found that in weaning-failure patients, expiratory muscles recruitment is as high as 24% of the total respiratory muscle effort at the end of the trial. Second, we found no difference in tonic activity of the diaphragm during expiration between weaning-failure patients and weaning-success patients. Finally, neuromechanical efficiency of the diaphragm is lower in weaning-failure patients compared to weaning-success patients, both during pressure support ventilation and during a spontaneous breathing trial.

Respiratory Muscle Effort during Expiration

The pressure-time product has been used to quantify breathing effort.^{45,46} Experimental studies have demonstrated that changes in inspiratory PTPes are correlated with changes in



Fig. 3. Changes in transdiaphragmatic pressure (Δ Pdi_{insp}), esophageal pressure (Δ Pes_{insp}), gastric pressure (Δ Pga_{insp}), diaphragm electrical activity (Δ EAdi_{insp}), and neuromechanical efficiency (NMEdi_{insp}) during inspiration for pressure support ventilation (PSV) and during the spontaneous breathing trial (SBT) in the weaning-success (*white circles*) and weaning-failure (*black circles*) groups. After removal of ventilator assist Δ Pdi_{insp}, Δ Pes_{insp}, and Δ EAdi_{insp} increased in both groups. During the SBT, data were analyzed at seven points in time: the first minute (start SBT) and last minute (end SBT) and five epochs of at least 1 min taken at equal time intervals in between (ticks in between). A *P* value table is inserted with the results of the repeated measures ANOVA for each variable during the SBT. **P* < 0.05 PSV *versus* start SBT.

diaphragmatic energy expenditure.^{49,50} In the present study, we found an increase in PTPes during inspiration with the transition from mechanical ventilation to unassisted breathing, with no differences between the success group and the failure group. At first sight, this seems surprising, but under conditions of increased loading and/or impaired diaphragm function, the expiratory muscles could be recruited.²³ Indeed, in the present study, expiratory muscle activity, as indicated by an expiratory rise in gastric pressure, was found in the weaning-failure group during the spontaneous breathing trial. This expiratory rise in gastric pressure is consistent with previous findings in patients weaning from mechanical ventilation with COPD.^{21,22} Expiratory muscle recruitment may increase the energy expenditure of respiration. In exercising healthy subjects, expiratory pressure generation is a significant

contributor to the perception of dyspnea.⁵¹ Therefore, muscle effort during expiration should be taken into account when calculating respiratory muscle effort. To this end, we calculated the expiratory pressure–time product from the gastric pressure curve. In the weaning-failure group, the contribution of the expiratory muscles to total respiratory muscle effort progressively increased during the spontaneous breathing trial, whereas it remained stable in successfully extubated patients. Whether the increase in expiratory muscles, expiratory rib cage muscles, or a combination of the two is unknown; the relative contribution of each muscle group during expiration cannot be derived from the expiratory rise in gastric pressure.²²

Pellegrini *et al.*³² have recently demonstrated that in lunginjured pigs, decreasing extrinsic PEEP increases diaphragm

497





Fig. 4. Expiratory muscle pressure–time product (*A*) and inspiratory muscle pressure–time product (*B*) for pressure support ventilation (PSV) and during the spontaneous breathing trial (SBT) in the weaning-success and weaning-failure groups. After removal of ventilator assist, esophageal pressure–time product during inspiration (PTPes_{insp}) increased in both groups. During the SBT, there were no effects of time and group on PTPes. The apparent increase in gastric pressure–time product during expiration (PTPga_{exp}) did not reach statistical significance for the interaction term of the mixed ANOVA (*P* = 0.099). Note that only data at start and end of the SBT are shown and not the time intervals in between. **P* < 0.05 PSV *versus* start SBT. PEEPi = intrinsic positive endexpiratory pressure.

activity during expiration. Diaphragmatic contraction during expiration will affect ΔPga_{exp} and $PTPga_{exp}$ in patients weaning from mechanical ventilation. In the current study, there was negligible low tonic diaphragmatic activity in the last quartile of the expiratory phase during the spontaneous breathing trial. However, in the first expiratory quartile, diaphragm activity was approximately 40% of its peak value. In our opinion, it is debatable whether the electrical



Fig. 5. The contribution of the expiratory muscles to total respiratory muscle effort for pressure support ventilation (PSV) and during the spontaneous breathing trial (SBT) in the weaning-success (*white circle*) and weaning-failure (*black circle*) groups. During the SBT, there was a significant interaction between the time and group (P = 0.049) on the ratio between the expiratory gastric pressure–time product and the total respiratory pressure–time product (PTPga_{exp}/PTPtot). # $P \le 0.05$ pairwise *post hoc* comparison between success and failure group.

activity of the diaphragm in the first expiratory quartile should be defined as persistent activity of the diaphragm in expiration^{26,32} or should be regarded as the termination of the inspiratory activity. Regardless, in our study there was no difference in activity of the diaphragm (diaphragm electrical activity and transdiaphragmatic pressure) between weaning-failure patients and weaning-success patients for all expiratory quartiles during the spontaneous breathing trial. Therefore, it is reasonable to assume that cyclic expiratory changes in gastric pressure at the end of the spontaneous breathing trial in weaning-failure patients are the result of expiratory muscle activity. In addition, the rise in gastric pressure was correlated with peak expiratory flow in the weaning-failure group, consistent with increased expiratory muscle activity.

It has been reasoned that an important goal of expiratory muscle recruitment is to assist the inspiratory muscles by decreasing end-expiratory lung volume.^{23,52} A decrease in end-expiratory lung volume places the diaphragm at a more optimal position of the length-tension curve. However, intrinsic PEEP (corrected for the expiratory rise in gastric pressure) in the current study, a surrogate of end-expiratory lung volume, was not lower in the failure group compared to the success group. On the contrary, intrinsic PEEP tended to be higher in the failure group. In addition, we found no improvements in NMEdi_{insp} in the failure group as a result of increased expiratory muscle recruitment. In line with these findings, it has been demonstrated that expiratory muscle recruitment does not increase NMEdi, in COPD patients.⁵² Thus, it is questionable whether expiratory muscle recruitment assists the inspiratory muscles in weaningfailure patients. Whether expiratory muscle recruitment aids weaning patients should be addressed in future studies.

					P Value			
				Main	Effect			
	Group	PSV [∗]	SBT End	Time	Group	(Time × Group)		
Pao ₂ (mmHg)	F S	91±26 102±21	101±18 99±26	0.225	0.127	0.595		
Paco ₂ (mmHg)	F S	46±11 38±4	50 ± 14 40 ± 5	0.014	0.045	0.128		
рН	F S	7.42 ± 0.07 7.45 ± 0.04	7.39 ± 0.08 7.44 ± 0.04	0.015	0.261	0.427		
MAP (mmHg)	F S	88 ± 20 96 ± 25	92 ± 23 94 ± 21	0.621	0.707	0.157		
Heart rate	F S	87±17 81±11	91±21 82±9	0.194	0.312	0.425		

Table 3. Blood Gas Values and Hemodynamic Variables before and after the SBT for the Failure Group and the Success Group

*Blood gas values were drawn just before start of the SBT.

F = weaning failure; MAP = mean arterial pressure; PSV = pressure support ventilation; S = weaning success; SBT = spontaneous breathing trial.

Neuromechanical Efficiency of the Diaphragm during Inspiration

As expected, removal of ventilator assist increased ΔPdi_{insp} and $\Delta EAdi_{insp}$ in both groups. NMEdi_{insp} was lower in weaning-failure patients compared to the success group. The neuromechanical efficiency of the diaphragm in weaning-failure patients and weaning-success patients in our study were approximately 30 and 50% of the values we reported for healthy subjects, respectively.³⁷ To the best of our knowledge, our study is the first to continuously measure NMEdi_{insp} during a T-tube trial using diaphragm electrical activity and transdiaphragmatic pressure. Previously, NMEdi, nsp has been calculated intermittently during a 30-min weaning trial as the decrease in airway pressure divided by $\Delta EAdi_{insp}$ during an inspiratory occlusion.¹⁸ In the latter study, the weaning trial was performed with 5 cm H₂O of continuous positive airway pressure. The addition of 5 cm H₂O of continuous positive airway pressure decreases the work of breathing by as much as 40% in ventilated patients.⁴⁶ Nevertheless, the reduced ability of the diaphragm to convert neural respiratory drive into pressure in weaning-failure patients is a consistent finding among studies.17,18

Numerous studies have demonstrated that diaphragm weakness develops in critically ill patients.^{9–14} Reduced pressure-generating capacity of the diaphragm may result from structural modifications but also from the development of muscle fatigue, altered force-velocity relation, or altered force-length relation of the muscle. These factors may play a role in the development of weakness during a spontaneous breathing trial. However, previously it was demonstrated that a 1-h spontaneous breathing trial did not cause long-lasting fatigue of the diaphragm ¹⁹ Increased end-expiratory lung volume may place the diaphragm at a less favorable position on a length-tension curve. An increase in lung volume from functional residual capacity to total lung capacity reduces transdiaphragmatic pressure by 60% for a given $\Delta EAdi_{insp}$.⁵³ Although we did not directly measure end-expiratory lung volume, corrected

intrinsic PEEP and lung mechanics were not significantly different between groups during weaning, suggesting no major differences in lung volume. Accordingly, reduced neuromechanical efficiency in the failure group may largely be explained by structural modifications, such as diaphragm atrophy, but we cannot completely rule out impaired respiratory mechanics.

In contrast to a previous study by Jubran and Tobin,⁴⁵ we did not find a difference in PTPes_{insp} between weaningsuccess and weaning-failure patients. This difference may be explained by patient selection and weaning failure criteria. In Jubran and Tobin's study, only patients with COPD were included. This provides an explanation for the fact that the higher PTPes_{insp} resulted from higher resistance and intrinsic PEEP. Furthermore, we defined weaning failure as failing the spontaneous breathing trial or reintubation within 48 h after extubation, whereas in their study, weaning failure was defined as failing the spontaneous breathing trial.

In the present study, increased respiratory muscle effort (including expiratory muscle effort) and reduced neuromechanical efficiency of the diaphragm appear to play a role in failure to wean from mechanical ventilation. However, it is important to note that cardiac dysfunction, cognitive dysfunction, and metabolic disorders are recognized causes for weaning failure as well^{6,7} and may have contributed to weaning failure in the current study.

Limitations

In the present study, raw EAdi signals were recorded and processed according to the methods of Sinderby *et al.*⁴⁰⁻⁴² Today, many studies record electrical activity of the diaphragm not in its raw format but as a processed signal *via* the SERVO-i (Maquet Critical Care) ventilator. Differences between algorithms to process EAdi can lead to differences in absolute values of Δ EAdi_{insp}. Therefore, absolute values of Δ EAdi_{insp} (and NMEdi_{insp}) obtained in our study cannot be directly compared to values obtained with recordings from the SERVO-i (Maquet Critical Care) ventilator.

We did not record electrical activity of the abdominal wall muscles. However, we are confident that ΔPga_{exp} reflects expiratory muscle recruitment in our study. First, we excluded the possibility of expiratory diaphragmatic activity contributing to ΔPga_{exp} . Second, we found a strong correlation between peak expiratory flow and ΔPga_{exp} in weaningfailure patients. Third, a strong correlation has previously been demonstrated between ΔPga_{exp} and electrical activity of the abdominal wall muscles.⁵⁴

In conclusion, the expiratory muscles significantly contribute to respiratory muscle effort in a mixed group of weaning-failure patients. Therefore, the expiratory pressure– time product should be measured when estimating energy expenditure of the respiratory muscles during weaning from mechanical ventilation. We did not find evidence of increased expiratory tonic diaphragmatic contraction during weaning failure. In addition, our findings confirm that impaired pressure-generating capacity of the diaphragm, regardless of its origin, plays a role in failure to wean from mechanical ventilation.

Acknowledgments

The authors thank Wiebe Pestman, Ph.D. (Department of Biostatistics, Radboud University Medical Center, Nijmegen, The Netherlands), for his statistical advice and review of the manuscript.

Research Support

This study was investigator-initiated and financed by institutional resources (Department of Critical Care Medicine, Radboud University Medical Center, Nijmegen, The Netherlands).

Competing Interests

The authors declare no competing interests.

Correspondence

Address correspondence to Dr. Heunks: VU University Medical Center Amsterdam, Department of Intensive Care Medicine, Postbox 7057, 1007 MB Amsterdam, The Netherlands. L.Heunks@VUmc.nl. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY'S articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

- Béduneau G, Pham T, Schortgen F, Piquilloud L, Zogheib E, Jonas M, Grelon F, Runge I, Nicolas Terzi, Grangé S, Barberet G, Guitard PG, Frat JP, Constan A, Chretien JM, Mancebo J, Mercat A, Richard JM, Brochard L; WIND (Weaning according to a New Definition) Study Group and the REVA (Réseau Européen de Recherche en Ventilation Artificielle) Network: Epidemiology of weaning outcome according to a new definition: The WIND study. Am J Respir Crit Care Med 2017; 195:772–83
- Peñuelas O, Frutos-Vivar F, Fernández C, Anzueto A, Epstein SK, Apezteguía C, González M, Nin N, Raymondos K, Tomicic V, Desmery P, Arabi Y, Pelosi P, Kuiper M, Jibaja M, Matamis

D, Ferguson ND, Esteban A; Ventila Group: Characteristics and outcomes of ventilated patients according to time to liberation from mechanical ventilation. Am J Respir Crit Care Med 2011; 184:430–7

- Funk GC, Anders S, Breyer MK, Burghuber OC, Edelmann G, Heindl W, Hinterholzer G, Kohansal R, Schuster R, Schwarzmaier-D'Assie A, Valentin A, Hartl S: Incidence and outcome of weaning from mechanical ventilation according to new categories. Eur Respir J 2010; 35:88–94
- Boles JM, Bion J, Connors A, Herridge M, Marsh B, Melot C, Pearl R, Silverman H, Stanchina M, Vieillard-Baron A, Welte T: Weaning from mechanical ventilation. Eur Respir J 2007; 29:1033–56
- 5. Dres M, Goligher EC, Heunks LMA, Brochard LJ: Critical illness-associated diaphragm weakness. Intensive Care Med 2017; 43:1441–52
- Doorduin J, van der Hoeven JG, Heunks LM: The differential diagnosis for failure to wean from mechanical ventilation. Curr Opin Anaesthesiol 2016; 29:150–7
- 7. Heunks LM, van der Hoeven JG: The ABC of weaning failure: A structured approach. Crit Care 2010; 14:245
- Goligher EC, Fan E, Herridge MS, Murray A, Vorona S, Brace D, Rittayamai N, Lanys A, Tomlinson G, Singh JM, Bolz SS, Rubenfeld GD, Kavanagh BP, Brochard LJ, Ferguson ND: Evolution of diaphragm thickness during mechanical ventilation. Impact of inspiratory effort. Am J Respir Crit Care Med 2015; 192:1080–8
- Hooijman PE, Beishuizen A, Witt CC, de Waard MC, Girbes AR, Spoelstra-de Man AM, Niessen HW, Manders E, van Hees HW, van den Brom CE, Silderhuis V, Lawlor MW, Labeit S, Stienen GJ, Hartemink KJ, Paul MA, Heunks LM, Ottenheijm CA: Diaphragm muscle fiber weakness and ubiquitin-proteasome activation in critically ill patients. Am J Respir Crit Care Med 2015; 191:1126–38
- 10. Hooijman PE, Beishuizen A, de Waard MC, de Man FS, Vermeijden JW, Steenvoorde P, Bouwman RA, Lommen W, van Hees HW, Heunks LM, Dickhoff C, van der Peet DL, Girbes AR, Jasper JR, Malik FI, Stienen GJ, Hartemink KJ, Paul MA, Ottenheijm CA: Diaphragm fiber strength is reduced in critically ill patients and restored by a troponin activator. Am J Respir Crit Care Med 2014; 189:863–5
- 11. Demoule A, Jung B, Prodanovic H, Molinari N, Chanques G, Coirault C, Matecki S, Duguet A, Similowski T, Jaber S: Diaphragm dysfunction on admission to the intensive care unit: Prevalence, risk factors, and prognostic impact: A prospective study. Am J Respir Crit Care Med 2013; 188:213–9
- 12. Jaber S, Petrof BJ, Jung B, Chanques G, Berthet JP, Rabuel C, Bouyabrine H, Courouble P, Koechlin-Ramonatxo C, Sebbane M, Similowski T, Scheuermann V, Mebazaa A, Capdevila X, Mornet D, Mercier J, Lacampagne A, Philips A, Matecki S: Rapidly progressive diaphragmatic weakness and injury during mechanical ventilation in humans. Am J Respir Crit Care Med 2011; 183:364–71
- 13. Hermans G, Agten A, Testelmans D, Decramer M, Gayan-Ramirez G: Increased duration of mechanical ventilation is associated with decreased diaphragmatic force: A prospective observational study. Crit Care 2010; 14:R127
- 14. Levine S, Nguyen T, Taylor N, Friscia ME, Budak MT, Rothenberg P, Zhu J, Sachdeva R, Sonnad S, Kaiser LR, Rubinstein NA, Powers SK, Shrager JB: Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. N Engl J Med 2008; 358:1327–35
- 15. De Jonghe B, Bastuji-Garin S, Durand MC, Malissin I, Rodrigues P, Cerf C, Outin H, Sharshar T; Groupe de Réflexion et d'Etude des Neuromyopathies en Réanimation: Respiratory weakness is associated with limb weakness and delayed weaning in critical illness. Crit Care Med 2007; 35:2007–15
- DiNino E, Gartman EJ, Sethi JM, McCool FD: Diaphragm ultrasound as a predictor of successful extubation from mechanical ventilation. Thorax 2014; 69:423–7

- Dres M, Schmidt M, Ferre A, Mayaux J, Similowski T, Demoule A: Diaphragm electromyographic activity as a predictor of weaning failure. Intensive Care Med 2012; 38:2017–25
- Liu L, Liu H, Yang Y, Huang Y, Liu S, Beck J, Slutsky AS, Sinderby C, Qiu H: Neuroventilatory efficiency and extubation readiness in critically ill patients. Crit Care 2012; 16:R143
- Laghi F, Cattapan SE, Jubran A, Parthasarathy S, Warshawsky P, Choi YS, Tobin MJ: Is weaning failure caused by low-frequency fatigue of the diaphragm? Am J Respir Crit Care Med 2003; 167:120–7
- Vassilakopoulos T, Zakynthinos S, Roussos Ch: Respiratory muscles and weaning failure. Eur Respir J 1996; 9:2383–400
- 21. Tobin MJ, Laghi F, Brochard L: Role of the respiratory muscles in acute respiratory failure of COPD: Lessons from weaning failure. J Appl Physiol 2009; 107:962–70
- Parthasarathy S, Jubran A, Laghi F, Tobin MJ: Sternomastoid, rib cage, and expiratory muscle activity during weaning failure. J Appl Physiol 2007; 103:140–7
- Laghi F, Tobin MJ: Disorders of the respiratory muscles. Am J Respir Crit Care Med 2003; 168:10–48
- 24. Kyroussis D, Polkey MI, Hamnegård CH, Mills GH, Green M, Moxham J: Respiratory muscle activity in patients with COPD walking to exhaustion with and without pressure support. Eur Respir J 2000; 15:649–55
- Kyroussis D, Mills GH, Polkey MI, Hamnegard CH, Wragg S, Road J, Green M, Moxham J: Effect of maximum ventilation on abdominal muscle relaxation rate. Thorax 1996; 51:510–5
- 26. Emeriaud G, Beck J, Tucci M, Lacroix J, Sinderby C: Diaphragm electrical activity during expiration in mechanically ventilated infants. Pediatr Res 2006; 59:705–10
- 27. Allo JC, Beck JC, Brander L, Brunet F, Slutsky AS, Sinderby CA: Influence of neurally adjusted ventilatory assist and positive end-expiratory pressure on breathing pattern in rabbits with acute lung injury. Crit Care Med 2006; 34:2997–3004
- Easton PA, Katagiri M, Kieser TM, Platt RS: Postinspiratory activity of costal and crural diaphragm. J Appl Physiol 1999; 87:582–9
- Meessen NE, van der Grinten CP, Luijendijk SC, Folgering HT: Continuous negative airway pressure increases tonic activity in diaphragm and intercostal muscles in humans. J Appl Physiol 1994; 77:1256–62
- Muller N, Bryan AC, Zamel N: Tonic inspiratory muscle activity as a cause of hyperinflation in histamine-induced asthma. J Appl Physiol Respir Environ Exerc Physiol 1980; 49:869–74
- Muller N, Volgyesi G, Becker L, Bryan MH, Bryan AC: Diaphragmatic muscle tone. J Appl Physiol Respir Environ Exerc Physiol 1979; 47:279–84
- 32. Pellegrini M, Hedenstierna G, Roneus A, Segelsjö M, Larsson A, Perchiazzi G: The diaphragm acts as a brake during expiration to prevent lung collapse. Am J Respir Crit Care Med 2017; 195:1608–16
- 33. Doorduin J, Sinderby CA, Beck J, van der Hoeven JG, Heunks LM: Assisted ventilation in patients with acute respiratory distress syndrome: Lung-distending pressure and patient-ventilator interaction. ANESTHESIOLOGY 2015; 123:181–90
- 34. Doorduin J, Nollet JL, Vugts MP, Roesthuis LH, Akankan F, van der Hoeven JG, van Hees HW, Heunks LM: Assessment of dead-space ventilation in patients with acute respiratory distress syndrome: A prospective observational study. Crit Care 2016; 20:121
- 35. Doorduin J, Nollet JL, Roesthuis LH, van Hees HW, Brochard LJ, Sinderby CA, van der Hoeven JG, Heunks LM: Partial neuromuscular blockade during partial ventilatory support in sedated patients with high tidal volumes. Am J Respir Crit Care Med 2017; 195:1033–42
- 36. Doorduin J, Sinderby CA, Beck J, van der Hoeven JG, Heunks LM: Automated patient-ventilator interaction analysis during

neurally adjusted non-invasive ventilation and pressure support ventilation in chronic obstructive pulmonary disease. Crit Care 2014; 18:550

- 37. Doorduin J, Sinderby CA, Beck J, Stegeman DF, van Hees HW, van der Hoeven JG, Heunks LM: The calcium sensitizer levosimendan improves human diaphragm function. Am J Respir Crit Care Med 2012; 185:90–5
- Beck J, Sinderby C, Lindström L, Grassino A: Influence of bipolar esophageal electrode positioning on measurements of human crural diaphragm electromyogram. J Appl Physiol 1996; 81:1434–49
- 39. Baydur A, Behrakis PK, Zin WA, Jaeger M, Milic-Emili J: A simple method for assessing the validity of the esophageal balloon technique. Am Rev Respir Dis 1982; 126:788–91
- 40. Sinderby C, Beck J, Spahija J, Weinberg J, Grassino A: Voluntary activation of the human diaphragm in health and disease. J Appl Physiol 1998; 85:2146–58
- Sinderby CA, Beck JC, Lindström LH, Grassino AE: Enhancement of signal quality in esophageal recordings of diaphragm EMG. J Appl Physiol 1997; 82:1370–7
- 42. Sinderby C, Lindström L, Grassino AE: Automatic assessment of electromyogram quality. J Appl Physiol 1995; 79:1803–15
- 43. Vogiatzis I, Georgiadou O, Golemati S, Aliverti A, Kosmas E, Kastanakis E, Geladas N, Koutsoukou A, Nanas S, Zakynthinos S, Roussos C: Patterns of dynamic hyperinflation during exercise and recovery in patients with severe chronic obstructive pulmonary disease. Thorax 2005; 60:723–9
- 44. Lessard MR, Lofaso F, Brochard L: Expiratory muscle activity increases intrinsic positive end-expiratory pressure independently of dynamic hyperinflation in mechanically ventilated patients. Am J Respir Crit Care Med 1995; 151:562–9
- Jubran A, Tobin MJ: Pathophysiologic basis of acute respiratory distress in patients who fail a trial of weaning from mechanical ventilation. Am J Respir Crit Care Med 1997; 155:906–15
- 46. Sassoon CS, Light RW, Lodia R, Sieck GC, Mahutte CK: Pressure-time product during continuous positive airway pressure, pressure support ventilation, and T-piece during weaning from mechanical ventilation. Am Rev Respir Dis 1991; 143:469–75
- 47. Fleury B, Murciano D, Talamo C, Aubier M, Pariente R, Milic-Emili J: Work of breathing in patients with chronic obstructive pulmonary disease in acute respiratory failure. Am Rev Respir Dis 1985; 131:822–7
- Frank NR, Mead J, Ferris BG Jr: The mechanical behavior of the lungs in healthy elderly persons. J Clin Invest 1957; 36:1680–7
- Field S, Sanci S, Grassino A: Respiratory muscle oxygen consumption estimated by the diaphragm pressure-time index. J Appl Physiol Respir Environ Exerc Physiol 1984; 57:44–51
- 50. Rochester DF, Bettini G: Diaphragmatic blood flow and energy expenditure in the dog: Effects of inspiratory airflow resistance and hypercapnia. J Clin Invest 1976; 57:661–72
- Kayser B, Sliwinski P, Yan S, Tobiasz M, Macklem PT: Respiratory effort sensation during exercise with induced expiratory-flow limitation in healthy humans. J Appl Physiol 1997; 83:936–47
- 52. Yan S, Sinderby C, Bielen P, Beck J, Comtois N, Sliwinski P: Expiratory muscle pressure and breathing mechanics in chronic obstructive pulmonary disease. Eur Respir J 2000; 16:684–90
- Beck J, Sinderby C, Lindström L, Grassino A: Effects of lung volume on diaphragm EMG signal strength during voluntary contractions. J Appl Physiol 1998; 85:1123–34
- 54. Parthasarathy S, Jubran A, Tobin MJ: Cycling of inspiratory and expiratory muscle groups with the ventilator in airflow limitation. Am J Respir Crit Care Med 1998; 158:1471–8