

APPROPRIATE USE CRITERIA

ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2016 Appropriate Use Criteria for Coronary Revascularization in Patients With Acute Coronary Syndromes

A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and the Society of Thoracic Surgeons

Coronary Revascularization Writing Group

Manesh R. Patel, MD, FACC, FAHA, FSCAI, *Chair*
 John H. Calhoon, MD
 Gregory J. Dehmer, MD, MACC, MSCAI, FACP, FAHA*
 James Aaron Grantham, MD, FACC
 Thomas M. Maddox, MD, MSc, FACC, FAHA

David J. Maron, MD, FACC, FAHA
 Peter K. Smith, MD, FACC†

*Society for Cardiovascular Angiography and Interventions Representative. †Society of Thoracic Surgeons Representative.

Rating Panel

Michael J. Wolk, MD, MACC, *Moderator*
 Manesh R. Patel, MD, FACC, FAHA, FSCAI, *Writing Group Liaison*
 Gregory J. Dehmer, MD, MACC, MSCAI, FACP, FAHA, *Writing Group Liaison**
 Peter K. Smith, MD, FACC, *Writing Group Liaison*
 James C. Blankenship, MD, MACC‡
 Alfred A. Bove, MD, PhD, MACC‡
 Steven M. Bradley, MD§
 Larry S. Dean, MD, FACC, FSCAI*
 Peter L. Duffy, MD, FACC, FSCAI*
 T. Bruce Ferguson, Jr, MD, FACC‡
 Frederick L. Grover, MD, FACC‡
 Robert A. Guyton, MD, FACC||

Mark A. Hlatky, MD, FACC‡
 Harold L. Lazar, MD, FACC¶
 Vera H. Rigolin, MD, FACC‡
 Geoffrey A. Rose, MD, FACC, FASE#
 Richard J. Shemin, MD, FACC||
 Jacqueline E. Tamis-Holland, MD, FACC‡
 Carl L. Tommaso, MD, FACC, FSCAI*
 L. Samuel Wann, MD, MACC**
 John B. Wong, MD‡

‡American College of Cardiology Representative. §American Heart Association Representative. ||Society of Thoracic Surgeons Representative. ¶American Association for Thoracic Surgery Representative. #American Society of Echocardiography Representative. **American Society of Nuclear Cardiology Representative.

This document was approved by the American College of Cardiology Board of Trustees in October 2016.

The American College of Cardiology requests that this document be cited as follows: Patel MR, Calhoon JH, Dehmer GJ, Grantham JA, Maddox TM, Maron DJ, Smith PK. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2016 appropriate use criteria for coronary revascularization in patients with acute coronary syndromes: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and the Society of Thoracic Surgeons. *J Am Coll Cardiol* 2016;XX:xxx-xx.

This document has been reprinted in *Catheterization and Cardiovascular Interventions* and the *Journal of Nuclear Cardiology*.

Copies: This document is available on the World Wide Web site of the American College of Cardiology (www.acc.org). For copies of this document, please contact Elsevier Reprint Department, fax (212) 633-3820 or e-mail reprints@elsevier.com.

Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the American College of Cardiology. Please contact healthpermissions@elsevier.com.

Appropriate Use Criteria Task Force

John U. Doherty, MD, FACC, *Co-Chair*
 Gregory J. Dehmer, MD, MACC, *Co-Chair*

Steven R. Bailey, MD, FACC, FSCAI, FAHA
 Nicole M. Bhave, MD, FACC
 Alan S. Brown, MD, FACC^{††}
 Stacie L. Daugherty, MD, FACC
 Milind Y. Desai, MBBS, FACC
 Claire S. Duvernoy, MD, FACC
 Linda D. Gillam, MD, FACC
 Robert C. Hendel, MD, FACC, FAHA^{††}
 Christopher M. Kramer, MD, FACC, FAHA^{††}
 Bruce D. Lindsay, MD, FACC^{††}

Warren J. Manning, MD, FACC
 Manesh R. Patel, MD, FACC, FAHA^{§§}
 Ritu Sachdeva, MBBS, FACC
 L. Samuel Wann, MD, MACC^{††}
 David E. Winchester, MD, FACC
 Michael J. Wolk, MD, MACC^{††}
 Joseph M. Allen, MA

^{††}Former Task Force member, current member during the writing effort. ^{†††}Former Task Force Co-Chair, current Co-Chair during the writing effort. ^{§§}Former Task Force Chair, current Chair during the writing effort.

TABLE OF CONTENTS

ABSTRACT	■	Table 1.3 STEMI - Revascularization of Nonculprit Artery During the Initial Hospitalization	■
PREFACE	■	Table 1.4 NSTEMI/Unstable Angina	■
1. INTRODUCTION	■	7. DISCUSSION	■
2. METHODS	■	REFERENCES	■
Indication Development	■	APPENDIX A	
Scope of Indications	■	Appropriate Use Criteria for Coronary Revascularization in Patients With Acute Coronary Syndromes: Participants	■
3. ASSUMPTIONS	■	APPENDIX B	
General Assumptions	■	Relationships With Industry and Other Entities	■
4. DEFINITIONS	■	ABSTRACT	
Indication	■	The American College of Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Thoracic Surgeons, and American Association for Thoracic Surgery, along with key specialty and subspecialty societies, have completed a 2-part revision of the appropriate use criteria (AUC) for coronary revascularization. In prior coronary revascularization AUC documents, indications for revascularization in acute coronary syndromes (ACS) and stable ischemic heart disease were combined into 1 document. To address the expanding clinical indications for coronary revascularization, and in an effort to align the subject matter with the most current American College of Cardiology/American Heart Association guidelines, the new AUC for coronary artery revascularization were separated into 2 documents addressing ACS and stable ischemic heart disease individually. This document presents the AUC for ACS.	
Cardiac Risk Factor Modification and Antianginal Medical Therapy	■	Clinical scenarios were developed to mimic patient presentations encountered in everyday practice and	
Culprit Stenosis	■		
Symptoms of Myocardial Ischemia	■		
Unstable Angina	■		
Stress Testing and Risk of Findings on Noninvasive Testing	■		
The Role of Patient Preference in the AUC	■		
Specific Acute Coronary Syndromes	■		
5. ABBREVIATIONS	■		
6. CORONARY REVASCLARIZATION IN PATIENTS WITH ACS: AUC (BY INDICATION)	■		
Table 1.1 STEMI - Immediate Revascularization by PCI	■		
Table 1.2 STEMI - Initial Treatment by Fibrinolytic Therapy	■		

included information on symptom status, presence of clinical instability or ongoing ischemic symptoms, prior reperfusion therapy, risk level as assessed by noninvasive testing, fractional flow reserve testing, and coronary anatomy. This update provides a reassessment of clinical scenarios that the writing group felt to be affected by significant changes in the medical literature or gaps from prior criteria. The methodology used in this update is similar to the initial document but employs the recent modifications in the methods for developing AUC, most notably, alterations in the nomenclature for appropriate use categorization.

A separate, independent rating panel scored the clinical scenarios on a scale of 1 to 9. Scores of 7 to 9 indicate that revascularization is considered appropriate for the clinical scenario presented. Scores of 1 to 3 indicate that revascularization is considered rarely appropriate for the clinical scenario, whereas scores in the mid-range (4 to 6) indicate that coronary revascularization may be appropriate for the clinical scenario. Seventeen clinical scenarios were developed by a writing committee and scored by the rating panel: 10 were identified as appropriate, 6 as may be appropriate, and 1 as rarely appropriate.

As seen with the prior coronary revascularization AUC, revascularization in clinical scenarios with ST-segment elevation myocardial infarction and non-ST-segment elevation myocardial infarction were considered appropriate. Likewise, clinical scenarios with unstable angina and intermediate- or high-risk features were deemed appropriate. Additionally, the management of nonculprit artery disease and the timing of revascularization are now also rated. The primary objective of the AUC is to provide a framework for the assessment of practice patterns that will hopefully improve physician decision making.

PREFACE

The American College of Cardiology (ACC), in collaboration with the Society for Cardiovascular Angiography and Interventions, Society for Thoracic Surgeons, American Association for Thoracic Surgery, and other societies, developed and published the first version of the appropriate use criteria (AUC) for coronary revascularization in 2009, with the last update in 2012. The AUC are an effort to assist clinicians in the rational use of coronary revascularization in common clinical scenarios found in everyday practice. The new AUC for coronary revascularization was developed as separate documents for acute coronary syndromes (ACS) and stable ischemic heart disease (SIHD). This was done to address the expanding clinical indications for coronary revascularization, include new literature published since

the last update, and align the subject matter with the ACC/American Heart Association guidelines. An additional goal was to address several of the shortcomings of the initial document that became evident as experience with the use of the AUC accumulated in clinical practice.

The publication of AUC reflects 1 of several ongoing efforts by the ACC and its partners to assist clinicians who are caring for patients with cardiovascular diseases and in support of high-quality cardiovascular care. The ACC/American Heart Association clinical practice guidelines provide a foundation for summarizing evidence-based cardiovascular care and, when evidence is lacking, provide expert consensus opinion that is approved in review by the ACC and American Heart Association. However, in many areas, variability remains in the use of cardiovascular procedures, raising questions of over- or under-use. The AUC provide a practical standard upon which to assess and better understand variability.

We are grateful to the writing committee for the development of the overall structure of the document and clinical scenarios and to the rating panel, a professional group with a wide range of skills and insights, for their thoughtful deliberation of the merits of coronary revascularization for various clinical scenarios. We would also like to thank the parent AUC Task Force and the ACC staff, Joseph Allen, Leah White, and specifically Maria Velasquez, for their skilled support in the generation of this document.

Manesh R. Patel, MD, FACC


*Chair, Coronary Revascularization Writing Group
Chair, Appropriate Use Criteria Task Force*

Michael J. Wolk, MD, MACC

Moderator, Appropriate Use Criteria Task Force

1. INTRODUCTION

In a continuing effort to provide information to patients, physicians, and policy makers, the Appropriate Use Task Force approved this revision of the 2012 coronary revascularization AUC (1). Since publication of the 2012 AUC document, new guidelines for ST-segment elevation myocardial infarction (STEMI) (2) and non-ST-segment elevation myocardial infarction (NSTEMI)/unstable angina (3) have been published with additional focused updates of the SIHD guideline and a combined focused update of the percutaneous coronary intervention (PCI) and STEMI guideline (4,5). New clinical trials have been published extending the knowledge and evidence around coronary revascularization, including trials that challenge earlier recommendations about the timing of nonculprit vessel PCI in the setting of STEMI (6-8). Additional studies related to coronary artery bypass graft surgery, medical therapy, and diagnostic technologies such as



fractional flow reserve (FFR) have emerged as well as analyses from The National Cardiovascular Data Registry (NCDR) on the existing AUC that provide insights into practice patterns, clinical scenarios, and patient features not previously addressed (9–11).

In an effort to make the AUC usable, meaningful, and as up-to-date as possible, the writing group was asked to develop AUC specifically for coronary revascularization in ACS including STEMI to coincide with the recently published focused update of the STEMI guidelines (5). A new separate AUC document specific to SIHD is under preparation and will be forthcoming. The goal of the writing group was to develop clinical indications (scenarios) that reflect typical situations encountered in everyday practice, which are then classified by a separate rating panel

using methodology previously described in detail (12) (Figure 1). In addition, step-by-step flow charts are provided to help use the criteria.

2. METHODS

Indication Development

A multidisciplinary writing group consisting of cardiovascular health outcomes researchers, interventional cardiologists, cardiothoracic surgeons, and general cardiologists was convened to review and revise the coronary revascularization AUC.

The revascularization AUC are on the basis of our current understanding of procedure outcomes plus the potential patient benefits and risks of the revascularization

strategies examined. The AUC are developed to identify many of the common clinical scenarios encountered in practice, but cannot possibly include every conceivable patient presentation. (In this document, the phrase “clinical scenario” is frequently used interchangeably with the term “indication.”) Some patients seen in clinical practice are not represented in these AUC or have additional extenuating features that would alter the appropriateness of treatment compared with the exact clinical scenarios presented.

AUC documents often contain more detailed clinical scenarios than the more generalized situations covered in clinical practice guidelines, and thus, subtle differences between these documents may exist. Furthermore, because recommendations for revascularization or the medical management of coronary artery disease (CAD) are found throughout several clinical practice guidelines, the AUC ratings herein are meant to unify related clinical practice guidelines and other data sources and provide a useful tool for clinicians. The AUC were developed with the intent to assist patients and clinicians, but are not intended to diminish the acknowledged complexity or uncertainty of clinical decision-making and should not be a substitute for sound clinical judgment. There are acknowledged evidence gaps in many areas where clinical judgement and experience must be blended with patient preferences, and the existing knowledge base must be defined in clinical practice guidelines.

It is important to emphasize that a rating of *appropriate care* does not mandate that a procedure or revascularization strategy be performed, *may be appropriate care* represents reasonable care and can be considered by the patient and provider, and finally, a rating of *rarely appropriate care* should not prevent a therapy from being performed. It is anticipated that there will be some clinical scenarios rated as rarely appropriate where an alternative therapy or performing revascularization may still be in the best interest of a particular patient. Situations where the clinician believes a therapy contrary to the AUC rating is best for the patient may require careful documentation as to the specific patient features not captured in the clinical scenario or the rationale for the chosen therapy. Depending on the urgency of care, obtaining a second opinion may be helpful in some of these settings.

The AUC can be used in several ways. As a clinical tool, the AUC assist clinicians in evaluating possible therapies under consideration and can help better inform patients about their therapeutic options. As an administrative and research tool, the AUC provide a means to compare utilization patterns across a large subset of providers to deliver an assessment of an individual clinician's management strategies with those of similar physicians. It is important to again emphasize that the AUC should be

used to measure overall patterns of clinical care rather than to adjudicate the appropriateness of individual cases. The ACC and its collaborators believe that an ongoing review of one's practice using these criteria will help guide more effective, efficient, and equitable allocation of healthcare resources, and ultimately lead to better patient outcomes. Under no circumstances should the AUC be used as the sole means to adjudicate or determine payment for individual patients—rather, the intent of the AUC is to provide a framework to evaluate overall clinical practice and to improve the quality of care.

In developing these AUC for coronary revascularization, the rating panel was asked to rate each indication using the following definition of appropriate use:

A coronary revascularization or antianginal therapeutic strategy is appropriate care when the potential benefits, in terms of survival or health outcomes (symptoms, functional status, and/or quality of life) exceed the potential negative consequences of the treatment strategy.

Although antianginal therapy is mentioned in this definition, the writing committee acknowledges that the focus of this document is revascularization, as it is the dominant therapy for patients with ACS. Medical therapy may have a role in the management of ongoing ischemic symptoms, but not to the extent that it does for SIHD.

The rating panel scored each indication on a scale from 1 to 9 as follows:

Score 7 to 9: Appropriate care

Score 4 to 6: May be appropriate care

Score 1 to 3: Rarely appropriate care

Appropriate Use Definition and Ratings

In rating these criteria, the rating panel was asked to assess whether the use of revascularization for each indication is “appropriate care,” “may be appropriate care,” or “rarely appropriate care” using the following definitions and their associated numeric ranges.

Median Score 7 to 9: Appropriate Care

An appropriate option for management of patients in this population due to benefits generally outweighing risks; an effective option for individual care plans, although not always necessary depending on physician judgment and patient-specific preferences (i.e., procedure is generally acceptable and is generally reasonable for the indication).

Median Score 4 to 6: May Be Appropriate Care

At times, an appropriate option for management of patients in this population due to variable evidence or agreement regarding the risk-benefit ratio, potential

benefit on the basis of practice experience in the absence of evidence, and/or variability in the population; effectiveness for individual care must be determined by a patient's physician in consultation with the patient on the basis of additional clinical variables and judgment along with patient preferences (i.e., procedure may be acceptable and may be reasonable for the indication).

Median Score 1 to 3: Rarely Appropriate Care

Rarely an appropriate option for management of patients in this population due to the lack of a clear benefit/risk advantage; rarely, an effective option for individual care plans; exceptions should have documentation of the clinical reasons for proceeding with this care option (i.e., procedure is not generally acceptable and is not generally reasonable for the indication).

Scope of Indications

The indications for coronary revascularization in ACS were developed considering the following common variables:

1. The clinical presentation (STEMI, NSTEMI, or other ACS);
2. Time from onset of symptoms;
3. Presence of other complicating factors (severe heart failure or cardiogenic shock; hemodynamic or electrical instability, presence of left ventricular dysfunction, persistent or recurring ischemic symptoms);
4. Prior treatment by fibrinolysis;
5. Predicted risk as estimated by the Thrombolysis In Myocardial Infarction score;
6. Relevant comorbidities; and
7. Extent of anatomic disease in the culprit and non-culprit arteries.

The writing group characterized ACS and their management into the 2 common clinical presentations: STEMI and NSTEMI/unstable angina. The anatomic construct for CAD is on the basis of the presence or absence of important obstructions in the coronary arteries categorized by the number of vessels involved 1-, 2-, and 3-vessel CAD) and the ability to identify the culprit artery responsible for the ACS. Although the culprit stenosis is frequently obvious from the coronary angiogram, there are situations where the location of the culprit stenosis is uncertain or where multiple culprit stenoses may exist.

After initial treatment of the patient with an ACS, it may be helpful to categorize the amount of myocardium at risk or affected by ischemia; thus, a minority of scenarios include noninvasive testing. The writing group characterized noninvasive test findings as low-risk versus intermediate- or high-risk, as these terms are routinely used in clinical practice. The use of FFR measurement is increasing in the setting of stable ischemic heart disease, but there are limited data on its utility in the setting of

ACS to evaluate nonculprit vessels (6). Nevertheless, the writing group provided some indications with invasive physiology testing (represented by FFR) in nonculprit vessels in patients with ACS.

3. ASSUMPTIONS

General Assumptions

Specific instructions and assumptions used by the rating panel to assist in the rating of clinical scenarios are listed in the following text:

1. Each clinical scenario is intended to provide the key information typically available when a patient presents with an ACS, recognizing that especially in the setting of an STEMI, the need for rapid treatment may prevent a complete evaluation.
2. Although the clinical scenarios should be rated on the basis of the published literature, the writing committee acknowledges that in daily practice, decisions about therapy are required in certain patient populations that are poorly represented in the literature. Therefore, rating panel members were instructed to use their best clinical judgment and experience in assigning ratings to clinical scenarios that have low levels of evidence.
3. In ACS, the percent luminal diameter narrowing of a stenosis may be difficult to assess. Determining the significance of a stenosis includes not only the percent luminal diameter narrowing, but also the angiographic appearance of the stenosis and distal flow pattern. For these clinical scenarios, a **coronary stenosis** in an artery is defined as:
 - **Severe:**
 - a. A $\geq 70\%$ luminal diameter narrowing of an epicardial stenosis made by visual assessment in the "worst view" angiographic projection; or
 - b. A $\geq 50\%$ luminal diameter narrowing of the left main artery made by visual assessment, in the "worst view" angiographic projection.
 - **Intermediate:**
 - c. A $\geq 50\%$ and $< 70\%$ diameter narrowing of an epicardial stenosis made by visual assessment in the "worst view" angiographic projection.
4. For scenarios reflecting later phases of care for patients with ACS (scenarios during hospitalization), assume that patients are receiving guideline-directed medical therapy for secondary prevention of cardiac events unless specifically noted and efforts to control other risk factors have started (13-17).
5. Operators performing percutaneous or surgical revascularization have appropriate clinical training and experience and have satisfactory outcomes as assessed by quality assurance monitoring (18-20).

6. Revascularization by either percutaneous or surgical methods is performed in a manner consistent with established standards of care at centers with quality/volume standards (18-20).
7. No unusual extenuating circumstances exist in the clinical scenarios such as but not limited to do-not-resuscitate status, advanced malignancy, unwillingness to consider revascularization, technical reasons rendering revascularization infeasible, or comorbidities likely to markedly increase procedural risk.
8. Assume that the appropriateness rating applies only to the specific treatment strategy outlined in the scenario and not additional revascularization procedures that may be performed later in the patient's course. Specifically, additional elective revascularization procedures (so called delayed staged procedures) performed after the hospitalization for ACS are evaluated and rated in the forthcoming AUC document on SIHD. For data collection purposes, this will require documenting that the procedure is staged (either PCI or hybrid revascularization with surgery).
9. As with all previously published clinical policies, deviations by the rating panel from prior published documents were driven by new evidence and/or implementation of knowledge that justifies such evolution. However, the reader is advised to pay careful attention to the wording of an indication in the present document and should avoid making comparisons to prior documents.
10. Indication ratings contained herein supersede the ratings of similar indications contained in previous AUC coronary revascularization documents.

4. DEFINITIONS

Definitions of terms used throughout the indication set are listed here. These definitions were provided to and discussed with the rating panel before the rating of indications. The writing group assumed that noninvasive assessments of coronary anatomy (i.e., cardiac computed tomography, cardiac magnetic resonance angiography) provide anatomic information that is potentially similar to X-ray angiography. However, these modalities do not currently provide information on ischemic burden and are not assumed to be present in the clinical scenarios.

Indication

A set of patient-specific conditions defines an "indication," which is used interchangeably with the phrase "clinical scenario."

Cardiac Risk Factor Modification and Antianginal Medical Therapy

The indications assume that patients are receiving guideline-directed medical therapies for their ACS

including antiplatelet and anticoagulant medications, beta-blockers, statins, and other medications as indicated by their clinical condition.

Culprit Stenosis

The phrase "culprit stenosis" is often used interchangeably with "infarct-related artery" to identify the coronary artery stenosis and/or artery responsible for the ACS. In this document, the phrase "culprit stenosis or culprit artery" is preferred, because in the setting of unstable angina there may be a culprit stenosis or culprit artery, but by definition, there is no evidence of a myocardial infarction.

Symptoms of Myocardial Ischemia

For the purposes of the clinical scenarios in this document, the AUC are intended to apply to patients who have the typical underlying pathology of an ACS, not simply an elevated troponin value in the absence of an appropriate clinical syndrome. The symptoms of an ACS may be described as both typical and atypical angina or symptoms felt to represent myocardial ischemia, such as exertional dyspnea, and are captured under the broad term "ischemic symptoms." Although previous AUC had used the Canadian Cardiovascular Society system for anginal classification, the writing group recognized that the broad spectrum of ischemic symptoms may limit patients' functional status in a variety of ways, and capturing the Canadian Cardiovascular Society status in clinical practice may also vary widely. Therefore, the presence or absence of ischemic symptoms are presented without specific scale. Additionally, post-ACS symptoms may persist and/or be easily provoked with minimal activity.

Unstable Angina

The definition of unstable angina is largely on the basis of the clinical presentation. Unstable angina is defined as **typical chest pain** or other ischemic symptoms occurring **at rest** or with **minimal exertion**, and presumed to be related to an acutely active coronary plaque. In **contrast** to **stable** angina, **unstable** angina is often described as **severe** and as a **frank pain**. Moreover, unstable angina may be new in onset or occur in a **crescendo** pattern in a patient with a previous stable pattern of angina. Unstable angina may be associated with new electrocardiographic changes such as **transient ST-segment elevation**, ST-segment **depression**, or T-wave **inversion**, but **may be** present in the **absence** of **electrocardiographic** changes. Several scoring systems exist for determining high-risk patients with ACS (Tables A and B).

Stress Testing and Risk of Findings on Noninvasive Testing

Stress testing and coronary CTA are commonly used for both diagnosis and risk stratification of patients with coronary artery disease or those with suspected ACS.

TABLE A

High-Risk Features for Short-Term Risk of Death or Nonfatal MI in Patients With NSTEMI/UA

At least 1 of the following:

- History—accelerating tempo of anginal symptoms in preceding 48 hours
- Character of pain—prolonged ongoing (>20 minutes) rest pain
- Clinical findings
 - Pulmonary edema, most likely due to ischemia
 - New or worsening MR murmur
 - S₃ or new/worsening rales
 - Hypotension, bradycardia, tachycardia
 - Age >75 years
- ECG
 - Transient ST-segment deviation >0.5 mm
 - Bundle-branch block, new or presumed new
 - Sustained ventricular tachycardia
- Cardiac marker
 - Elevated cardiac TnT, TnI, or CK-MB (e.g., TnT or TnI >0.1 ng per mL)

High-risk features were defined as in the ACS guidelines (21).

CK-MB = creatine kinase, MB isoenzyme; ECG = electrocardiogram; MI = myocardial infarction; MR = mitral regurgitation; NSTEMI = non-ST segment elevation myocardial infarction; TnI = troponin I; TnT = troponin T; UA = unstable angina.

Although often contraindicated in ACS, stress testing may be performed for further risk stratification later during the index hospitalization. Risk stratification by noninvasive testing is defined as (4):

Low-risk stress test findings: associated with a <1% per year cardiac mortality rate.

Intermediate-risk stress test findings: associated with a 1% to 3% per year cardiac mortality rate.

High-risk stress test findings: associated with a >3% per year cardiac mortality rate.

The Role of Patient Preference in the AUC

Patients often make decisions about medical treatments without a complete understanding of their options. Patient participation or shared decision-making describes a collaborative approach where patients are provided evidence-based information on treatment choices and are encouraged to use the information in an informed dialogue with their provider to make decisions that not only use the scientific evidence, but also align with their values, preferences, and lifestyle (23-25). The alternative

TABLE B

Thrombolysis In Myocardial Infarction Risk Score—For Patients With Suspected ACS (22)

Variables (1 point each)

- Age ≥65 years
- ≥3 risk factors (HTN, DM, FH, lipids, smoking)
- Known CAD (stenosis ≥50%)
- Aspirin use in past 7 days
- Severe angina (≥2 episodes within 24 hours)
- ST-segment deviation ≥0.5 mm
- Elevated cardiac markers

Risk of death or ischemic event through 14 days

- Low: 0-2 (<8.3% event rate)
- Intermediate: 3-4 (<19.3% event rate)
- High: 5-7 (41% event rate)

ACS = acute coronary syndrome; CAD = coronary artery disease; DM = diabetes mellitus; FH = family history; HTN = hypertension.

decision paradigm, often referred to as medical paternalism, places decision authority with physicians and gives the patient a more passive role (26).

Shared decision-making respects both the provider's knowledge and the patient's right to be fully informed of all care options with their associated risks and benefits. It also suggests that the healthcare team has educated the patient to the extent the patient desires with regard to the risk and benefits of different treatment options. The patient is given the opportunity to participate in the decision regarding the preferred treatment. Especially regarding primary PCI for STEMI, the need for rapid treatment will often preclude a detailed discussion of the risks and benefits of invasive therapy or other possible treatment decisions. However, patient preferences should be considered when the treatment of a nonculprit stenosis is contemplated later during the hospitalization.

Specific Acute Coronary Syndromes

The writing group developed these clinical scenarios around the common clinical situations in which coronary revascularization is typically considered on the basis of evidence and recommendations from the 2013 STEMI guideline (2) and 2014 NSTEMI/unstable angina guideline (3). Because of 3 recent studies and the 2015 update to the PCI/STEMI guidelines, treatment of nonculprit related arteries at the time of the initial procedure or during the initial hospitalization is also explored (5-8). Previously, treatment of nonculprit stenoses during the initial procedure or during the same hospitalization in the absence of clinical instability or further testing documenting ischemia was assigned a Class III recommendation in guideline documents and is thus considered inappropriate using the original terminology for the AUC. The 3 new randomized studies have challenged this concept, leading to a focused update of the PCI/STEMI guideline and the new Class IIb assignment for treatment of nonculprit stenoses in the setting of primary PCI.

However, the timing of treatment and criteria for nonculprit stenosis treatment varied among these 3 studies as shown in Table C.

In PRAMI (Preventive Angioplasty in Acute Myocardial Infarction Trial), the nonculprit stenosis needed to have a diameter stenosis >50% and be deemed treatable by the operator. There were exclusions to immediate nonculprit PCI, such as left main stenosis, ostial left anterior descending coronary artery and circumflex stenoses, and prior coronary artery bypass graft surgery. Treatment at any time other than during the primary PCI was discouraged. In CvLPRIT (Complete Versus Lesion-Only Primary PCI Trial), the nonculprit stenosis was required to have >70% diameter stenosis in 1 angiographic plane or >50% in 2 planes and in an artery >2 mm suitable for stent implantation. Treatment of the nonculprit stenosis

TABLE C Treatment of Nonculprit Stenoses in the Patient With STEMI

	PRAMI (n = 465)	CvLPRIT (n = 296)	DANAMI3-PRIMULTI (n = 627)
Randomization	After primary PCI	"During" primary PCI	After primary PCI
Lesion criteria	>50% DS	>70% DS or >50% DS in 2 views	>50% DS and FFR <0.80 or >90% DS
Strategy for non-IRA lesions	Immediate—at time of primary PCI	Immediate or staged within index admission	Staged within index admission (average day 2)

CvLPRIT = Complete Versus Lesion-Only Primary PCI Trial; DANAMI3-PRIMULTI = The Third Danish Study of Optimal Acute Treatment of Patients with STEMI: Primary PCI in Multivessel Disease; DS = diameter stenosis; FFR = fractional flow reserve; IRA = infarct-related artery; PCI = percutaneous coronary intervention; PRAMI = Preventive Angioplasty in Acute Myocardial Infarction Trial.

immediately following the primary PCI was encouraged, but could be deferred to later during the same hospitalization. In DANAMI3-PRIMULTI (The Third Danish Study of Optimal Acute Treatment of Patients with STEMI: Primary PCI in Multivessel Disease), nonculprit stenoses were treated if the diameter stenosis was >50% and the FFR <0.80 or if the diameter stenosis alone was >90%. Treatment of the nonculprit stenoses was planned for 2 days after the primary PCI during the index hospitalization. These variations in the criteria for nonculprit stenosis treatment and timing of treatment from these 3 relatively small studies make it challenging to develop clinical scenarios. This is an evolving shift in the treatment paradigm for patients presenting with STEMI that, at present, is incompletely understood. Scenarios were developed to allow the rating panel to evaluate clinical situations that mirror the evidence provided in these new trials.

This AUC only covers clinical scenarios where the culprit artery and additional nonculprit arteries are treated at the time of primary PCI or later during the initial hospitalization. The writing group recognizes there may be circumstances where treatment of a nonculprit artery is deferred beyond the initial hospitalization. That specific circumstance was not studied in the 3 recent trials

of nonculprit stenosis treatment. However, if the characteristics of the patient are such that treatment of nonculprit stenoses are deferred beyond the initial hospitalization, it is assumed the patient is clinically stable. These clinical scenarios will be evaluated in the forthcoming SIHD document.

5. ABBREVIATIONS

ACS = acute coronary syndrome

AUC = appropriate use criteria

CAD = coronary artery disease

FFR = fractional flow reserve

NSTEMI = non-ST-segment elevation myocardial infarction

PCI = percutaneous coronary intervention

SIHD = stable ischemic heart disease

STEMI = ST-segment elevation myocardial infarction

6. CORONARY REVASCULARIZATION IN PATIENTS WITH ACS: AUC (BY INDICATION)

Scenarios 1 to 3 in Table 1.1 specifically address treatment of the culprit stenosis at the time intervals and with the

TABLE 1.1 STEMI—Immediate Revascularization by PCI

Indication	Appropriate Use Score (1-9)
Revascularization of the Presumed Culprit Artery by PCI (Primary PCI)	
1. ■ Less than or equal to 12 hours from onset of symptoms	A (9)
2. ■ Onset of symptoms within the prior 12-24 hours AND ■ Severe HF, persistent ischemic symptoms, or hemodynamic or electrical instability present	A (8)
3. ■ Onset of symptoms within the prior 12-24 hours AND ■ Stable without severe HF, persistent ischemic symptoms, or hemodynamic or electrical instability	M (6)
Successful Treatment of the Culprit Artery by Primary PCI Followed by Immediate Revascularization of 1 or More Nonculprit Arteries During the Same Procedure	
4. ■ Cardiogenic shock persisting after PCI of the presumed culprit artery ■ PCI or CABG of 1 or more additional vessels	A (8)
5. ■ Stable patient immediately following PCI of the presumed culprit artery ■ One or more additional severe stenoses	M (6)
6. ■ Stable patient immediately following PCI of the presumed culprit artery ■ One or more additional intermediate (50%-70%) stenoses	M (4)

The number in parenthesis next to the rating reflects the median score for that indication.

A = appropriate; CABG = coronary artery bypass graft; HF = heart failure; M = may be appropriate; PCI = percutaneous coronary intervention; R = rarely appropriate; STEMI = ST-segment elevation myocardial infarction.

TABLE 1.2 STEMI—Initial Treatment by Fibrinolytic Therapy

Indication	Appropriate Use Score (1-9)
PCI of the Presumed Culprit Artery After Fibrinolysis	
7. ■ Evidence of failed reperfusion after fibrinolysis (e.g., failure of ST-segment resolution, presence of acute severe HF, ongoing myocardial ischemia, or unstable ventricular arrhythmias)	A (9)
8. ■ Stable after fibrinolysis AND ■ Asymptomatic (no HF, myocardial ischemia, or unstable ventricular arrhythmias) AND ■ PCI performed 3-24 hours after fibrinolytic therapy	A (7)
9. ■ Stable after fibrinolysis AND ■ Asymptomatic (no HF, myocardial ischemia, or unstable ventricular arrhythmias) AND ■ PCI >24 hours after onset of STEMI	M (5)

The number in parenthesis next to the rating reflects the median score for that indication.

A = appropriate; CABG = coronary artery bypass graft; HF = heart failure; M = may be appropriate; PCI = percutaneous coronary intervention; R = rarely appropriate; STEMI = ST-segment elevation myocardial infarction.

TABLE 1.3 STEMI—Revascularization of Nonculprit Artery During the Initial Hospitalization

Indication	Appropriate Use Score (1-9)
Successful Treatment of the Culprit Artery by Primary PCI or Fibrinolysis Revascularization of 1 or More Nonculprit Arteries During the Same Hospitalization	
Revascularization by PCI or CABG	
10. ■ Spontaneous or easily provoked symptoms of myocardial ischemia ■ One or more additional severe stenoses	A (8)
11. ■ Asymptomatic ■ Findings of ischemia on noninvasive testing ■ One or more additional severe stenoses	A (7)
12. ■ Asymptomatic (no additional testing performed) ■ One or more additional severe stenoses	M (6)
13. ■ Asymptomatic (no additional testing performed) ■ One or more additional intermediate stenoses	R (3)
14. ■ Asymptomatic ■ One or more additional intermediate (50%-70%) stenoses ■ FFR performed and ≤ 0.80	A (7)

The number in parenthesis next to the rating reflects the median score for that indication.

A = appropriate; CABG = coronary artery bypass graft; FFR = fractional flow reserve; M = may be appropriate; PCI = percutaneous coronary intervention; R = rarely appropriate; STEMI = ST-segment elevation myocardial infarction.

presence or absence of symptoms as noted. Scenarios 4 to 6 in Table 1.1 specifically address treatment of 1 or more nonculprit stenoses during the same procedure as treatment of the culprit stenosis. Because these scenarios are specific for nonculprit treatment immediately following

primary PCI, the criteria for treatment used in DANAMI3-PRIMULTI cannot be applied in this table.

As noted in Table 1.1, treatment of the nonculprit artery can occur at several different times after treatment of the culprit stenosis. Because Table 1.1 covers those scenarios

TABLE 1.4 NSTEMI/Unstable Angina

Indication	Appropriate Use Score (1-9)
Revascularization by PCI or CABG	
15. ■ Evidence of cardiogenic shock ■ Immediate revascularization of 1 or more coronary arteries	A (9)
16. ■ Patient stabilized ■ Intermediate- OR high-risk features for clinical events (e.g., TIMI score 3-4) ■ Revascularization of 1 or more coronary arteries	A (7)
17. ■ Patient stabilized after presentation ■ Low-risk features for clinical events (e.g., TIMI score ≤ 2) ■ Revascularization of 1 or more coronary arteries	M (5)

The number in parenthesis next to the rating reflects the median score for that indication.

A = appropriate; CABG = coronary artery bypass graft; M = may be appropriate; NSTEMI = non-ST-segment elevation myocardial infarction; PCI = percutaneous coronary intervention; R = rarely appropriate; TIMI = Thrombolysis In Myocardial Infarction.

where nonculprit treatment occurs immediately after the primary PCI, this table is specific for treatment of nonculprit stenoses after the initial procedure, but during the initial hospitalization.

Unstable angina/NSTEMI category—in patients with **Thrombolysis In Myocardial Infarction 3 flow** and multiple coronary artery stenoses, consideration should be given for heart team evaluation in patients with a high burden of CAD, such as 2-vessel disease with **proximal left anterior descending** coronary artery stenosis or more severe disease.


7. DISCUSSION

The new AUC ratings for ACS are consistent with existing guidelines for STEMI and NSTEMI-ACS (Figure 2). For patients with ACS, revascularization by either PCI or

CABG is the most commonly used therapy, and this is reflected in the ratings of “appropriate care” or “may be appropriate care” for all but 1 of the 17 scenarios presented. Although these AUC ratings do not compare the merits of PCI versus CABG for revascularization in ACS, in clinical practice, patients presenting with STEMI typically are treated by PCI of the culprit stenosis. However, the option of surgical revascularization should be considered for patients with ACS but less acute presentation, especially in those with complex multivessel CAD.

The current AUC rate revascularization as “appropriate care” for patients presenting within 12 hours of the onset of STEMI or up to 24 hours if there is clinical instability. For STEMI patients presenting more than 12 and up to 24 hours from symptom onset but with no signs of clinical instability, revascularization was rated as “may be appropriate,” indicating that many on the technical panel

FIGURE 2 Flow Diagram for the Determination of Appropriate Use in Patients With Acute Coronary Syndromes



Asx = asymptomatic; CABG = coronary artery bypass graft; FFR = fractional flow reserve; HF = heart failure; NSTEMI = non-ST-segment elevation myocardial infarction; PCI = percutaneous coronary intervention; STEMI = ST-segment elevation myocardial infarction; UA = unstable angina.

consider it reasonable to revascularize such patients. Furthermore, nonculprit artery revascularization at the time of primary PCI was rated as “may be appropriate,” but because this is an emerging concept on the basis of relatively small studies, clinical judgment by the operator is encouraged.

For STEMI patients initially treated with fibrinolysis, revascularization was rated as “appropriate therapy” in the setting of suspected failed fibrinolytic therapy or in stable and asymptomatic patients from 3 to 24 hours after fibrinolysis. In the setting of suspected failed fibrinolysis, the need for revascularization is usually immediate, whereas in stable patients with apparent successful fibrinolysis, revascularization can be delayed for up to 24 hours. For stable patients >24 hours after fibrinolysis, revascularization was rated as “may be appropriate.” Revascularization soon after apparent successful fibrinolysis is supported by data and guideline recommendations about the management of patients transferred from centers where PCI is not available.

Nonculprit artery revascularization during the index hospitalization after primary PCI or fibrinolysis was also rated as appropriate and reasonable for patients with 1 or more severe stenoses and spontaneous or easily provoked ischemia or for asymptomatic patients with ischemic findings on noninvasive testing. In the presence of an intermediate-severity nonculprit artery stenosis, revascularization was rated as “appropriate therapy” provided that the FFR was ≤ 0.80 . For patients who are stable and asymptomatic after primary PCI, revascularization was rated as “may be appropriate” for 1 or more severe stenoses even in the absence of further testing. The only “rarely appropriate” rating in patients with ACS occurred for asymptomatic patients with intermediate-severity nonculprit artery stenoses in the absence of any additional testing to demonstrate the functional significance of the stenosis.

For patients with NSTEMI/unstable angina, and consistent with existing guidelines and the available

evidence, revascularization was rated as “appropriate care” in the setting of cardiogenic shock or in a patient with intermediate- or high-risk features. For stable patients with low-risk features, revascularization was rated as “may be appropriate.” Decisions around the timing of revascularization, management of multivessel disease, and concomitant pharmacotherapy should all be on the basis of evidence from the relevant practice guidelines.

In conclusion, the AUC for ACS are consistent with the large body of evidence and guideline recommendations that support invasive strategies to define anatomy and revascularize patients with STEMI and NSTEMI-ACS. The evolving evidence around nonculprit stenosis revascularization has led to ratings that revascularization may be appropriate after primary PCI in selected asymptomatic patients with severe stenoses, defined herein as $\geq 70\%$ diameter narrowing, or in patients with intermediate-severity stenosis if FFR testing is abnormal. As in prior versions of the AUC, these revascularization ratings should be used to reinforce existing management strategies and identify patient populations that need more information to identify the most effective treatments.

ACC PRESIDENT AND STAFF

Richard A. Chazal, MD, FACC, President
Shalom Jacobovitz, Chief Executive Officer
William J. Oetgen, MD, FACC, Executive Vice President, Science, Education, and Quality
Joseph M. Allen, MA, Team Leader, Clinical Policy and Pathways
Leah White, MPH, CCRP, Team Leader, Appropriate Use Criteria
María Velásquez, Senior Research Specialist, Appropriate Use Criteria
Amelia Scholtz, PhD, Publications Manager, Clinical Policy and Pathways

REFERENCES

1. Patel MR, Dehmer GJ, Hirshfeld JW, Smith PK, Spertus JA. ACCF/SCAI/STS/AATS/AHA/ASNC/HFSA/SCCT 2012 appropriate use criteria for coronary revascularization focused update: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, Society of Thoracic Surgeons, American Association for Thoracic Surgery, American Heart Association, American Society of Nuclear Cardiology, and the Society of Cardiovascular Computed Tomography. *J Am Coll Cardiol*. 2012; 59:857-81.
2. O’Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61:e78-140.
3. Jneid H, Anderson JL, Wright RS, et al. 2012 ACCF/AHA focused update of the guideline for the management of patients with unstable angina/non-ST-elevation myocardial infarction (updating the 2007 guideline and replacing the 2011 focused update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2012;60:645-81.
4. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2014;64: 1929-49.
5. Levine GN, O’Gara PT, Bates ER, et al. 2015 ACC/AHA/SCAI focused update on primary percutaneous coronary intervention for patients with ST-elevation myocardial infarction: an update of the 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention and the 2013 ACCF/AHA Guideline for the Management of

ST-Elevation Myocardial Infarction: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *J Am Coll Cardiol*. 2016; 67:1235-50.

6. Engstrom T, Kelbaek H, Helqvist S, et al. Complete revascularisation versus treatment of the culprit lesion only in patients with ST-segment elevation myocardial infarction and multivessel disease (DANAMI-3-PRIMULTI): an open-label, randomised controlled trial. *Lancet*. 2015;386:665-71.

7. Gershlick AH, Khan JN, Kelly DJ, et al. Randomized trial of complete versus lesion-only revascularization in patients undergoing primary percutaneous coronary intervention for STEMI and multivessel disease: the CvLPRIT trial. *J Am Coll Cardiol*. 2015;65:963-72.

8. Wald DS, Morris JK, Wald NJ, et al. Randomized trial of preventive angioplasty in myocardial infarction. *N Engl J Med*. 2013;369:1115-23.

9. Bradley SM, Spertus JA, Kennedy KF, et al. Patient selection for diagnostic coronary angiography and hospital-level percutaneous coronary intervention appropriateness: insights from the National Cardiovascular Data Registry. *JAMA Intern Med*. 2014;174:1630-9.

10. Chan PS, Patel MR, Klein LW, et al. Appropriateness of percutaneous coronary intervention. *JAMA*. 2011; 306:53-61.

11. Desai NR, Bradley SM, Parzynski CS, et al. Appropriate use criteria for coronary revascularization and trends in utilization, patient selection, and appropriateness of percutaneous coronary intervention. *JAMA*. 2015;314:2045-53.

12. Hendel RC, Patel MR, Allen JM, et al. Appropriate use of cardiovascular technology: 2013 ACCF appropriate use criteria methodology update. *J Am Coll Cardiol*. 2013;61:1305-17.

13. Smith SC Jr, Allen J, Blair SN, et al. AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update endorsed by the National Heart, Lung, and Blood Institute. *J Am Coll Cardiol*. 2006;47: 2130-9.

14. Pearson TA, Blair SN, Daniels SR, et al. AHA guidelines for primary prevention of cardiovascular disease and stroke: 2002 update: Consensus Panel Guide to Comprehensive Risk Reduction for Adult Patients Without Coronary or Other Atherosclerotic Vascular Diseases. American Heart Association Science Advisory and Coordinating Committee. *Circulation*. 2002;106:388-91.

15. Buse JB, Ginsberg HN, Bakris GL, et al. Primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Circulation*. 2007;115:114-26.

16. Adult Treatment Panel III. Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002; 106:3143-421.

17. Chobanian AV, Bakris GL, Black HR, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42:1206-52.

18. Harold JG, Bass TA, Bashore TM, et al. ACCF/AHA/SCAI 2013 update of the clinical competence statement on coronary artery interventional procedures: a report of the American College of Cardiology Foundation/American Heart Association/American College of Physicians Task Force on Clinical Competence and Training (Writing Committee to Revise the 2007 Clinical Competence Statement on Cardiac Interventional Procedures). *J Am Coll Cardiol*. 2013; 62:357-96.

19. Hillis LD, Smith P, Anderson J, et al. 2011 ACCF/AHA guidelines for coronary artery bypass graft surgery: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58:e123-210.

20. Levine GN, Bates ER, Blankenship JC, et al. 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *J Am Coll Cardiol*. 2011;58:e44-122.

21. Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;64: e139-228.

22. TIMI Study Group. Available at: <http://www.timi.org>. Accessed March 15, 2011.

23. Chewning B, Bylund CL, Shah B, Arora NK, Gueguen JA, Makoul G. Patient preferences for shared decisions: a systematic review. *Patient Educ Couns*. 2012;86:9-18.

24. Lin GA, Fagerlin A. Shared decision making: state of the science. *Circ Cardiovasc Qual Outcomes*. 2014;7: 328-34.

25. Ting HH, Brito JP, Montori VM. Shared decision making: science and action. *Circ Cardiovasc Qual Outcomes*. 2014;7:323-7.

26. Buchanan A. Medical paternalism. *Philos Public Aff*. 1978;7:370-90.

KEY WORDS ACC Appropriate Use Criteria, coronary revascularization, imaging, medical therapy, multimodality

APPENDIX A. APPROPRIATE USE CRITERIA FOR CORONARY REVASCULARIZATION IN PATIENTS WITH ACUTE CORONARY SYNDROMES: PARTICIPANTS

Writing Group

Manesh R. Patel, MD, FACC, FAHA, FSCAI—Associate Professor of Medicine, Director Interventional Cardiology and Catheterization Labs, Duke University Health System, Duke Clinical Research Institute, Durham, NC

John H. Calhoun, MD—Professor and Chair, Presidents Council Chair for Excellence in Surgery, Department of Cardiothoracic Surgery, Director, University of Texas Health Science Center at San Antonio, Heart and Vascular Institute, San Antonio, TX

Gregory J. Dehmer, MD, MACC, MSCAI, FACP, FAHA—Clinical Professor of Medicine, Texas A&M Health Science Center College of Medicine, Medical Director, Cardiovascular Services, Central Texas Division, Director, Cardiology Division, Baylor Scott & White—Temple Memorial, Temple, TX

James Aaron Grantham, MD, FACC—Associate Clinical Professor, University of Missouri-Kansas City School of Medicine, Director, Cardiovascular Disease Fellowship Program, University of Missouri-Kansas City School of Medicine, Director, Cardiovascular Medical Education, Saint Luke's Hospital, Kansas City, MO

Thomas M. Maddox, MD, MSc, FACC, FAHA—National Director, VA CART Program Cardiology, VA Eastern Colorado Health Care System, Associate Professor, Department of Medicine, Cardiology, University of Colorado, Colorado Cardiovascular Outcomes Research Consortium, Denver, CO

David J. Maron, MD, FACC, FAHA—Clinical Professor of Medicine, Cardiovascular, Director, Preventive Cardiology, ISCHEMIA Trial Co-Chair, Principal Investigator, Stanford University School of Medicine, Stanford, CA

Peter K. Smith, MD, FACC—Professor of Surgery, Division Chief, Cardiovascular and Thoracic Surgery, Duke University, Durham, NC

Rating Panel

Michael J. Wolk, MD, MACC, Moderator—Past President, American College of Cardiology, Clinical Professor of Medicine, Weill Medical College of Cornell University, New York Cardiology Associates, New York, NY

Manesh R. Patel, MD, FACC, FAHA, FSCAI, Writing Committee Liaison—Associate Professor of Medicine, Director Interventional Cardiology and Catheterization Labs, Duke University Health System, Duke Clinical Research Institute, Durham, NC

Gregory J. Dehmer, MD, MACC, MSCAI, FAHA, Writing Committee Liaison—Clinical Professor of Medicine, Texas

A&M Health Science Center College of Medicine, Medical Director, Cardiovascular Services, Central Texas Division, Director, Cardiology Division, Baylor Scott and White, Temple Memorial, Temple, TX

Peter K. Smith, MD, FACC, Writing Committee Liaison—Professor of Surgery, Division Chief, Cardiovascular and Thoracic Surgery, Duke University, Durham, NC

James C. Blankenship, MD, MACC—Staff Physician, Director, Cardiac Catheterization Laboratory, Division of Cardiology, Geisinger Medical Center, Danville, PA

Alfred A. Bove, MD, PhD, MACC—Past President, American College of Cardiology, Professor Emeritus, Lewis Katz School of Medicine, Heart and Vascular, Temple University, Philadelphia, PA

Steven M. Bradley, MD—Staff Cardiologist, VA Eastern Colorado Health Care System, Assistant Professor of Medicine, Division of Cardiology at the University of Colorado, Denver, CO

Larry S. Dean, MD, FACC, FSCAI—Professor of Medicine and Surgery, University of Washington School of Medicine, Director, University of Washington, Medicine Regional Heart Center, Seattle, WA

Peter L. Duffy, MD, FACC, FSCAI—Director of Quality for the Cardiovascular Service Line, First Health of the Carolinas, Reid Heart Institute/Moore Regional Hospital, Pinehurst, NC

T. Bruce Ferguson, Jr., MD, FACC—Professor of Thoracic Surgery, Department of Cardiovascular Sciences, Cardiothoracic Surgery, East Carolina Heart Institute, East Carolina University, Greenville, NC

Frederick L. Grover, MD, FACC—Professor of Cardiothoracic Surgery, Department of Cardiothoracic Surgery, University of Colorado, Denver, CO

Robert A. Guyton, MD, FACC—Chief of Cardiothoracic Surgery, Professor of Surgery, Division of Cardiothoracic Surgery, Department of Surgery, Director, Thoracic Surgery Residency Program, Emory University School of Medicine, Atlanta, GA

Mark A. Hlatky, MD, FACC—Professor of Health Research and Policy, Health Services Research, Professor of Medicine, Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA

Harold L. Lazar, MD, FACC—Director, Cardiothoracic Research Program, Professor of Cardiothoracic Surgery, Boston University School of Medicine, Boston, MA

Vera H. Rigolin, MD, FACC—Professor, Cardiology, Northwestern University Feinberg School of Medicine, Chicago, IL

Geoffrey A. Rose, MD, FACC, FASE—Chief, Division of Cardiology, Sanger Heart and Vascular Institute, Charlotte, NC

Richard J. Shemin, MD, FACC—Robert and Kelly Day Professor, Chief of Cardiothoracic Surgery, Executive Vice Chair of Surgery, Co-Director of the Cardiovascular Center, Director of Cardiac Quality at the Ronald Reagan UCLA Medical Center, Los Angeles, CA

Jacqueline E. Tamis-Holland, MD, FACC—Director, Interventional Cardiology Fellowship, Mount Sinai, Saint Luke's Hospital Director, Women's Heart NY Assistant Professor of Medicine, Icahn School of Medicine at Mount Sinai Hospital, New York, NY

Carl L. Tommaso, MD, FACC, FSCAI—Director of the Cardiac Catheterization Laboratory at Skokie Illinois Hospital, part of the Northshore University Health System, Associate Professor of Medicine at Rush Medical College in Chicago, Chicago, IL

L. Samuel Wann, MD, MACC—Past President, American College of Cardiology, Clinical Cardiologist, Columbia St. Mary's Healthcare, Medical Director, Heart Failure Program, Milwaukee, WI

John B. Wong, MD—Chief, Division of Clinical Decision Making, Primary Care Physician, Principal Investigator, Institute for Clinical Research and Health Policy Studies, Professor, Tufts University School of Medicine, Boston, MA

Reviewers

Jeffrey L. Anderson, MD, FACC—Associate Chief of Cardiology, Intermountain Medical Center, Murray, UT

James C. Blankenship, MD, MACC—Staff Physician, Director, Cardiac Catheterization Laboratory, Geisinger Medical Center, Division of Cardiology, Danville, PA

Jeffrey A. Brinker, MD, FACC—Professor of Medicine, Johns Hopkins Hospital, Baltimore, MD

Alexandru I. Costea, MD—Associate Professor, University of Cincinnati Medical Center, Cincinnati, OH

Ali E. Denktas, MD, FACC—Assistant Professor, Baylor College of Medicine, Houston, TX

Lloyd W. Klein, MD, FACC—Professor of Medicine, Melrose Park, IL

Frederick G. Kushner, MD, FACC—Clinical Professor, Tulane University Medical Center, Medical Director, Heart Clinic of Louisiana, Marrero, LA

Glenn N. Levine, MD, FACC—Professor, Baylor College of Medicine, Cardiology, Pearland, TX

David Joel Maron, MD, FACC—Professor of Medicine and Emergency Medicine, Stanford University School of Medicine, Stanford, CA

James B. McClurken, MD, FACC—Director of Thoracic Surgery, Professor of Surgery Emeritus, Temple University, School of Medicine, Richard A Reif Heart Institute, Doylestown Hospital, Doylestown, PA

Robert N. Piana, MD, FACC—Professor of Medicine, Cardiology, Vanderbilt University Medical Center, Nashville, TN

John A. Spertus, M.D, MPH, FACC—Adjunct Professor of Medicine, Washington University School of Medicine, St. Louis, MO

Raymond F. Stainback, MD, FACC—Medical Director, Non-Invasive Cardiology Texas Heart Institute at Baylor St. Luke's Medical Center, Houston, TX

Robert C. Stoler, MD, FACC—Director of Cardiac Catheterization Laboratory, Cardiology Consultants of Texas, Dallas, TX

Todd C. Villines, MD, FACC—Co-Director of Cardiovascular Computed Tomography and Assistant Chief, Cardiology Service at Walter Reed Army Medical Center, Rockville, MD

David H. Wiener, MD, FACC—Professor of Medicine, Jefferson Medical College, Jefferson Heart Institute, Philadelphia, PA

ACC Appropriate Use Criteria Task Force

John U. Doherty, MD, FACC, FAHA—Co-Chair, AUC Task Force, Professor of Medicine, Jefferson Medical College of Thomas Jefferson University, Philadelphia, PA

Gregory J. Dehmer, MD, MACC—Co-Chair, AUC Task Force, Medical Director, Cardiovascular Services, Central Texas Division, Baylor Scott & White Health, Temple, TX

Steven R. Bailey, MD, FACC, FSCAI, FAHA—Chair, Division of Cardiology, Professor of Medicine and Radiology, Janey Briscoe Distinguished Chair, University of Texas Health Sciences Center, San Antonio, TX

Nicole M. Bhave, MD, FACC—Clinical Assistant Professor, Department of Internal Medicine, Division of Cardiovascular Medicine, University of Michigan Cardiovascular Center, Ann Arbor, MI

Alan S. Brown, MD, FACC—Medical Director, Midwest Heart Disease Prevention Center, Advocate Lutheran General Hospital, Director, Division of Cardiology, Park Ridge, IL

Stacie L. Daugherty, MD, FACC—Associate Professor, Division of Cardiology, Department of Medicine, University of Colorado School of Medicine, Denver, CO

Milind Y. Desai, MBBS, FACC—Associate Director, Clinical Investigations Heart and Vascular Institute, Cleveland Clinic, Cleveland, OH

Claire S. Duvernoy, MD, FACC—Cardiology Section Chief, Division of Cardiology, University of Michigan Health System, Ann Arbor, MI

Linda D. Gillam, MD, FACC—Chair, Department of Cardiovascular Medicine, Morristown Medical Center, Morristown, NJ

Robert C. Hendel, MD, FACC, FAHA—Director of Cardiac Imaging and Outpatient Services, Division of Cardiology, Miami University School of Medicine, Miami, FL

Christopher M. Kramer, MD, FACC, FAHA—Former Co-Chair, AUC Task Force, Ruth C. Heede Professor of Cardiology & Radiology, and Director, Cardiovascular Imaging Center, University of Virginia Health System, Charlottesville, VA

Bruce D. Lindsay, MD, FACC—Professor of Cardiology, Cleveland Clinic Foundation of Cardiovascular Medicine, Cleveland, OH

Warren J. Manning, MD, FACC—Professor of Medicine and Radiology, Beth Israel Deaconess Medical Center, Division of Cardiology, Boston, MA

Manesh R. Patel, MD, FACC, FAHA—Former Chair, AUC Task Force, Assistant Professor of Medicine, Division

of Cardiology, Duke University Medical Center, Durham, NC

Ritu Sachdeva, MBBS, FACC—Associate Professor, Division of Pediatric Cardiology, Department of Pediatrics, Emory University School of Medicine, Children's Health Care of Atlanta, Sibley Heart Center Cardiology, Atlanta, GA

L. Samuel Wann, MD, MACC—Staff Cardiologist, Columbia St. Mary's Healthcare, Milwaukee, WI

David E. Winchester, MD, FACC—Assistant Professor of Medicine, University of Florida, Division of Cardiology, Gainesville, FL

Joseph M. Allen, MA—Team Leader, Clinical Policy and Pathways, American College of Cardiology, Washington, DC

APPENDIX B. RELATIONSHIPS WITH INDUSTRY (RWI) AND OTHER ENTITIES

The College and its partnering organizations rigorously avoid any actual, perceived, or potential conflicts of interest that might arise as a result of an outside relationship or personal interest of a member of the rating panel. Specifically, all panelists are asked to provide disclosure statements of all relationships that might be perceived as real or potential conflicts of interest. These statements were reviewed by the Appropriate Use Criteria Task Force,

discussed with all members of the rating panel at the face-to-face meeting, and updated and reviewed as necessary. The following is a table of relevant disclosures by the rating panel and oversight working group members. In addition, to ensure complete transparency, a full list of disclosure information—including relationships not pertinent to this document—is available in the [Online Appendix](#).

APPROPRIATE USE CRITERIA FOR CORONARY REVASCULARIZATION IN PATIENTS WITH ACUTE CORONARY SYNDROMES: MEMBERS OF THE WRITING GROUP, RATING PANEL, INDICATION REVIEWERS, AND AUC TASK FORCE—RELATIONSHIPS WITH INDUSTRY AND OTHER ENTITIES (RELEVANT)

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Writing Group							
Manesh R. Patel (Chair)	Duke University Health System, Duke Clinical Research Institute—Associate Professor of Medicine, Director Interventional Cardiology and Catheterization Labs	None	None	None	None	None	None
John H. Calhoon	University of Texas Health Science Center at San Antonio Department of Cardiothoracic Surgery, Heart and Vascular Institute Director—Professor and Chair, Presidents Council Chair for Excellence in Surgery	None	None	None	None	None	None
Gregory J. Dehmer	Baylor Scott & White-Temple Memorial, Texas A&M Health Science Center College of Medicine, Central Texas Division—Clinical Professor of Medicine, Medical Director, Cardiovascular Services, Director, Cardiology Division	None	None	None	None	None	None
James Aaron Grantham	Saint Luke's Hospital—Associate Clinical Professor, University of Missouri-Kansas City School of Medicine—Director, Cardiovascular Disease Fellowship Program, Director, Cardiovascular Medical Education	<ul style="list-style-type: none"> ■ Abbott Vascular† ■ Asahi-Intecc† ■ Boston Scientific† ■ Bridgepoint Medical Systems† ■ Medtronic† 	None	None	<ul style="list-style-type: none"> ■ Abbott Vascular† ■ Asahi-Intecc† ■ Boston Scientific† ■ Bridgepoint Medical Systems† ■ Medtronic† 	None	None
Thomas M. Maddox	VA Eastern Colorado Health Care System—National Director, Associate Professor, Department of Medicine, Cardiology, University of Colorado, Colorado Cardiovascular Outcomes Research Consortium	None	None	None	None	None	None

Continued on the next page

APPENDIX B. CONTINUED

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
David J. Maron	Stanford University School of Medicine—Clinical Professor of Medicine, Cardiovascular, Director, Preventive Cardiology	None	None	None	None	None	None
Peter K. Smith	Cardiovascular and Thoracic Surgery, Duke University—Professor of Surgery, Division Chief	None	None	None	None	None	None
Rating Panel							
James C. Blankenship	Geisinger Medical Center, Division of Cardiology—Staff Physician, Director, Cardiac Catheterization Laboratory	None	None	None	<ul style="list-style-type: none"> ■ Abbott Vascular* ■ AstraZeneca* ■ Boston Scientific* ■ GlaxoSmithKline* ■ Hamilton Health Services* ■ Medinol LTD* ■ Orexigen Therapeutics/Takeda* ■ Stentys, Inc.* ■ Takeda Pharmaceuticals 	None	None
Alfred A. Bove	Temple University, Lewis Katz School of Medicine, Heart and Vascular—Professor Emeritus	None	None	None	<ul style="list-style-type: none"> ■ Merck Schering-Plough† 	None	None
Steven M. Bradley	VA Eastern Colorado Health Care System, Division of Cardiology at the University of Colorado—Staff Cardiologist, Assistant Professor of Medicine	None	None	None	None	None	None
Larry S. Dean	Medicine Regional Heart Center University of Washington School of Medicine—Professor of Medicine and Surgery, Director	<ul style="list-style-type: none"> ■ Philips Medical† 	None	None	<ul style="list-style-type: none"> ■ Edwards Lifesciences† 	None	None
Peter L. Duffy	First Health of the Carolinas, Reid Heart Institute/Moore Regional Hospital—Director of Quality for the Cardiovascular Service Line	None	<ul style="list-style-type: none"> ■ Volcano Corp† 	None	None	None	None
T. Bruce Ferguson, Jr.	East Carolina Heart Institute, East Carolina University, Department of Cardiovascular Sciences, Cardiothoracic Surgery—Professor of Thoracic Surgery	None	None	<ul style="list-style-type: none"> ■ RFPI* 	<ul style="list-style-type: none"> ■ Novadaq Technologies† 	None	None
Frederick L. Grover	University of Colorado, Department of Cardiothoracic Surgery—Professor of Cardiothoracic Surgery	<ul style="list-style-type: none"> ■ Somalution 	None	None	None	None	None
Robert A. Guyton	Emory University School of Medicine, Division of Cardiothoracic Surgery, Department of Surgery, Thoracic Surgery Residency Program—Chief of Cardiothoracic Surgery, Professor of Surgery, Director	<ul style="list-style-type: none"> ■ Medtronic† 	None	None	None	None	None

Continued on the next page

APPENDIX B. CONTINUED

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Mark A. Hlatky	Stanford University School of Medicine, Cardiovascular Medicine, Health Services Research—Professor of Health Research and Policy, Professor of Medicine	None	None	None		■ Sanofi-Aventis	None
Harold L. Lazar	Boston University School of Medicine, Cardiothoracic Research Program—Director Professor of Cardiothoracic Surgery	None	None	None	None	None	None
Vera H. Rigolin	Northwestern University Feinberg School of Medicine, Cardiology—Professor	None	None	None	None	■ Pfizer†	None
Geoffrey A. Rose	Division of Cardiology, Sanger Heart and Vascular Institute—Chief	None	None	None	None	■ Medtronic	None
Richard J. Shemin	Ronald Reagan UCLA Medical Center, Cardiovascular Center—Director of Cardiac Quality, Robert and Kelly Day Professor, Chief of Cardiothoracic Surgery, Executive Vice Chair of Surgery	■ Edwards Lifesciences ■ Sorin Group	None	None	None	None	None
Jacqueline E. Tamis-Holland	Saint Luke's Hospital, Icahn School of Medicine at Mount Sinai Hospital Mount Sinai—Director, Women's Heart NY, Assistant Professor of Medicine, Director, Interventional Cardiology Fellowship	None	None	None	None	None	None
Carl L. Tommaso	Rush Medical College in Chicago, Skokie Illinois Hospital, part of the Northshore University Health System—Director of the Cardiac Catheterization Laboratory, Associate Professor of Medicine	None	None	None	None	None	None
L. Samuel Wann	Columbia St. Mary's Healthcare—Clinical Cardiologist, Medical Director, Heart Failure Program	■ United Healthcare	None	None	None	None	None
John B. Wong	Tufts University School of Medicine—Chief, Division of Clinical Decision Making, Primary Care Physician, Principal Investigator, Institute for Clinical Research and Health Policy Studies, Professor	None	None	None	None	None	None

Continued on the next page

APPENDIX B. CONTINUED

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Reviewers							
Jeffrey L. Anderson	Intermountain Medical Center—Associate Chief of Cardiology	■ Sanofi-Aventis ■ The Medicines Company	None	None	None	None	None
Jeffrey A. Brinker	Johns Hopkins Hospital—Professor of Medicine	None	None	None	None	None	None
Alexandru I. Costea	University of Cincinnati Medical Center—Associate Professor	None	None	None	None	■ Boston Scientific*	None
Ali E. Denktas	Baylor College of Medicine—Assistant Professor	None	None	None	■ AstraZeneca ■ Edwards Lifesciences	None	None
Lloyd W. Klein	Melrose Park—Professor of Medicine	None	None	None	None	None	None
Frederick G. Kushner	Tulane University Medical Center, Heart Clinic of Louisiana—Clinical Professor, Medical Director	None	None	None	None	None	None
Glenn N. Levine	Baylor College of Medicine, Cardiology—Professor	None	None	None	None	None	None
David J. Maron	Stanford University School of Medicine—Professor of Medicine and Emergency Medicine	None	None	None	None	None	None
James B. McClurken	Temple University, School of Medicine, Richard A Reif Heart Institute, Doylestown Hospital—Director of Thoracic Surgery, Professor of Surgery Emeritus	None	None	None	None	None	None
Robert N. Piana	Vanderbilt University Medical Center—Professor of Medicine, Cardiology	■ Axio Research ■ Harvard Clinical Research Institute ■ W.L. Gore & Associates, Inc.	None	None	None	None	None
John A. Spertus	Washington University School of Medicine—Adjunct Professor of Medicine	■ Amgen ■ Bayer Health-care Pharmaceuticals ■ Janssen ■ Novartis ■ Regeneron	None	■ Health Outcomes Sciences	None	None	None
Raymond F. Stainback	Texas Heart Institute at Baylor St. Luke's Medical Center, Non-Invasive Cardiology—Medical Director	None	None	None	None	None	None
Robert C. Stoler	Cardiology Consultants of Texas—Director of Cardiac Catheterization Laboratory	■ Boston Scientific ■ Medtronic	None	None	None	None	None

Continued on the next page

APPENDIX B. CONTINUED

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Todd C. Villines	Cardiology Service at Walter Reed Army Medical Center—Co-Director of Cardiovascular Computed Tomography and Assistant Chief	■ Boehringer Ingelheim†	None	None	None	None	None
David H. Wiener	Jefferson Medical College, Jefferson Heart Institute—Professor of Medicine	None	None	None	None	None	None
Appropriate Use Criteria Task Force							
Steven R. Bailey	University of Texas Health Sciences Center—Chair, Division of Cardiology, Professor of Medicine and Radiology, Janey Briscoe Distinguished Chair	None	None	None	None	None	None
Nicole M. Bhave	University of Michigan Cardiovascular Center, Department of Internal Medicine, Division of Cardiovascular Medicine—Clinical Assistant Professor	None	None	None	None	None	None
Alan S. Brown	Midwest Heart Disease Prevention Center, Advocate Lutheran General Hospital—Director, Division of Cardiology—Medical Director	None	None	None	None	None	None
Stacie L. Daugherty	University of Colorado School of Medicine, Division of Cardiology, Department of Medicine—Associate Professor	None	None	None	None	None	None
Gregory J. Dehmer	Baylor Scott & White, Central Texas Division, Cardiovascular Services Health—Medical Director	None	None	None	None	None	None
Milind Y. Desai	Cleveland Clinic, Clinical Investigations, Heart and Vascular Institute—Associate Director	None	None	None	None	None	None
John U. Doherty	Thomas Jefferson University, Jefferson Medical College—Professor of Medicine	None	None	None	None	None	None
Claire S. Duvernoy	University of Michigan Health System, Division of Cardiology—Cardiology Section Chief	None	None	None	None	None	None
Linda D. Gillam	Morristown Medical Center, Department of Cardiovascular Medicine—Chair	■ Edwards Lifesciences* ■ Medtronic*	None	None	None	None	None
Robert C. Hendel	Miami University School of Medicine, Division of Cardiology—Director of Cardiac Imaging and Outpatient Services	None	None	None	None	None	None

Continued on the next page

APPENDIX B. CONTINUED

Participant	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Christopher M. Kramer	University of Virginia Health System—Ruth C. Heede Professor of Cardiology & Radiology, Director, Cardiovascular Imaging Center	None	None	None	None	None	None
Bruce D. Lindsay	Cleveland Clinic Foundation of Cardiovascular Medicine—Professor of Cardiology	None	None	None	None	None	None
Warren J. Manning	Beth Israel Deaconess Medical Center, Division of Cardiology—Professor of Medicine and Radiology	■ Merck	None	None	■ Philips Medical Systems†	None	None
Manesh R. Patel	Duke University Medical Center, Division of Cardiology—Assistant Professor of Medicine	None	None	None	None	None	None
Ritu Sachdeva	Emory University School of Medicine, Children's Health Care of Atlanta, Sibley Heart Center Cardiology, Division of Pediatric Cardiology, Department of Pediatrics—Associate Professor	None	None	None	None	None	None
L. Samuel Wann	Columbia St. Mary's Healthcare—Staff Cardiologist	None	None	None	None	None	None
David E. Winchester	University of Florida, Division of Cardiology—Assistant Professor of Medicine	None	None	None	None	None	None
Joseph M. Allen	American College of Cardiology—Team Leader, Clinical Policy and Pathways	None	None	None	None	None	None

Note: A standard exemption to the ACC relationship with industry policy is extended to AUC writing groups, because they do not make recommendations but rather prepare background materials and typical clinical scenarios/indications that are rated independently by a separate panel of experts. This table represents relevant relationships of participants with industry and other entities that were reported by reviewers at the time this document was under development. The table does not necessarily reflect relationships with industry at the time of publication. A person is deemed to have a significant interest in a business if the interest represents ownership of $\geq 5\%$ of the voting stock or share of the business entity, or ownership of $\geq \$5,000$ of the fair market value of the business entity; or if funds received by the person from the business entity exceed 5% of the person's gross income for the previous year. Relationships that exist with no financial benefit are also included for the purpose of transparency. Relationships in this table are modest unless otherwise noted. Please refer to <http://www.acc.org/guidelines/about-guidelines-and-clinical-documents/relationships-with-industry-policy> for definitions of disclosure categories or additional information about the ACC Disclosure Policy for Writing Committees.

*No financial benefit.

†Significant relationship.

ACC = American College of Cardiology; AUC = appropriate use criteria.