

Non-occlusive mesenteric ischemia (NOT) to be missed

Christiaan Boerma




mcl
medisch centrum
leeuwarden


What is NOMI?


Christiaan Boerma





mcl
medisch centrum
leeuwarden














Knichwitz, Der Anaesthetist 2005; 54:41-48










Duplex Ultrasound Measurement of Postprandial Intestinal Blood Flow: Effect of Meal Composition

GASTROENTEROLOGY 1988;95:1294-301





Exercise-Induced Splanchnic Hypoperfusion Results in Gut Dysfunction in Healthy Men

Kim van Wijck^{1,2,3}, Kaatje Lenaerts^{1,2,3*}, Luc J. C. van Loon^{1,2,4}, Wilbert H. M. Peters⁵, Wim A. Buurman^{2,3}, Cornelis H. C. Dejong^{1,2,3}





Exercise-Induced Splanchnic Hypoperfusion Results in Gut Dysfunction in Healthy Men

Kim van Wijck^{1,2,3}, Kaatje Lenaerts^{1,2,3*}, Luc J. C. van Loon^{1,2,4}, Wilbert H. M. Peters⁵, Wim A. Buurman^{2,3}, Cornelis H. C. Dejong^{1,2,3}






Exercise-Induced Splanchnic Hypoperfusion Results in Gut Dysfunction in Healthy Men

Kim van Wijck^{1,2,3}, Kaatje Lenaerts^{1,2,3*}, Luc J. C. van Loon^{1,2,4}, Wilbert H. M. Peters⁵, Wim A. Buurman^{2,3}, Cornelis H. C. Dejong^{1,2,3}




Exercise-Induced Splanchnic Hypoperfusion Results in Gut Dysfunction in Healthy Men

Kim van Wijck^{1,2,3}, Kaatje Lenaerts^{1,2,3*}, Luc J. C. van Loon^{1,2,4}, Wilbert H. M. Peters⁵, Wim A. Buurman^{2,3}, Cornelis H. C. Dejong^{1,2,3}





- 
14. Schaub N, Spichtin P, Stalder GA. Ischemic colitis as a cause of intestinal bleeding after marathon running. *Schweiz Med Wochenschr* 1985;115:454–7.
15. Heer M, Repond F, Hany A, Sulser H, Jager K. Acute ischaemic colitis in a female long distance runner. *Gut* 1987;28:896–9.
16. Moses F, Brewer TG, Peura DA. Running associated proximal hemorrhagic colitis. *Ann Intern Med* 1988;108:385–6.
17. Beaumont AC, Teare JP. Subtotal colectomy following marathon running in a female patient. *J R Soc Med* 1991;84:439–40.
18. Schwartz AE, Vanagunas A, Kamel PL. Endoscopy to evaluate gastrointestinal bleeding in marathon runners. *Ann Intern Med* 1990;113:632–3.
19. Clausen JP. Effect of physical training on cardiovascular adjustments to exercise in man. *Physiol Rev* 1997;77:779–815.
20. Kehl O, Jager K, Munch R, et al. Mesenteric ischemia as a cause of jogging anemia? *Schweiz Med Wochenschr* 1986;116:974–6.
21. Lucas W, Schroy PC. Reversible ischemic colitis in a high endurance athlete. *Am J Gastroenterol* 1998;93:2231–4.



Hemodynamics of Pancreatic Ischemia in Cardiogenic Shock in Pigs

GASTROENTEROLOGY 1997;113:938–945

PATRICK M. REILLY, THOMAS J. TOUNG, MASAHIKO MIYACHI, HENRY J. SCHILLER,
and GREGORY B. BULKLEY

- Total vascular resistance
- Pancreatic vascular resistance

Shock




Hemodynamics of Pancreatic Ischemia in Cardiogenic Shock in Pigs


GASTROENTEROLOGY 1997;113:938-945

PATRICK M. REILLY, THOMAS J. TOUNG, MASAHIKO MIYACHI, HENRY J. SCHILLER,
and GREGORY B. BULKLEY

- Total vascular resistance
- Pancreatic vascular resistance



Angiotensin II








Nonocclusive Mesenteric Hypoperfusion Syndromes: Recognition and Treatment

Martin Björck and Anders Wanhainen, Semin Vasc Surg 23:54-64 2010





	R	L	p Value
<i>Symptoms</i>			
Abdominal pain	78.6%	71.9%	0.266
Rectal bleeding	44.9%	87.4%	<0.001
Diarrhea	55.3%	45.5%	0.160
Nausea	57.6%	39.4%	0.013
Weight loss	55.6%	18.8%	<0.001
<i>Physical examination</i>			
Temperature $\geq 38^\circ$	22.3%	15.2%	0.178
Hypotension (<100 mmHg systolic pressure)	12.5%	15.9%	0.481
Tachycardia (>100/min)	33.0%	25.0%	0.207
Paralytic ileus	31.5%	17.9%	0.018
Peritoneal signs	27.1%	6.7%	<0.001

	R	L	p Value
<i>Symptoms</i>			
Abdominal pain	78.6%	71.9%	0.266
Rectal bleeding	44.9%	87.4%	<0.001
Diarrhea	55.3%	45.5%	0.160
Nausea	57.6%	39.4%	0.013
Weight loss	55.6%	18.8%	<0.001
<i>Physical examination</i>			
Temperature $\geq 38^\circ$	22.3%	15.2%	0.178
Hypotension (<100 mmHg systolic pressure)	12.5%	15.9%	0.481
Tachycardia (>100/min)	33.0%	25.0%	0.207
Paralytic ileus	31.5%	17.9%	0.018
Peritoneal signs	27.1%	6.7%	<0.001



Outcome of acute mesenteric ischemia in the intensive care unit: a retrospective, multicenter study of 780 cases


Intensive Care Med (2015) 41:667–676




Nonocclusive Mesenteric Hypoperfusion Syndromes: Recognition and Treatment

Seminars in
**VASCULAR
SURGERY**

Martin Björck and Anders Wanhainen, Semin Vasc Surg 23:54-64 2010









FUJINON

17/11/2005
16:31:11

PEAK

FILM 03
1/100







D

E

F




C




Markus Trompeter
Thurid Brazda
Christopher T. Remy
Thomas Vestring
Peter Reimer

Non-occlusive mesenteric ischemia: etiology, diagnosis, and interventional therapy



a



b








Gut ischaemia

U Haglund


Gut 1994; supplement 1: S73–S76







Inspiration






Expiration

CA

SMA



IMA




CA

SMA


IMA




CA



SMA






IMA



CA

SMA

IMA



DON'T rule out



Clinical symptoms

Lactate

Endoscopy/laparoscopy

Christiaan Boerma



mcl
medisch centrum
leeuwarden

Avoid



Vasopressors/NSAIDS



Intra abdominal hypertension



Enteral feeding

Christiaan Boerma



mcl
medisch centrum
leeuwarden

DO

-  History
-  Biopsy
-  Hybrid

Christiaan Boerma



mcl
medisch centrum
leeuwarden